

# MUCINEX® GOLD & FLU FAST-RELEASE LIQUID GELS FOR THOSE WITH HBP AND DIABETES

## Drug Facts

### Active ingredients (in each liquid gel)

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Guaifenesin 200 mg.....Expectorant

### Purposes

### Uses

- temporarily relieves these common cold and flu symptoms:
  - minor aches and pains
  - headache      ■ sore throat
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 liquid gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening      ■ blisters      ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### When using this product do not use more than directed

#### Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

## Drug Facts (continued)

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 liquid gels in any 24-hour period
- adults and children 12 years of age and over: take 2 liquid gels every 4 hours
- children under 12 years of age: do not use

### Other information

- store at 20-25°C (68-77°F)
- avoid excessive heat

### Inactive ingredients

FD&C blue no. 1, gelatin, glycerin, hypromellose, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

### Questions? 1-866-MUCINEX (1-866-682-4639)