

### Drug Facts (continued)

- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

#### Mucinex Sinus-Max Day Pressure, Pain & Cough

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

#### Mucinex Nightshift Night Sinus

- do not take more than directed (see Overdose warning)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

**Other information** ■ store at 20-25°C (68-77°F)

### Inactive ingredients

#### Mucinex Sinus-Max Day Pressure, Pain & Cough

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

### Inactive ingredients

#### Mucinex Nightshift Night Sinus

croscarmellose sodium, crospovidone, ferric oxide, hydroxypropyl cellulose, mica, microcrystalline cellulose, polyvinyl alcohol, polyvinyl alcohol polyethylene glycol copolymer, povidone, silicon dioxide, stearic acid, talc, titanium dioxide

**Questions?** 1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

### Drug Facts (continued)

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease ■ heart disease ■ diabetes
- high blood pressure ■ thyroid disease
- glaucoma (*Nightshift Night Sinus only*)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (*Nightshift Night Sinus only*)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

- cough that occurs with too much phlegm (mucus)

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (*Nightshift Night Sinus only*)

#### When using this product

- do not use more than directed
- excitability may occur, especially in children (*Nightshift Night Sinus only*)
- marked drowsiness may occur (*Nightshift Night Sinus only*)
- alcohol, sedatives, and tranquilizers may increase drowsiness (*Nightshift Night Sinus only*)
- avoid alcoholic drinks (*Nightshift Night Sinus only*)
- use caution when driving a motor vehicle or operating machinery (*Nightshift Night Sinus only*)

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

### Drug Facts

#### Active ingredients (in each caplet)

#### Mucinex Sinus-Max Day Pressure, Pain & Cough

Acetaminophen 325 mg .....	Pain reliever
Dextromethorphan HBr 10 mg .....	Cough suppressant
Guaifenesin 200 mg .....	Expectorant
Phenylephrine HCl 5 mg .....	Nasal decongestant

#### Purposes

#### Active ingredients (in each caplet)

#### Mucinex Nightshift Night Sinus

Acetaminophen 325 mg .....	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg .....	Cough suppressant
Phenylephrine HCl 5 mg .....	Nasal decongestant
Triprolidine HCl 1.25 mg .....	Antihistamine

#### Purposes

### Uses

#### Mucinex Sinus-Max Day Pressure, Pain & Cough

- temporarily relieves:
  - nasal congestion ■ headache
  - minor aches and pains ■ cough
  - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

#### Mucinex Nightshift Night Sinus

- temporarily relieves these common cold and flu symptoms:
  - cough ■ nasal congestion ■ minor aches and pains
  - sore throat ■ headache ■ sneezing
  - sinus congestion and pressure ■ runny nose
  - itching of the nose or throat
  - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

### Warnings

**Liver warning:** This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning (*Nightshift Night Sinus only*):** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.