



DIRECT TO RESERVOIR

Mixing Instructions and Feed Charts



More Info

METRIC - mL/L and g/L


1. Fill reservoir to target volume; begin agitation.
2. Add **Front Row Si***; agitate 3-5 minutes.
3. Add **Part A**; agitate 3-5 minutes.
4. Add **Part B**; agitate 3-5 minutes.
5. Add **Bloom**; agitate 3-5 minutes.
6. Add **Clean Up** in 0.01 g/L steps until target pH is achieved.
7. Validate pH/EC and adjust as necessary.

*Only use Front Row Si if reservoir will be fully used within 48 hours.

DIRECT TO RESERVOIR NOTES

- When using Front Row Si, reservoirs should be fully used within 48 hours.
- Without Front Row Si, reservoirs should be used within 5-7 days.
- Avoid mixing strong oxidizers, especially peroxides into reservoirs. If running a sterile reservoir, use calcium hypochlorite at 0.26 to 0.79 g / 100 liters.
- All feed charts are based on using RO water. If your starting water has any EC, be sure to account for that in the total EC.
- If using PhosZyme, add with Part B.

STEP-BY-STEP



FEED CHART NOTES

These feed charts are not a prescription, but an example of the general ranges and relationship of EC and recipes that can be used. Each facility and cultivation methodology will require customization of EC values. See "EC Considerations".

EC CONSIDERATIONS






Given the variance in facility infrastructure, cultivation methods, and cultivars, it's impossible to give a specific EC prescription that applies to all scenarios. Most facilities feed at 2.0-3.0 EC with Front Row Ag, and our "Standard" and "High Strength" Feed Charts reflect effective feeding strategies in this range.

FEED EC vs Si USAGE RATE	
Feed EC	Si Usage Rate (mL/L)
< 2.3	0.13
2.3-2.7	0.10
2.7-3.1	0.07
3.1-3.5	0.03
> 3.5	0

HIGHER EC	LOWER EC
Smaller pots	Larger pots
Frequent irrigation	Infrequent irrigation
Consistent runoff	Less runoff
Substrate monitoring	No substrate monitoring
Higher PPFD	Lower PPFD
Heavy feeding strains	Lower feeding strains
Higher CO2	Lower CO2
Tight environmental control	Less environmental control

DIRECT TO RESERVOIR (DTR) FEED CHARTS

DTR STANDARD STRENGTH

Week of Flower		1	2	3	4	5	6	7	8	Strain Dependant
Chart Units g/L	Phase	Veg/Moms	Week 1-2	Week 3-5	Week 6-8/9	Final 1-2 Weeks				
	Recipe									
		Veg	Stretch	Stack*	Swell	Ripen				
Base Fertilizer:	EC	2.6	2.4	2.2	2.0	1.6				
PART A	g/L	1.4	1.1	0.9	0.7	0.5				
	Part A EC	1.7	1.3	1.1	0.9	0.6				
PART B	g/L	0.9	0.7	0.6	0.5	0.5				
	Part B EC	0.9	0.6	0.5	0.5	0.5				
BLOOM	g/L		0.5	0.6	0.9	0.7				
	Bloom EC		0.4	0.5	0.7	0.6				

Optional Inputs:

Si (mL)	mL/L	0.1	0.1	0.1	0.1	0.1				
PhosZyme	g/L	0.1	0.1	0.1	0.1	0.1				
	PhosZyme EC	0.1	0.1	0.1	0.1	0.1				

DTR HIGH STRENGTH

Chart Units g/L	Phase	Veg/Moms	Week 1-2	Week 3-5	Week 6-8/9	Final 1-2 Weeks				
		Recipe	Veg	Stretch	Stack*	Swell	Ripen			
	EC	3.0	3.0	2.7	2.4	1.8				
PART A	g/L	1.6	1.4	1.1	0.9	0.5				
	Part A EC	2.0	1.7	1.4	1.1	0.6				
PART B	g/L	1.1	0.9	0.8	0.6	0.6				
	Part B EC	1.0	0.9	0.7	0.6	0.5				
BLOOM	g/L		0.6	0.8	1.0	0.8				
	Bloom EC		0.5	0.6	0.8	0.6				
Si (mL)	mL/L	0.1	0.1	0.1	0.1	0.1				
PhosZyme	g/L	0.1	0.1	0.1	0.1	0.1				
	PhosZyme EC	0.1	0.1	0.1	0.1	0.1				

* For facilities that want to run one recipe throughout flower, use "Stack" recipe.

ADDITIVE USAGE RATES

Additive	Usage Rate	Notes
Front Row Si	0 - 0.13 mL/L	Si usage rate depends on feed EC, please refer to SI vs EC Table.
Triologic	0.26-0.53 mL/L	Recommended to be used 1x per week.
BioFlo	8 mL/L	Use as necessary to remove biofilm from irrigation lines.

PART	CONTRIBUTED EC/g/L
PART A	1.219
PART B	0.965
BLOOM	0.757

