

GUARANTEEDANA	11313
Total Nitrogen (N)	4.0%
3.3% Urea Nitrogen*	
0.35% Ammoniacal Nitro	gen
0.35% Nitrate Nitrogen	
Soluble Potash (K2O)	2.0%
Sulfur (S)	1.0%
Iron (Fe)	1.5%

Derived from: Urea, Ammonium Nitrate, Sulfate of Potash, Ferrous Sulfate, Ascophyllum Nodosum

*2% urea nitrogen stabilized with N-(n-butyl) thiophosphoric triamide ALSO CONTAINS NON-PLANT FOOD INGREDIENT(S): 2.0% Humic Acids

(Derived from Leonardite) 2.0% Kelp (Microbe food) Purpose Statement:

Purpose Statement:
May increase micronutrient uptake,
May increase microbe food.
Information regarding the contents and levels

May increase microbe rood.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

1.0 at = 2.4 lbs at 68° F

 $1.0 \text{ qt} = 2.4 \text{ ms at 65}^{\circ} \text{ P}$ net volume 1.0 qt (.95 l)

DIV Directions to Use

Do not apply to foliage if temperatures are above 80° F or below 40 ° F.

Avoid direct contact with flower blooms to prevent staining.

SHAKE WELL! The mixed product should be agitated prior to and during application.

Healthy Trees: Apply at a rate of 1/2 oz. per caliper inch of the diameter at shoulder height. Dilute with water at a rate of 6 oz. per gal. Spray the trunk of the tree up to a height of 6 ft. or to the first fork. Apply two times/yr. in spring and fall.

Sick or Diseased Trees: Apply at a rate of 1 oz. per caliper inch of diameter at shoulder height. Dilute at a rate of 6 oz. per gal. of water. Spray the trunk of the tree up to the height of 6 ft. or to the first fork. Apply every 30 days as needed.

Fruit Trees: Apply at 1/2 oz. per caliper inch of the diameter at shoulder height before fruit sets. Dilute at 6 oz. per gal. of water. Spray trunk to a height of 6 ft. or to the first fork. No application after fruit sets.

Plants and Ornamentals: Apply at a rate of 6 oz. per 1,000 sq. ft. of bedding material. Dilute at a rate of 6 oz. per gal. of water; apply as a mist to foliage or as a soil drench at the base of the plant.

Palms: Apply at a rate of 1 oz. per caliper inch of diameter at shoulder height.

Dilute at a rate of 6 oz. per gal. of water. Apply as a soil drench at the base of the palm.

Potted Plants and Vegetables: Dilute at 1 oz. per gal. of water and apply weekly as a light soaking to the base of the plant.

Page 1 of 2 | See Back Label

Greene County Fertilizer Company







Manufactured and Guaranteed by Greene County Fertilizer Company, Inc. P.O. BOX 1346, Greensboro, GA 30642 1-855-606-3378 • greenecountyfert.com F2735

1.0 qt = 2.4 lbs at 68° F net volume 1.0 qt (.95 l)



Greene County Fertilizer Company



Fertility Forwards

Store in a cool dry place away from sunlight. Product will store for 2 yrs, in normal warehouse conditions.

Compatibility: These concentrated materials are compatible with most fertilizers and chemicals but should not be mixed directly with other chemicals. Conduct a jar test as needed to ensure compatibility before tank mixing.

Apply in accordance with best management practices (BMP's) established by your Cooperative Extension Service. Observe any State or Local fertilizer application regulations.

Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn/garden.

Caution: This product contains the secondary micronutrient iron. Iron may stain concrete, wood and other pervious and impervious surfaces. Apply only to turf, plants, and soil. If plants are flowering, apply to base of the plant to avoid staining of blooms.

Keep out of reach of children. If product comes in contact with skin or eyes flush with water immediately.

Page 2 of 2 | See Front Label for Application Rates

VISIT
GREENECOUNTYFERT.COM/DIY/
OR MEI PEUL APPUICATION TIPS AND MORE



CONDITIONS OF SALE Seller warmins that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions to the. No one, other than the officer or Seller, is suthorized to page any warming guaranties, or directions concerning this product. Because the time, place, rate of application and other confinioners of the are beyond Seller's control Seller's lability from handling, stonge and use of this product is limited to leplacement of product or refund of purchase pice, buyer assumed it reports that it is not use not in accordance with label in structure. The product names are registered undernative of Greene County Fertilizer Company, inc.



Derived from: Urea, Ammonium Nitrate, Sulfate of Potash, Ferrous Sulfate, Ascophyllum Nodosum

*2% urea nitrogen stabilized with

N-(n-butyl) thiophosphoric triamide ALSO CONTAINS NON-PLANT FOOD

INGREDIENT(S):

2.0% Humic Acids (Derived from Leonardite)
2.0% Kelp (Microbe food)

Purpose Statement:

May increase micronutrient uptake.

May increase microbe food.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html







DIV DIRECTIONS FOR USE

Do not apply to foliage if temperatures are above 80° F or below 40 ° F.

Avoid direct contact with flower blooms to prevent staining.

SHAKE WELL! The mixed product should be agitated prior to and during application.

Healthy Trees: Apply at a rate of 1/2 oz. per caliper inch of the diameter at shoulder height. Dilute with water at a rate of 6 oz. per gal. Spray the trunk of the tree up to a height of 6 ft. or to the first fork. Apply two times per year in the spring and fall.

Sick or Diseased Trees: Apply at a rate of 1 oz. per caliper inch of diameter at shoulder height. Dilute at a rate of 6 oz. per gal. of water. Spray the trunk of the tree up to the height of 6 ft. or to the first fork. Apply every 30 days as needed.

Fruit Trees: Apply at 1/2 oz. per caliper inch of the diameter at shoulder height before fruit sets. Dilute at 6 oz. per gal. of water. Spray trunk to a height of 6 ft. or to the first fork. No application after fruit sets.

Plants and Ornamentals: Apply at a rate of 6 oz. per 1,000 sq. ft. of bedding material. Dilute at a rate of 6 oz. per gal. of water; apply as a mist to foliage or as a soil drench at the base of the plant.

Palms: Apply at a rate of 1 oz. per caliper inch of diameter at shoulder height.

Dilute at a rate of 6 oz. per gal. of water. Apply as a soil drench at the base of the palm.

Potted Plants and Vegetables: Dilute at 1 oz. per gal. of water and apply weekly as a light soaking to the base of the plant.

Store in a cool dry place away from sunlight. Product will store for 2 yrs. in normal warehouse conditions.

Page 1 of 2 See Back Label



1.0 gal = 9.7 lbs at 68° F net volume 1.0 gal (3.8 l)



Manufactured and Guaranteed by Greene County Fertilizer Company, Inc. P.O. BOX 1346, Greensboro, GA 30642 1-855-606-3378 • greenecountyfert.com F2735

1.0 gal = 9.7 lbs at 68° F net volume 1.0 gal (3.8 l)



Compatibility: These concentrated materials are compatible with most fertilizers and chemicals but should not be mixed directly with other chemicals. Conduct a jar test as needed to ensure compatibility before tank mixing. Mix in spray tanks with water and proper agitation. Apply in accordance with best management practices (BMP's) established by your Cooperative Extension Service. Observe any State or Local fertilizer application regulations.

Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn/garden.

Caution: This product contains the secondary micronutrient iron. Iron may stain concrete, wood and other pervious and impervious surfaces. Apply only to turf, plants, and soil. If plants are flowering, apply to base of the plant to avoid staining of blooms.

Keep out of reach of children. If product comes in contact with skin or eyes flush with water immediately.

Page 2 of 2 | See Front Label for Application Rates



VISIT
GREENECOUNTYFERT.COM/DIY/
FOR HELPFUL APPLICATION TIPS AND MORE.



CONDITIONS OF SALE Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than the officer or Seller, is authorized to make any warranty, guarantee, or directions concerning this product. Because the time, place. No one of the product or and other conditions of use are beyond Seller's control Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price. Buyer assumes all responsibility for safety and use not in accordance with label instructions. The product names are registered trademarks of Greene County Fertilizer Company, Inc.



Total Nitrogen (N)4.0%
3.3% Urea Nitrogen*
0.35% Ammoniacal Nitrogen
0.35% Nitrate Nitrogen
Soluble Potash (K ₂ O)2.0%
Sulfur (S)1.0%
Iron (Fe)1.5%

Derived from: Urea, Ammonium Nitrate, Sulfate of Potash, Ferrous Sulfate,

Ascophyllum Nodosum

*2% urea nitrogen stabilized with N-(butyl) thiophosphoric triamide

ALSO CONTAINS NON-PLANT FOOD INGREDIENT(S):

2.0% Humic Acids (Derived from Leonardite)

2.0% Kelp (Microbe food) Purpose Statement:

May increase micronutrient uptake.

May increase microbe food.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html



1.0 gal = 9.7 lbs at 68° F net volume 2.5 gal (9.51)

DIY Directions for Use

Do not apply to foliage if temperatures are above 80° F or below 40° F. Avoid direct contact with flower blooms to prevent staining.

SHAKE WELL! The mixed product should be agitated prior to and during application.

Healthy Trees: Apply at a rate of 1/2 oz. per caliper inch of the diameter at shoulder height. Dilute with water at a rate of 6 oz. per gal. Spray the trunk of the tree up to a height of 6 ft. or to the first fork. Apply two times per year in the spring and fall.

Sick or Diseased Trees: Apply at a rate of 1 oz. per caliper inch of diameter at shoulder height. Dilute at a rate of 6 oz. per gal, of water, Spray the trunk of the tree up to the height of 6 ft. or to the first fork. Apply every 30 days as needed.

Fruit Trees: Apply at 1/2 oz, per caliper inch of the diameter at shoulder height before fruit sets. Dilute at 6 oz. per gal. of water. Spray trunk to a height of 6 ft. or to the first fork. No application after fruit sets.

Plants and Ornamentals: Apply at a rate of 6 oz. per 1,000 sq. ft. of bedding material. Dilute at a rate of 6 oz. per gal. of water; apply as a mist to foliage or as a soil drench at the base of the plant.

Palms: Apply at a rate of 1 oz. per caliper inch of diameter at shoulder height. Dilute at a rate of 6 oz. per gal. of water. Apply as a soil drench at the base of the palm.

Potted Plants and Vegetables: Dilute at 1 oz. per gal. of water and apply weekly as a light soaking to the base of the plant.

Store in a cool dry place away from sunlight. Product will store for 2 yrs. in normal warehouse conditions.

Compatibility: These concentrated materials are compatible with most fertilizers and chemicals but should not be mixed directly with other chemicals. Conduct a jar test as needed to ensure compatibility before tank mixing. Mix in spray tanks with water and proper agitation. Apply in accordance with best management practices (BMP's) established by your Cooperative Extension Service. Observe any State or Local fertilizer application regulations. Do not apply near water, storm drains, or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn/garden, and sweep any product that lands on the driveway, sidewalk, or street back onto your lawn/garden. Caution: This product contains the secondary micronutrient iron. Iron may stain concrete, wood and other pervious and impervious surfaces. Apply only to turf, plants, and soil. If plants are flowering, apply to base of the plant to avoid staining of blooms. Keep out of reach of children. If product comes in contact with skin or eyes flush with water immediately.

Greene County Fertilizer Company







GREENECOUNTYFERT.COM/DIY/

SDS & Labels

Fertility Forward®

Manufactured and Guaranteed by Greene County Fertilizer Company, Inc. • P.O. Box 1346, Greensboro, GA 30642 • 1-855-606-3378 • F2735

CONDITIONS OF SALE Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than the officer or Seller, is authorized to make any warranty, guarantee, or directions concerning this product. Because the time, place, rate of application and other conditions of use are beyond Seller's control Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price. Buyer assumes all responsibility for safety and use not in accordance with label instructions. The product names are registered trademarks of Greene County Fertilizer Company, Inc.









Total Nitrogen (N)	.4.0%
3.3% Urea Nitrogen*	
0.35% Ammoniacal Nitroge	en
0.35% Nitrate Nitrogen	
Soluble Potash (K2O)	
Sulfur (S)	
Iron (Fe)	1.5%

Derived from: Urea, Ammonium Nitrate, Sulfate of Potash, Ferrous Sulfate, Ascophyllum Nodosum

*2% urea nitrogen stabilized with N-(n-butyl) thiophosphoric triamide

ALSO CONTAINS NON-PLANT FOOD INGREDIENT(S):

2.0% Humic Acids (Derived from Leonardite) 2.0% Kelp (Microbe food)

Purpose Statement:

May increase micronutirent uptake. May increase microbe food.

Information regarding the contents and levels of metals in this product is available on the internet at http://aapfco.org/metals.html.

Manufactured and Guaranteed by Greene County Fertilizer Company, Inc. PO Box 1346 • Greensboro, GA 30642 1-855-606-3378 • greenecountyfert.com



PRODUCT DESCRIPTION

4-0-2 FloraGreene[™] *Fertilizer* is engineered to be a complete product for all varieties of landscape plants, ornamentals, tropicals, shrubs, and trees.

APPLICATION RECOMMENDATIONS While soil application is acceptable, **4-0-2** is formulated for foliar application.

- 4-0-2 FloraGreene[™] will absorb through leaf, stem, and cambium tissue and be distributed to the roots through the circulatory systems of plants, ornamentals, shrubs, and trees.
- Application rates of **4-0-2** are very low.
- There is no risk of tissue burn.
- Product can stain. Special care should be taken when applying near homes or hardscapes.

DIY Directions • Storage & Handling

SHAKE WELL! The mixed product should be agitated prior to and during application.

Do not apply to foliage if temperatures are above 80° F or below 40 ° F.

Healthy Trees: Apply at a rate of 1/2 ounce per caliper inch of the diameter at shoulder height. Dilute with water at a rate of 6 ounces per gallon. Spray the trunk of the tree up to a height of 6 feet or to the first fork. Apply two times per year in the spring and fall.

Sick or Diseased Trees: Apply at a rate of 1 ounce per caliper inch of diameter at should height. Dilute at a rate of 6 ounces per gallon of water. Spray the trunk of the tree up to the height of 6 feet or to the first fork. Apply every 30 days as needed.

Fruit Trees: Apply at 1/2 ounce per caliper inch of the diameter at shoulder height before fruit sets. Dilute at 6 ounces per gallon of water. Spray trunk to a height of 6 feet or to the first fork. No application after fruit sets.

Plants and Ornamentals: Apply at a rate of 6 ounces per 1,000 sq. ft. of bedding material. Dilute at a rate of 6 ounces per gallon of water; apply as a mist to foliage or as a soil drench at the base of the plant. Avoid direct contact with flower blooms to prevent staining.

Palms: Apply at a rate of 1 ounce per caliper inch of diameter at shoulder height. Dilute at a rate of 6 ounces per gallon of water. Apply as a soil drench at the base of the palm.

Potted Plants and Vegetables: Dilute at 1 ounce per gallon of water and apply weekly as a light soaking to the base of the plant. Avoid direct contact with flower blooms to prevent staining.

Storage: in a cool dry place away from direct sunlight. Product will store for 2 years under warehouse conditions.

Apply in accordance with best management practices (BMP's) established by your Cooperative Extension Service. Observe any State or Local fertilizer application regulations.

Caution: This product contains the secondary micronutrient iron. Iron may stain concrete, wood and other pervious and impervious surfaces. **Apply** only to turf, plants, and soil. If plants are flowering, apply to base of the plant to avoid staining of blooms.

Keep out of Reach of Children: If product comes in contact with skin or eyes, flush with water immediately.



PACKAGING: 1.0-gal bottle and 2.5-gal bottle

Always read and follow label directions.

