

Drug Facts**Active ingredient (in each brown tablet)**

Ibuprofen USP, 200 mg (NSAID)* Pain reliever/fever reducer

Purpose

*nonsteroidal anti-inflammatory drug

Uses:

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
 - muscular aches ■ backache ■ toothache ■ menstrual cramps ■ the common cold
 - headache ■ minor pain of arthritis

Warnings**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.Symptoms may include: ■ shock ■ facial swelling ■ asthma (wheezing) ■ rash
■ blisters ■ skin reddening ■ hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ take more or for a longer time than directed

- are age 60 or older ■ take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.**Do not use:** ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery**Ask a doctor before use if:** ■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn ►**Drug Facts (continued)**

- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are:

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product: take with food or milk if stomach upset occurs**Stop use and ask a doctor if:**

- you experience any of the following signs of stomach bleeding:
 - have stomach pain that does not get better ■ feel faint
 - have bloody or black stools ■ vomit blood
- you have symptoms of heart problems or stroke:
 - chest pain ■ slurred speech ■ leg swelling
 - trouble breathing ■ weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breastfeeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. ►**Drug Facts (continued)****Directions**

- **do not take more than directed ■ the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- If pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED**
- store between 20°-25°C (68°-77°F) ■ avoid excessive heat 40°C (104°F)
- see vial label for expiration date and lot number ■ use by expiration date on package

Inactive ingredients

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? 1-833-BE-WELLY (1-833-239-3559)Distributed by Welly Health PBC, Minneapolis, MN 55402
1-833-BE-WELLY**Drug Facts****Active ingredients (in each gram)**Bacitracin Zinc (400 units Bacitracin) First aid antibiotic
Neomycin Sulfate (3.5 mg Neomycin) First aid antibiotic
Polymyxin B Sulfate (Polymyxin B 5,000 units) First aid antibiotic**Purpose****Uses** First aid to help prevent infection in minor cuts, scrapes and burns**Warnings****For external use only****Do not use:**

- over large areas of the body ■ in the eyes
- if you are allergic to any of the ingredients ■ longer than 1 week unless directed by a doctor

Ask a doctor before use:

in case of deep or puncture wounds, animal bites, or serious burns

Stop use and ask a doctor if:

- condition persists or gets worse ■ a rash or other allergic reaction develops.

Keep this and all drugs out of the reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.**Directions**

- Clean affected area
- Apply a small amount of this product (an amount equal to the surface area of tip of finger) on area 1 to 3 times daily
- May be covered with a sterile bandage

Other information

- Store at room temperature (do not freeze)
- Do not use any opened or torn packets

Inactive ingredient

white petrolatum

Questions or comments? 1-833-BE-WELLY (1-833-239-3559)

Distributed by Welly Health PBC, Minneapolis, MN 55402 1-833-BE-WELLY

Drug Facts**Active ingredient**

Hydrocortisone 1.0% Anti-itch

Purpose

- Uses** ■ for temporary relief of itching associated with minor skin irritations, inflammation and rashes due to eczema, insect bites, poison ivy, poison oak, poison sumac, soaps, detergents, cosmetics, jewelry, seborrheic dermatitis and psoriasis
- external anal itching
 - other uses of this product should be only under the advice and supervision of a doctor

Warnings**For external use only****Do not use:** for the treatment of diaper rash. Consult a doctor**When using this product:**

- avoid contact with the eyes ■ do not use more than directed unless directed by a doctor
- do not put into the rectum by using fingers or any mechanical device or applicator

Stop use and ask a doctor if:

- condition worsens, or symptoms persist for more than 7 days or clear up and occur again within a few days, and do not begin use of any other Hydrocortisone product unless you have consulted a doctor
- rectal bleeding occurs.

Keep this and all drugs out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.**Directions**

- For adults and children 2 years of age and older: apply to affected area not more than 3 to 4 times daily.
- Children under 2 years of age: do not use, consult a doctor
- Adults for external anal itching: when practical, cleanse the affected area with mild soap and warm water and rinse thoroughly or by patting or blotting with an appropriate cleansing pad. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product
- Children under 12 years of age: for external anal itching, consult a doctor

Other information

- Store at room temperature (do not freeze) ■ Do not use any opened or torn packets

Inactive ingredients

emulsifying wax, ethanol, methylparaben, mineral oil, paraffin, petrolatum, propylparaben, purified water, white wax

Questions or comments? 1-833-BE-WELLY (1-833-239-3559)

Distributed by Welly Health PBC, Minneapolis, MN 55402 1-833-BE-WELLY