

Drug Facts**Active ingredients (in each gram)**

	<i>Purpose</i>
Bacitracin Zinc (400 units Bacitracin).....	First aid antibiotic
Neomycin Sulfate (3.5 mg Neomycin).....	First aid antibiotic
Polymyxin B Sulfate (Polymyxin B 5,000 units).....	First aid antibiotic

Uses

First aid to help prevent infection in minor cuts, scrapes and burns

Warnings

For external use only

Do not use:

- in the eyes ■ over large areas of the body ■ if you are allergic to any of the ingredients
- longer than 1 week unless directed by a doctor

Ask a doctor before use:

in case of deep or puncture wounds, animal bites, or serious burns

Stop use and ask a doctor if:

- condition persists or gets worse
- a rash or other allergic reaction develops.

Keep this and all drugs out of the reach of children.

In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Directions

- Clean affected area
- Apply a small amount of this product (an amount equal to the surface area of tip of finger) on area 1 to 3 times daily ■ May be covered with a sterile bandage

Other information

- Store at room temperature (do not freeze) ■ Do not use any opened or torn packets

Inactive ingredient

white petrolatum

Questions or comments? 1-833-BE-WELLY (1-833-239-3559)

Drug Facts**Active ingredient**

	<i>Purpose</i>
Hydrocortisone 1.0%	Anti-itch

Uses ■ for temporary relief of itching associated with minor skin irritations, inflammation and rashes due to eczema, insect bites, poison ivy, poison oak, poison sumac, soaps, detergents, cosmetics, jewelry, seborrheic dermatitis and psoriasis

- external anal itching
- other uses of this product should be only under the advice and supervision of a doctor

Warnings

For external use only

Do not use: for the treatment of diaper rash. Consult a doctor

When using this product:

- avoid contact with the eyes ■ do not use more than directed unless directed by a doctor
- do not put into the rectum by using fingers or any mechanical device or applicator

Stop use and ask a doctor if:

- condition worsens, or symptoms persist for more than 7 days or clear up and occur again within a few days, and do not begin use of any other Hydrocortisone product unless you have consulted a doctor
- rectal bleeding occurs.

Keep this and all drugs out of the reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- For adults and children 2 years of age and older: apply to affected area not more than 3 to 4 times daily.
- Children under 2 years of age: do not use, consult a doctor
- Adults for external anal itching: when practical, cleanse the affected area with mild soap and warm water and rinse thoroughly or by patting or blotting with an appropriate cleansing pad. Gently dry by patting or blotting with toilet tissue or a soft cloth before application of this product
- Children under 12 years of age: for external anal itching, consult a doctor

Other information

- Store at room temperature (do not freeze) ■ Do not use any opened or torn packets

Inactive ingredients emulsifying wax, ethanol, methylparaben, mineral oil, paraffin, petrolatum, propylparaben, purified water, white wax

Questions or comments? 1-833-BE-WELLY (1-833-239-3559)