



STOMACH SOOTHER

Children's Stomach Soother

Drug Facts
Active ingredient Purpose (in each tablet)
 Calcium carbonate 400 mgAntacid

Uses for the relief of
 ■ heartburn ■ sour stomach
 ■ acid indigestion
 ■ upset stomach associated with these symptoms

Warnings
 Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

When using this product
 ■ do not take more than 3 chewable tablets (ages 2-5) or 6 chewable tablets (ages 6-11) in a 24-hour period
 ■ do not use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor
Keep out of reach of children.

Directions
 ■ find the right dose on the chart on the next page based on weight (preferred); otherwise use age ▶

Drug Facts (continued)
 ■ chew or crush tablets completely before swallowing; do not swallow tablets whole
 ■ repeat dose as needed, not more than 3 times a day

Weight (lb)	Age (yr)	Dose (chewable tablets)*
under 24	under 2	ask a doctor
24-47	2-5	1 tablet
48-95	6-11	2 tablets

*or as directed by a doctor

Other information
 ■ Each chewable tablet contains: calcium 160 mg
 ■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ avoid excessive humidity
 ■ see label for expiration date and lot number
 ■ use by expiration date on package

Drug Facts (continued)
Inactive Ingredients acacia, cornstarch, D&C red# 27 aluminum lake, D&C red #30 aluminum lake, FD&C red #40 aluminum lake, flavor, magnesium stearate, maltodextrin, silicon dioxide, sodium lauryl sulfate, sucrose, talc

Questions or comments?
 1-833-BE-WELLY (1-833-239-3559)



ALLERGY RELIEF

Children's Allergy Relief

Drug Facts
Active ingredient Purpose (in each chewable tablet)
 Diphenhydramine HCl 12.5 mg.Antihistamine

Use
 temporarily relieves runny nose and relieves sneezing, itching of the nose or throat, and itchy, watery eyes due to hay fever or other upper respiratory allergies. ▶

Drug Facts (continued)
Warnings
Do not use
 ■ with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have
 ■ a breathing problem such as emphysema or chronic bronchitis
 ■ glaucoma
 ■ difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product
 ■ may cause excitability in children
 ■ may cause marked drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness effect
 ■ avoid alcoholic beverages
 ■ use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. ▶

Drug Facts (continued)
Directions
 ■ find right dose on the chart below
 ■ chew or crush tablets completely before swallowing; do not swallow tablets whole
 ■ take every 4 to 6 hours, or as directed by a doctor
 ■ do not take more than 6 times in 24 hours

Adults and children 12 years of age and over	2 to 4 chewable tablets (25 to 50 mg)
Children 6 to under 12 years of age	1 to 2 chewable tablets (12.5 to 25 mg)
Children under 6 years of age	Consult a doctor

Other information
 ■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ avoid high humidity
 ■ see label for expiration date and lot number
 ■ use by expiration date on package ▶

Drug Facts (continued)
Inactive Ingredients
 D&C Red #27 Aluminum Lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, FD&C Blue #1 aluminum lake, flavor, hydroxypropyl cellulose, magnesium stearate, mannitol, stearic acid, sucralose, sucrose

Questions or comments?
 1-833-BE-WELLY (1-833-239-3559)



PAIN AND FEVER RELIEF

Children's Pain & Fever Relief

Drug Facts
Active ingredient Purpose (in each chewable tablet)
 Acetaminophen 160 mg.....Pain Reliever/
 Fever Reducer

Uses
 ■ for the temporary relief of minor aches and pains associated with
 ■ the common cold ■ sore throat
 ■ headache ■ toothache ■ flu
 ■ and to reduce fever ▶

Drug Facts (continued)
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes
 ■ more than 5 doses in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin. ▶

Drug Facts (continued)
Stop use and ask a doctor if
 ■ pain persists for more than 5 days or gets worse
 ■ fever persists for more than 3 days or gets worse.
 ■ new symptoms occur
 ■ redness or swelling is present
 These could be signs of a serious condition
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical even if you do not notice any signs or symptoms.

Directions
 ■ this product does not contain directions or complete warnings for adult use
 ■ do not give more than directed
 ■ find the right dose on the chart below. If possible, use weight to dose; otherwise, use age.
 ■ chew or crush tablets completely before swallowing; do not swallow tablets whole
 ■ repeat dose every 4 hours while symptoms last
 ■ do not give more than 5 times in 24 hours
 ■ do not give for more than 5 days unless directed by a doctor ▶

Drug Facts (continued)

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 ½ tablets
48-59	6-8	2 tablets
60-71	9-10	2 ½ tablets
72-95	11	3 tablets

*or as directed by a doctor

Other information
 ■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ avoid high humidity
 ■ see label for expiration date and lot number
 ■ use by expiration date on package

Inactive ingredients citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, FD&C blue #1 aluminum lake, flavors, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions or comments?
 1-833-BE-WELLY (1-833-239-3559)