

MOTOMCO


FARM & HOME®

NEW

# RAMPAGE® SOFT BAIT



Photo may not be representative of a label-consistent bait placement



**KILLS RATS, MICE & MEADOW VOLES\***

Net Weight:  
**8 lbs (3.6 kg)**

ACTIVE INGREDIENT:	
Bromethalin (CAS #63333-35-7):	0.01%
OTHER INGREDIENTS†:	99.99%
†Contains Denatonium Benzoate	<b>TOTAL 100.00%</b>

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See back panels for First Aid, additional  
Precautionary Statements, and Directions for Use.

\*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

FIRST AID HAVE LABEL WHEN OBTAINING TREATMENT ADVICE	
<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.</li> </ul>
<b>If in Eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> </ul>
<b>TREATMENT FOR PET POISONING</b>	
If an animal eats this bait, call a veterinarian or 1-877-854-2494 at once.	
<b>NOTE TO PHYSICIAN OR VETERINARIAN</b>	
Contains the nerve poison bromethalin. This product is not an anticoagulant type rodenticide. For humans or animals ingesting bait and/or showing poisoning signs (such as muscle tremors, loss of hind limb use, or seizures for animals), limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.	

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**  
**CAUTION:** Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Keep away from children, domestic animals and pets. All handlers (including applicators) must wear shoes plus socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear waterproof gloves.

**User Safety Requirements**  
 Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing them. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

**ENVIRONMENTAL HAZARDS**  
 This product is extremely toxic to mammals and birds. Dogs and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

**DIRECTIONS FOR USE**  
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for the sites, pests, and application methods described on this label.

**IMPORTANT:** Do not expose children, pets, or nontarget animals to rodenticides. To help prevent exposure:

1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hooled livestock, raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled or unconsumed bait as specified on this label.

# RAMPAGE<sup>®</sup> SOFT BAIT



**FOR INDOOR AND OUTDOOR USE**

**KILLS RATS, MICE & MEADOW VOLES\***

**KILLS ANTICOAGULANT RESISTANT NORWAY RATS, ROOF RATS AND HOUSE MICE**

Due to Stop Feed Action, Rats and Mice Typically Feed Less or Cease Feeding Altogether After Consuming a Lethal Dose

**ACTIVE INGREDIENT:**  
 Bromethalin (CAS #63333-35-7): . . . . . 0.01%  
**OTHER INGREDIENTS:** . . . . . 99.99%  
 †Contains Denatonium Benzoate **TOTAL 100.00%**

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See side panels for First Aid, additional Precautionary Statements, and Directions for Use.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container in a cool, dry place inaccessible to children and pets.

**Pesticide Disposal:** Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

**Plastic:** Offer for recycling or reconditioning; or puncture and dispose of container in a sanitary landfill.

**Paper:** Dispose of empty container by placing in trash, at an approved disposal facility or by incineration.

**WARRANTY:** To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

\*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

Available Exclusively From:  
**MOTOMCO** **Net Weight: 8 lbs (3.6 kg)**  
 3699 Kinsman Blvd. **0.28 oz (8g) Sachets**  
 Madison, WI 53704 U.S.A. EPA REG. NO. 12455-149-3240  
 www.motomco.com EPA EST. NOs. 12455-WI-1k  
**MADE IN USA** 12455-WI-2P, 12455-WI-3W  
**The World Leader In Rodent Control Technology<sup>®</sup>** Superscript is first letter of the lot number.

**DIRECTIONS FOR USE** (Continued from other panel)

Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used wherever children, pets, non-target mammals or birds may have access to the bait placement location.

**USE RESTRICTIONS:** This product may only be used to control the following rodent pests in and around man-made structures: House mouse (*Mus musculus*), Norway rat (*Rattus norvegicus*), Roof rat (*Rattus rattus*), Cotton rat\* (*Sigmodon hispidus*), Eastern harvest mouse\* (*Reithrodontomys humuli*), Golden mouse\* (*Ochrotomys nuttalli*), Polynesian rat\* (*Rattus exulans*), Meadow vole\* (*Microtus pennsylvanicus*), White-throated woodrat\* (*Neotoma albigula*), Southern plains woodrat\* (*Neotoma micropsus*), Mexican woodrat\* (*Neotoma mexicana*). This product must be used in and within 100 feet of man-made harboring or attracting rodent infestations. Examples of such structures include homes and/or permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or baling, beyond 100 feet from a structure, as defined above, is prohibited. This product must not be applied directly to food or feed crops.

Burrow baiting with RAMPAGE<sup>®</sup> Soft Bait is prohibited.

Do not place bait near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment. Do not apply directly to food or feed crops. Do not broadcast bait.

Do not sell this product in individual containers holding less than 4 pounds of bait.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats, mice, or meadow voles\* will most likely find and consume bait. Generally, these are along walls, by gnawed openings, in corners and signs have been observed. Remove as much alternative food as possible.

**APPLICATION DIRECTIONS:**  
 The active ingredient in this product is different from anticoagulant rodenticides. Individual rats and mice typically feed less or cease feeding altogether after consuming a toxic dose. Bait may be removed from the sachet and placed in a feeding device that secures the unwrapped bait in a bait station. Discard empty wrapper in the trash.

**Rats:**  
 Place 4 to 24 sachets per placement (usually at intervals of 15- to 30- feet) in infested areas. Adjust the amount of bait applied to the level of rat feeding expected at each bait placement.

**Mice and Meadow Voles\*:**  
 Apply 1 to 2 sachets per placement. Space placements at intervals of 8- to 12- feet in infested areas. Larger placements, up to 4 sachets, may be needed at points of extremely high mouse or meadow vole\* activity.

**Baiting Strategy:** Highest bait consumption is expected to occur on the first day or two after treatment begins. After several days, inspect the placement and replenish bait at sites where there is evidence of heavy feeding or where there is continued evidence of rodent activity. Normally, the volume of bait consumed will be significantly less with bromethalin baits than with anticoagulant baits. Although one night's feeding will usually provide a lethal dose for mice and rats, maintain bait for at least one week to provide all individuals a chance to feed on the bait. Continue baiting until all signs of feeding have ceased.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Wearing waterproof gloves, collect and dispose of all dead, exposed animals and leftover bait. To discourage reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish them as needed.



576RS

120919/09-19 Product Code: 31588