

MaxMicrobe



52515154500

Application and Uses: SHAKE WELL BEFORE EACH USE

MaxMicrobe can be used in a multitude of agriculture environments. Applications can be made/done by the use of hand sprayers, hose end sprayers, garden sprayers, or automatic irrigation system injection. Can be used for foliar feeding; mixed with water for soil or coco; or mixed into nutrients and added to reservoir for hydroponics. Safe for use on all equipment. Can be used as pre-plant or pre-emergence soil spray, as well as used as a direct foliar spray. Can be used on seeds, in soil, and directly on plants. PH Neutral

Guaranteed Analysis

CONTENTS:

NON PLANT INGREDIENTS:

Mixture of microbial strains. 4.55% (nominal)

Azospirillum brasilense. 1.00e+07 (10,000,000) cfu/ml

Bacillus altitudinis. 1.00e+08 (100,000,000) cfu/ml

Bacillus amyloliquefaciens. 1.00e+07 (10,000,000) cfu/ml

Bacillus licheniformis 1.00e+07 (10,000,000) cfu/ml

Cellulomonas cellasea 1.00e+07 (10,000,000) cfu/ml

Pseudomonas fluorescens 1.00e+07 (10,000,000) cfu/ml

Pseudomonas putida 1.00e+08 (100,000,000) cfu/ml

Pseudomonas stutzeri 1.00e+08 (100,000,000) cfu/ml

Streptomyces albidoflavus 1.00e+05 (100,000) cfu/ml

TOTAL350,100,000 cfu/mL

INERT INGREDIENT:

Water 95.1%

Ferment Residue 0.35% (nominal) (Non Animal)

MaxMicrobe guarantees product conforms to description on the label. MaxMicrobe guarantees information is reasonable for the purposes stated on label when used in accordance with directions. MaxMicrobe does not warranty or guarantee any particular level of results, as conditions beyond our control may affect results.

MaxMicrobe limits, excludes, and disclaims liability for incidental and consequential damages arising out of use of this product and any use contrary to the label directions. Buyer accepts these conditions with use of product. Warranty is limited to replacement of product that is deemed to be defective in manufacturing or packaging.

CAUTION: Not for internal consumption. Keep away from children and pets. If swallowed, contact a physician and call your local poison control center.

STORAGE: Store in cool dry place away from direct sunlight. Avoid freezing and temperature above 110°

Suggested Application Rates:

Expiration Date: _____

FOLIAR FEED: 1-2 ml/Gallon

GROWTH PHASE

WEEK	GROWTH STAGE	SOIL/COCO	HYDRO
1 & 2	CLONE	0.5 ml/GAL	0.5 ml/GAL
3 & 4	VEGETATION	0.5 ml/GAL	0.5 ml/GAL
5, 6 & 7	AGGRESSIVE VEGETATION	0.8 ml/GAL	0.6 ml/GAL

BLOOM PHASE

WEEK	GROWTH STAGE	SOIL/COCO	HYDRO
TRANSITION	TRANSITION	1.2 ml/GAL	0.5 ml/GAL
1 & 2	BLOOM	1.2 ml/GAL	0.5 ml/GAL
3 & 4	BLOOM	1.5 ml/GAL	0.5 ml/GAL
5, 6 & 7	AGGRESSIVE BLOOM	3.5 ml/GAL	0.6 ml/GAL
8	FLUSH	1.0 ml/GAL	0.5 ml/GAL

Note: Certain types of fungicides may reduce effectiveness. May cause undesired results if not used as directed, i.e. foliar burn, low yield. Use only as intended. Use only for purpose intended as described on label. For best results, apply in early morning or late evening. Best if applied without direct sunlight.

Distributed by MaxMicrobe, 113 North 700 West, Building 5C, Bay 5 Ogden UT 84404



Grown and blended
in the United States

www.maxmicrobe.com