GEL MAX FLAVOR FUSION

SH□EK

14- 14 14-

44-44

- 44

DENTAL WARRANTY DISCLAIMER

CERTIFICATE OF LIMITED WARRANTY AND DISCLAIMER

WARRANTY: Shock Doctor, Inc. ("Shock Doctor") warrants the within mouthguard unit to be free from defects in material and workmanship which could cause damage or injury to the sound, natural teeth of the purchaser herein when properly used in a supervised athletic contest or coach-supervised training session. This warranty shall extend for a period of one (1) year from the date of purchase.

Shock Doctor's obligations under this warranty shall be limited to the lesser of:

- i. The actual dental expense incurred by the user; or
- ii. \$1094 per injured or replaced tooth, up to a maximum liability of \$35,000 per incident.

Shock Doctor's obligations hereunder shall be reduced by the amount of any insurance collected or payable for the damage or injury claimed. This certificate does not provide payment or indemnity for orthodontic treatments or preparation of a tooth to be used as an anchor for a bridge.

Shock Doctor shall have no responsibility or liability hereunder unless:

- i. The mouthguard was fitted by claimant in accordance with the printed instructions accompanying this warranty; and
- ii. A complete written notice of the injury or damage is received by Shock Doctor within thirty (30) days after the occurrence of the injury or damage; and
- iii. The notice of injury is substantiated by a statement of a licensed dentist providing necessary dental treatment within thirty (30) days after the date of injury or damage and a statement of a participating game official, if a supervised contest, or a supervising coach or school superintendent, if a practice session, that the mouthguard unit was in proper use at the time of injury; and
- iv. The within mouthguard unit is returned to Shock Doctor with the written notice of injury. The mouthguard unit shall become the property of Shock Doctor upon receipt of return.

This warranty is made in lieu of all other warranties, express or implied, including but not limited to the implied warranty for a particular purpose and any implied warranty arising out of a course of a dealing, custom, usage or trade. Except as is otherwise expressly provided herein, Shock Doctor disclaims all liability for loss, injury or damage based upon negligent design or defects in materials and workmanship for the within mouthguard.

Keep this dental warranty, disclaimer and store receipt in your records. In case of injury, mail mouthguard, store receipt, the notice of injury and accompanying statements to Shock Doctor. Inc.

Shock Doctor, Inc., 11488 Slater Avenue, Fountain Valley, CA 92708 Phone: (800) 233-6956 | Fax: (888) 446-5999 | www.shockdoctor.com

GEL MAX FLAVOR FUSION



14- 44

10- 10

44- 44

44

10- 10

40- 40

10- 10 10-

44-44

-

44

FITTING INSTRUCTIONS

WARNING!

NOT FOR USE WITH BRACES DO NOT CHEW YOUR MOUTHGUARD

In order for a mouthguard to protect, fit, and last it must be worn properly on your teeth and gums as indicated in the fitting instructions. If any mouthguard is chewed or not worn properly, its protection, fit and durability will be greatly diminished.

TO FIT THIS MOUTHGUARD YOU WILL NEED:

> Boiling water > Saucepan > Mirror > Slotted spoon > Cold tap water > Timer

FITTING INSTRUCTIONS:

Insert Tether into Gel Max Flavor Fusion Strapped mouthguard as an aid in holding the mouthguard in hot water. For a Strapless version, use a slotted spoon to insert and remove mouthguard from hot water.

- 1. Boil at least 4 cups of water in a sauce pan or similar. The water must reach a rolling boil.
- When water reaches rolling boil, remove pan from heat source. Let water stand for 20 seconds before immersing mouthguard
- Immerse the mouthguard into the hot water, holding it under water with the
 mouthguard tether as a handle, or with the tip of the slotted spoon. Turn the mouthguard
 so the gel layer is facing down (this is not necessary, but is recommended.)
- Carefully remove mouthguard from hot water with a slotted spoon after 60 seconds and shake excess water from mouthguard
- 5. Lick your lips before placing mouthguard in your mouth.
- Watching in mirror, line up mouthguard carefully with the centerline of upper teeth. Fit
 mouthguard onto and around upper teeth. Press bottom of mouthguard into upper teeth,
 also pressing thumb into front of arch, behind front teeth, up to palate.
- 7. Bite down firmly with mouth closed. Suck in strongly and press along lips and cheeks to help form the mouthguard around the teeth.
- Continue to wear mouthguard for at least 45 seconds and then remove. The mouthguard
 can be immersed into cold water with ice cubes to speed cooling, but mouthguard will
 have it's best retention and fit once it cools and sets after a few hours.
- 9. Repeat steps 1 through 8 if fit is not accurate.

NOTE: Mouthguards are best stored in mouthguard cases while not in use. Clean your mouthguard regularly by washing under hot water. If using soap, rinse the mouthguard thoroughly before wearing.