

## Artificial sweeteners linked to risk of weight gain, heart disease and other health issues



Tea with sweetener in a spoon (stock image).

Credit: © photosiber / Fotolia

Artificial sweeteners may be associated with long-term weight gain and increased risk of obesity, diabetes, high blood pressure and heart disease, according to a new study published in *CMAJ (Canadian Medical Association Journal)*

Consumption of artificial sweeteners, such as aspartame and sucralose, is widespread and increasing. Emerging data indicate that artificial, or nonnutritive, sweeteners may have negative effects on metabolism, gut bacteria and appetite, although the evidence is conflicting.

To better understand whether consuming artificial sweeteners is associated with negative long-term effects on weight and heart disease, researchers from the University of Manitoba's George & Fay Yee Centre for Healthcare Innovation conducted a systematic review of 37 studies that followed over 400,000 people for an average of 10 years. Only 7 of these studies were randomized controlled trials (the gold standard in clinical research), involving 1003 people followed for 6 months on average.

The trials did not show a consistent effect of artificial sweeteners on weight loss, and the longer observational studies showed a link between consumption of artificial sweeteners and relatively higher risks of weight gain and obesity, high blood pressure, diabetes, heart disease and other health issues.

"Despite the fact that millions of individuals routinely consume artificial sweeteners, relatively few patients have been included in clinical trials of these products," said author Dr. Ryan Zarychanski, Assistant Professor, Rady Faculty of Health Sciences, University of Manitoba. "We found that data from clinical trials do not clearly support the intended benefits of artificial sweeteners for weight management."

"Caution is warranted until the long-term health effects of artificial sweeteners are fully characterized," said lead author Dr. Meghan Azad, Assistant Professor, Rady Faculty of Health Sciences, University of Manitoba. Her team at the Children's Hospital Research Institute of Manitoba is undertaking a new study to understand how artificial sweetener consumption by pregnant women may influence weight gain, metabolism and gut bacteria in their infants.

"Given the widespread and increasing use of artificial sweeteners, and the current epidemic of obesity and related diseases, more research is needed to determine the long-term risks and benefits of these products," said Azad.

### Story Source:

Materials provided by **Canadian Medical Association Journal**. *Note: Content may be edited for style and length.*

### Journal Reference:

1. Meghan B. Azad, Ahmed M. Abou-Setta, Bhupendrasinh F. Chauhan, Rasheda Rabbani, Justin Lys, Leslie Copstein, Amrinder Mann, Maya M. Jeyaraman, Ashleigh E. Reid, Michelle Fiander, Dylan S. MacKay, Jon McGavock, Brandy Wicklow, Ryan Zarychanski. **Nonnutritive sweeteners and cardiometabolic health: a systematic review and meta-analysis of randomized controlled trials and prospective cohort studies.** *Canadian Medical Association Journal*, 2017; 189 (28): E929 DOI: 10.1503/cmaj.161390

### Cite This Page:

MLA	APA	Chicago
-----	-----	---------

Canadian Medical Association Journal. "Artificial sweeteners linked to risk of weight gain, heart disease and other health issues." ScienceDaily. ScienceDaily, 17 July 2017.

<[www.sciencedaily.com/releases/2017/07/170717091043.htm](http://www.sciencedaily.com/releases/2017/07/170717091043.htm)>.

### RELATED STORIES

Severe Obesity Revealed as a Stand-Alone High-Risk Factor for Heart Failure

Aug. 22, 2016 — A study of more than 13,000 people has found that even after accounting for such risk factors as high blood pressure, high cholesterol and diabetes, so-called morbid obesity appears to stand alone as ...

**read more »**

Obesity and Type 2 Diabetes Harm Bone Health

Nov. 17, 2015 — In a new animal study, researchers examined how the development of obesity and insulin resistance contribute to bone-fracture risk and whether exercise prevents weight gain and diabetes and protects ...

**read more »**

Women With Hypertension in Pregnancy and Their Siblings Face Increased Risk of Heart Disease

Aug. 27, 2015 — High blood pressure during pregnancy is a risk factor for future hypertension and cardiovascular disease, but it's not clear if this increased risk is because these women are more likely to have a ... **read more »**

## Excess Weight at One Year Postpartum Increases Moms' Risk for Diabetes, Heart Problems

Mar. 25, 2014 — Watch out for weight gain within a year of giving birth, to prevent new risk of diabetes and cardiovascular disease, experts say. While it has long been believed that not losing 'baby ... **read more »**

### FROM AROUND THE WEB

---

*Below are relevant articles that may interest you. ScienceDaily shares links with scholarly publications in the TrendMD network and earns revenue from third-party advertisers, where indicated.*