

8 WEEK HYDROPONIC FEED SCHEDULE FOR HIGH VALUE CROPS IN HYDROPONIC SYSTEMS

Measurements are in mL / US Gallon (500 PPM SCALE)	VEGETATIVE STAGE						FLOWERING - RIPENING STAGE							
	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8
MICRO	4	5	6	8	8	8	6	7	7	8	8	8	6	6
GROW	5	7	10	12	14	14	3	3	3	3	3	2	1	1
BLOOM	2	3	3	4	4	4	9	9	9	17	17	19	21	21
CALNESIUM	1.5	1.5	1.5	1.5	1.5	1.5	2.5	2.5	5	5	5	5	5	5
SUPER B+	6	6	6	6	6	6	8	8	8	8	8	8	8	8
ROYAL GOLD	7	7	7	9	g	9	11	11	11	11	11	11	11	11
PLANTACILLIN	2	2	2	2	4	4	6	6	6	6	6	6	6	6
CARBO BLAST	2	2	2	2	4	4	8	8	8	8	8	8	8	8
PROP-O-GATOR	5	5	5	5	10	10	11	10	5	5				
TON O BUD										8	8			
BUD BOOM												8	8	8
SUGGESTED PPM RANGE	500-700	700-900	900-1100	1100-1200	1100-1400	1100-1400	800-1100	900-1200	900-1200	1100-1500	1100-1500	1000-1400	1000-1400	1000-1400

- 1) Always mix Micro first before adding Grow or Bloom, allowing additional mixing time between each product while maintaining a pH between 5.8-6.3.
- 2) At water levels between 200-350 PPM, use Hard Water Micro instead of Micro. At levels above 350 PPM, use Hard Water Micro without Calnesium.
- 3) If vegging longer than 6 weeks, repeat Week 6 for additional weeks. If flowering longer than 8 weeks, repeat Week 8 for each additional week of flower.
- 4) For cuttings and clones, use 1/4 strength of Week 1 for Micro, Grow, and Bloom with 4mL/Gal of Super B+ and 5mL/Gal of Prop-O-Gator until rooted.
- 5) In the case of tip burn or nutrient lock up, scale back measurements by 10-20%. In the last 3-5 days before harvest, flush with pH adjusted water alone.