

8 WEEK HYDROPONIC FEED SCHEDULE FOR HIGH VALUE CROPS IN HYDROPONIC SYSTEMS

Measurements are in mL / Litre 3.785L = 1 Gallon	VEGETATIVE STAGE						FLOWERING - RIPENING STAGE							
	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 1	WEEK 2	WEEK 3	WEEK 4	WEEK 5	WEEK 6	WEEK 7	WEEK 8
MICRO	1	1.25	1.5	2	2	2	1.5	1.75	1.75	2	2	2	1.5	1.5
GROW	1.25	1.75	2.5	3	3.5	3.5	0.75	0.75	0.75	0.75	0.75	0.5	0.25	0.25
BLOOM	0.5	0.75	0.75	1	1	1	2.25	2.25	2.25	4.25	4.25	4.75	<i>5.25</i>	<i>5.25</i>
CALNESIUM	0.4	0.4	0.4	0.4	0.4	0.4	0.6	0.6	1.25	1.25	1.25	1.25	1.25	1.25
SUPER B+	1.5	1.5	1.5	1.5	1.5	1.5	2	2	2	2	2	2	2	2
ROYAL GOLD	1.75	1.75	1.75	2.25	2.25	2.25	2.75	2.75	2.75	2.75	2.75	2.75	2.75	2.75
PLANTACILLIN	0.5	0.5	0.5	0.5	1	1	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
CARBO BLAST	0.5	0.5	0.5	0.5	1	1	2	2	2	2	2	2	2	2
PROP-O-GATOR	1.25	1.25	1.25	1.25	2.5	2.5	2.75	2.5	1.25	1.25				
TON O BUD										2	2			
BUD BOOM												2	2	2
SUGGESTED EC RANGE	1 - 1.4	1.4 - 1.8	1.8 - 2.2	2.2 - 2.4	2.2 - 2.8	2.2 - 2.8	1.1 - 2.2	1.8 - 2.4	1.8 - 2.4	2.2 - 3	2.2 - 3	2 - 2.8	2 - 2.8	2 - 2.8

- 1) Always mix Micro first before adding Grow or Bloom, allowing additional mixing time between each product while maintaining a pH between 5.8-6.3.
- 2) At water levels between 0.4-0.7 EC, use Hard Water Micro instead of Micro. At levels above 0.7 EC, use Hard Water Micro without Calnesium.
- 3) If vegging longer than 6 weeks, repeat Week 6 for additional weeks. If flowering longer than 8 weeks, repeat Week 8 for each additional week of flower.
- 4) For cuttings and clones, use 1/4 strength of Week 1 for Micro, Grow, and Bloom with 1mL/Gal of Super B+ and 1.25mL/Gal of Prop-O-Gator until rooted.
- 5) In the case of tip burn or nutrient lock up, scale back measurements by 10-20%. In the last 3-5 days before harvest, flush with pH adjusted water alone.