leturn	<b>Authorization For</b>	rm			Return Aut	horization #:
					Date Reque	sted:
. Cus	stomer Details					
Name: Address:		Company:			Pł	ione:
					Fa	Fax:
					Eı	nail:
City:		State:	Zip:		Co	ountry:
. Pro	duct Details	·			•	
RunO	110	☐ With Time Control☐ Without Time Contro	Seria	l#:	In	voice # :
	n for Return:		<u> </u>		Da	nte of Purchase:
□ The l	LED light is blinking and there does not seem to be any			rrent.		
Pleas	earch Conditions se provide answers to the You are working with Protein RNA	• .	nich may h	•	sually REUS	ause to the problem. Than  ■ the electrophoresis gel
Pleas	se provide answers to the You are working with   Protein  RNA	□ DNA □ Other:	•	Do you u running b	sually REUS	E the electrophoresis gel  □ NO
Pleas	se provide answers to the You are working with   Protein  RNA  When running electrons	□ DNA □ Other: phoresis gels, you use	5)	Do you u running b runsing b runsing b	sually <b>REUS</b> uffer?	E the electrophoresis gel  NO nes?
Pleas	se provide answers to the You are working with   Protein  RNA	□ DNA □ Other:  phoresis gels, you use □ 1 X TAE buffer	5)	Do you urunning buryES If YES, h If you RE make up urundad run	sually REUS uffer? ow many time USE the gel for the buffer ning buffer	E the electrophoresis gel  NO nes?
Pleas	se provide answers to the You are working with  Protein RNA  When running electro 0.5 X TAE buffer  Other: Other:	□ DNA □ Other:  phoresis gels, you use □ 1 X TAE buffer □ 1 X TBE buffer	5)	Do you u running b  YES If YES, h  If you RE make up  add run  Other:	sually <b>REUS</b> uffer?  ow many time USE the gel for the buffer ning buffer	E the electrophoresis gel  □ NO  nes?  running buffer, how do yo evaporation? □ add DI water
Pleas 1) 2)	se provide answers to the You are working with  Protein RNA  When running electro 0.5 X TAE buffer Other:  Obyou make your ow running buffer?	□ DNA □ Other:  phoresis gels, you use □ 1 X TAE buffer □ 1 X TBE buffer	5)	Do you u running b YES If YES, h If you RE make up add run Other: In genera	sually <b>REUS</b> uffer? ow many time USE the gel for the buffer ning buffer I, how long d	E the electrophoresis gel  □ NO nes? running buffer, how do you evaporation? □ add DI water  ooes it take you to run a ge
Pleas 1) 2)	you are working with Protein RNA When running electro 0.5 X TAE buffer 0.5 X TBE buffer Other: Do you make your ow running buffer? YES	□ DNA □ Other:  phoresis gels, you use □ 1 X TAE buffer □ 1 X TBE buffer  //n electrophoresis gel □ NO	5)	Do you urunning buryES If YES, h  If you RE make up add run urun Other:  In genera urun less tha	sually <b>REUS</b> uffer? ow many time USE the gel for the buffer ning buffer I, how long den 20 mins	E the electrophoresis gel  □ NO nes?  running buffer, how do you evaporation? □ add DI water  loes it take you to run a gel □ 20 to 40 mins
Pleas 1) 2)	what stock concentrate	DNA Dother: DNA Dother: Phoresis gels, you use 1 X TAE buffer 1 X TBE buffer No electrophoresis gel NO ouy from?	5)	Do you urunning b  YES If YES, h  If you RE make up add run Other: In genera less that 41 to 60	sually <b>REUS</b> uffer?  ow many time USE the gel for the buffer ning buffer  I, how long den 20 mins of mins cettrophoresis	DE the electrophoresis gel  □ NO  nes?  running buffer, how do you revaporation?  □ add DI water  loes it take you to run a gel □ 20 to 40 mins □ Other: □ system hot after a run?
Pleas 1) 2) 3)	what stock concentrate electrophoresis gel run	DNA Dother:  phoresis gels, you use 1 X TAE buffer 1 X TBE buffer  NO electrophoresis gel NO ouy from?  tion is your	<ul><li>5)</li><li>6)</li><li>7)</li></ul>	Do you urunning buryES If YES, h  If you RE make up to add run uruning buryES In generaurung less that urunung 41 to 66 Is your elurung slightly	sually <b>REUS</b> uffer? ow many time USE the gel for the buffer ning buffer  I, how long don 20 mins o mins cettrophoresis above room	E the electrophoresis gel  □ NO nes?  running buffer, how do you evaporation? □ add DI water  loes it take you to run a gel □ 20 to 40 mins □ Other: □ system hot after a run? temperature
Pleas 1) 2) 3)	what stock concentrate electrophoresis gel run	DNA Dother:  phoresis gels, you use 1 X TAE buffer 1 X TBE buffer  NO electrophoresis gel NO ouy from?  tion is your	<ul><li>5)</li><li>6)</li><li>7)</li></ul>	Do you urunning by YES If YES, h  If you RE make up add run Other:  In genera less that 41 to 60  Is your el slightly feels w	sually REUS uffer? ow many time USE the gel for the buffer ning buffer I, how long den 20 mins mins mins ectrophoresis above room arm ( around	E the electrophoresis gel  □ NO nes?  running buffer, how do you evaporation? □ add DI water  loes it take you to run a gel □ 20 to 40 mins □ Other: □ system hot after a run? temperature 30 – 45 °C)
Pleas 1) 2) 3)	what stock concentrate	DNA Dother:  phoresis gels, you use 1 X TAE buffer 1 X TBE buffer  NO electrophoresis gel NO ouy from?  tion is your	<ul><li>5)</li><li>6)</li><li>7)</li></ul>	Do you urunning by YES If YES, h  If you RE make up add run Other:  In genera less that 41 to 60  Is your el slightly feels w hot to t	sually <b>REUS</b> uffer? ow many time USE the gel for the buffer ning buffer  I, how long don 20 mins o mins cettrophoresis above room	E the electrophoresis gel  □ NO nes?  running buffer, how do you evaporation? □ add DI water  loes it take you to run a gel □ 20 to 40 mins □ Other: □ system hot after a run? temperature 30 – 45 °C)
Pleas 1) 2) 3)	Se provide answers to the You are working with   □ Protein   □ RNA  When running electro   □ 0.5 X TAE buffer   □ 0.5 X TBE buffer   □ Other:   □ Do you make your ow running buffer?   □ YES   If NO, where do you be   What stock concentrate electrophoresis gel run   □ 5 X   □ 40 X   □ Other:   □ O	□ DNA □ Other:  phoresis gels, you use □ 1 X TAE buffer □ 1 X TBE buffer  In electrophoresis gel □ NO Douy from?  tion is your Inning buffer? □ 10 X □ 50 X	<ul><li>5)</li><li>6)</li><li>7)</li></ul>	Do you urunning by YES If YES, h  If you RE make up add run Other:  In genera less that 41 to 60  Is your el slightly feels w hot to t	sually REUS uffer? ow many time USE the gel for the buffer ning buffer l, how long den 20 mins o mins ectrophoresis above room arm ( around ouch (above	E the electrophoresis gel  □ NO nes?  running buffer, how do your evaporation? □ add DI water  loes it take you to run a gel □ 20 to 40 mins □ Other: □ system hot after a run? temperature 30 – 45 °C)
Pleas 1) 2) 3)	Se provide answers to the You are working with   □ Protein   □ RNA    When running electro   □ 0.5 X TAE buffer   □ 0.5 X TBE buffer   □ Other:   □ Do you make your ow running buffer?   □ YES   If NO, where do you be   What stock concentrate electrophoresis gel run   □ 5 X   □ 40 X   □ Other:   □ OT Internal Use On	□ DNA □ Other:  phoresis gels, you use □ 1 X TAE buffer □ 1 X TBE buffer  In electrophoresis gel □ NO Douy from?  tion is your Inning buffer? □ 10 X □ 50 X	<ul><li>5)</li><li>6)</li><li>7)</li></ul>	Do you urunning by YES If YES, h  If you RE make up add run Other:  In genera less that 41 to 60  Is your el slightly feels who to	sually REUS uffer? ow many time USE the gel for the buffer ning buffer l, how long den 20 mins o mins ectrophoresis above room arm ( around ouch (above	□ NO nes? running buffer, how do you revaporation? □ add DI water  loes it take you to run a gel' □ 20 to 40 mins □ Other: system hot after a run? temperature 30 – 45 °C)
Pleas 1) 2) 3) 4)	Se provide answers to the You are working with   □ Protein   □ RNA    When running electro   □ 0.5 X TAE buffer   □ 0.5 X TBE buffer   □ Other:   □ Do you make your ow running buffer?   □ YES   If NO, where do you be   What stock concentrate electrophoresis gel run   □ 5 X   □ 40 X   □ Other:   □ OT Internal Use On	□ DNA □ Other:  phoresis gels, you use □ 1 X TAE buffer □ 1 X TBE buffer  /n electrophoresis gel □ NO  buy from?  tion is your  nning buffer? □ 10 X □ 50 X	<ul><li>5)</li><li>6)</li><li>7)</li></ul>	Do you urunning b  YES If YES, h  If you RE make up add run Other: In genera less tha 41 to 60  Is your el slightly feels w hot to t Other:	sually REUS uffer? ow many time USE the gel for the buffer ning buffer I, how long den 20 mins mins ectrophoresis above room arm ( around ouch (above	E the electrophoresis gel  □ NO les?  running buffer, how do you revaporation? □ add DI water  loes it take you to run a gel □ 20 to 40 mins □ Other:  s system hot after a run? temperature 30 – 45 °C) ~ 48°C)

Replacement Send Date (if any):