Section I – Analysis of Inorganic Compounds

Use the following information to answer Questions 16 and 17

An atomic absorption spectrometer can be used to determine the level of copper in soils. The calibration curve below plots the absorbance of four standard copper solutions against the concentration of copper ions in ppm.

The concentrations of copper ions in the standard solutions were 1.0, 2.0, 3.0 and 4.0 mg L^{-1} . (1 mg $L^{-1} = 1$ ppm)



Copper calibration curve

VCE 2019 Q16

1.

The concentration of copper in a test solution can be determined most accurately from the calibration curve if it is between

- A. 0.0 ppm and 5.0 ppm.
- B. 0.0 ppm and 4.0 ppm.
- C. 1.0 ppm and 4.0 ppm.
- D. 1.0 ppm and 5.0 ppm.

2. VCE 2019 Q17

If the test solution gave an absorbance reading of 0.40, what would be the concentration of copper ions in the solution in mol L^{-1} ?

A. 2.5 B. 3.9×10^{-2} C. 3.9×10^{-5} D. 2.5×10^{-6}

3. VCE 2013 Q8

A forensic chemist tests mud from a crime scene to determine whether the mud contains zinc.

Which one of the following analytical techniques would be best suited to this task?

- A. infrared spectroscopy
- B. mass spectrometry
- C. atomic absorption spectroscopy
- D. nuclear magnetic resonance spectroscopy

4. VCE 2011 Q20

The amount of copper in a solution of copper(II) sulfate can be determined using atomic absorption spectroscopy.

When a blue copper(II) sulfate solution is introduced into an atomic absorption spectrometer, a green flame is observed.

Consider the following statements

- I. A copper (II) sulfate solution appears blue because it absorbs red light.
- II. The metal species undergoes oxidation in the flame.
- III. The flame is green due to electron transitions from a higher energy state to a lower energy state.

Which of the above statements are true?

A. I only

- B. I and III only
- C. II and III only
- D. I, II and III

5. VCE 2010 Q9

The graph shows the absorption spectra of three food dyes: Blue No. 1, Red No. 2 and Yellow No. 4.



Which one of the following is the best wavelength to determine the concentration of Red No. 2 dye in a solution containing a mixture of all three dyes?

- A. 430 nm B. 500 nm
- C. 540 nm
- D. 620 nm

VCE 2012 Question 6 (7 marks)

The iron content in multivitamin tablets was determined using atomic absorption spectroscopy. The absorbances of four standards were measured.

Three multivitamin tablets were selected. Each tablet was dissolved in 100.0 mL of water. The absorbance of each of the three solutions was then measured.

The following absorbances were obtained.

Solution	Concentration mg/L	Absorbance
Standard 1	0.00	0.06
Standard 2	100.0	0.16
Standard 3	200.0	0.25
Standard 4	300.0	0.36
Standard 5	400.0	0.46
Tablet 1	_	0.39
Tablet 2	_	0.42
Tablet 3	_	0.45

(a)

(i) Use the grid below to construct a calibration graph of the absorbances of the standard solutions.



VCE 2012 Question 6 (continued)	
(ii) Determine the average iron content, in milligrams, of the multivitamin tablets.	2
Spectroscopic techniques work on the principle that, under certain conditions, atoms, molecules or ions will interact with electromagnetic radiation. The type of interaction depends on the wavelength of the electromagnetic radiation.	
(b) Name one spectroscopic technique that you have studied this year.	
(i) Which part of the electromagnetic spectrum does this technique use?	1
(ii) How does this part of the electromagnetic spectrum interact with matter? What information does this spectroscopic technique provide?	2

Section II – Analysis of Organic Compounds

0

1. VCE 2019 Q3

A compound has the following skeletal formula

The compound is analysed using mass spectrometry.

The molecular ion peak of the compound is at

- A. m/z = 71
- B. m/z = 74
- C. m/z = 85D. m/z = 86

2. VCE 2019 Q27

An organic compound has a molar mass of 88 g mol⁻¹. The ¹³C NMR spectrum of the organic compound shows four distinct peaks. The organic compound is most likely

- A. butan-1-ol.
- B. 2-methyl-butan-1-ol.
- C. 2-methyl-butan-2-ol.
- D. 2,2-dimethyl-propan-1-ol.

3.

VCE 2019 NHT Q10

A researcher uses a combination of spectroscopic techniques to determine the structure of a molecule. Which combination of spectroscopic techniques provides the most information about the molecule's functional groups and number of carbon environments?

А.	Mass spectrometry	¹³ C NMR
B.	Infrared	¹³ C NMR
C.	Infrared	Mass spectrometry
D.	Mass spectrometry	¹ H NMR

VCE 2017 NHT Q3 (6 marks)

A compound has the molecular formula $C_5H_{10}O_2$. It is analysed by infrared (IR), ¹³C NMR and ¹H NMR spectroscopy. The resulting spectra are shown below.



Data: SDBS Web, http://sdbs.db.aist.go.jp, National Institute of Advanced Industrial Science and Technology

VCE Past Exam Questions for Module 8: Applying Chemical Ideas SAMPLE

VCE 2017 NHT Q3 (continued)

Information for the splitting pattern of the ¹H NMR spectrum is shown below

	Chemical shift	Relative peak area	Splitting pattern	
	4.1	2	4 peaks	
	2.3	2	4 peaks	
	1.3	3	3 peaks	
	1.1	3	3 peaks	
) Usi nun	ng the IR spectrum pronter.	ovided, identify a bon	d in a molecule of C	$H_{10}O_2$ and give its wave
) Usi mo	ng the ¹ H NMR spectr lecule of $C_5H_{10}O_2$.	rum provided, state th	e number of hydroge	n environments in a
) Wh spli	at specific information itting pattern in the ¹ H	n about the structure o NMR spectrum?	f a molecule of C₅H₁	$_{0}O_{2}$ is provided by the
) Wh spli) Dra stru con	hat specific information itting pattern in the ¹ H	n about the structure of NMR spectrum?	f a molecule of C ₅ H ₁ ded for a molecule of nce from the ¹³ C NM	₀ O ₂ is provided by the
) Wh spli) Dra stru con	hat specific information itting pattern in the ¹ H	n about the structure of NMR spectrum?	f a molecule of C ₅ H ₁ ded for a molecule of nce from the ¹³ C NM	₀ O ₂ is provided by the