

## Drug Facts

<b>Active ingredients (in each tablet)</b>	<b>Purpose</b>
--	----------------

Aspirin 500 mg (NSAID*).....	Pain reliever
Caffeine 60 mg .....	Alertness aid/ Pain reliever aid

\*Nonsteroidal anti-inflammatory drug

### Uses

- for the temporary relief of minor aches and pains associated with a hangover
- helps restore mental alertness or wakefulness when experiencing fatigue or drowsiness associated with a hangover
- also for the temporary relief of headaches or body aches and pains alone

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

**Stomach bleeding warning:** This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing

## Drug Facts (continued)

prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

**Do not use** if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer

### Ask a doctor before use if

• stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have a sodium-restricted diet

### Ask a doctor or pharmacist before use if you are

taking a prescription drug for • diabetes • gout • arthritis

### Caffeine warning

• the recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heart beat. • for occasional use only.

## Drug Facts (continued)

### Stop use and ask a doctor if

• an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain lasts for more than 10 days or gets worse • new symptoms occur • redness or swelling is present • ringing in the ears or a loss of hearing occurs

### If pregnant or breastfeeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

• Upon waking, fully dissolve 1 or 2 tablets in 16 oz. of water and drink

• Adults and children 12 years and over: 1 or 2 tablets every 6 hours, as needed. Do not exceed 8 tablets in 24 hours. • Children under 12 years: do not use

## Drug Facts (continued)

### Other Information

• **each tablet contains:** sodium 406mg

• phenylketonurics: contains phenylalanine 12.6 mg per tablet

• store at room temperature (59-86°F). Protect from excessive heat.

### Inactive Ingredients

acesulfame potassium, anhydrous citric acid, aspartame, docusate sodium, flavors, mannitol, povidone, sodium benzoate, sodium bicarbonate

### Questions or Comments?

1-800-970-1793 (weekdays)

DO NOT USE IF PRINTED PACKETS  
ARE TORN OR PUNCTURED

forhangovers.com