

2016-2322
2016-05-13

PRO®
PROFESSIONAL ANT AND ROACH PYRODUST

INSECTICIDE FOR HOME AND GARDEN

READY TO USE

DOMESTIC

REG. NO. 13074 P.C.P. ACT

GUARANTEE:

Pyrethrins	0.2%
Piperonyl Butoxide	1.0%

READ THE LABEL BEFORE USING
KEEP OUT OF REACH OF CHILDREN

200 g – 600 g

PREMIER TECH BRIGHTON LTD.
1 avenue Premier
Rivière-du-Loup QC G5R 6C1
1-844-475-1262

PRO[®] Professional Ant and Roach Pyrodust is made from an African flower (*Chrysanthemum cinerariaefolium*) and has a very low mammalian toxicity.

Effective against ants, cockroaches, silverfish, spiders, earwigs, sowbugs, crickets, certain beetles, boxelder bugs, fleas, bed bugs, chewing and sucking insects such as leaf hoppers, exposed thrips and certain aphids in gardens and houses.

DIRECTIONS FOR USE:

DIRECTIONS FOR HOUSES:

Lightly coat the areas where the insects crawl and hide such as cracks and crevices, underneath built-ins, around pipes and drains, in storage cabinets, behind medicine cabinets, in electrical outlet boxes, garbage cans and in attics and basements. Repeat as necessary, probably once a month. Contact insects directly with the dust when possible. Do not dust pets. Bed bugs: dust thoroughly both sides of mattresses and beds. Fleas: thoroughly dust sleeping quarters of pets and floors.

DIRECTIONS FOR GARDENS:

Ornamental and Flower Gardens: To control Aphids, Caterpillars, Flies, Leafhoppers, Leafrollers and Thrips (exposed), lightly apply the insecticide dust when plants are dew covered or previously moistened with a spray of water. Coat both upper and lower surfaces of leaves. Repeat weekly. For the protection of bees, early or late dusting hours (while blossoms are closed) are recommended. For best results, apply in evening or cloudy day when air is still. Do not apply in direct sunlight or when temperature exceeds 32°C. Contact insects directly with the dust.

DIRECTIONS FOR USE FOR VEGETABLES: Dust the foliage thoroughly; repeat as necessary. Do not apply if rain is expected.

BEANS: To control Bean aphids, Mexican bean beetle larvae and Potato leafhoppers, use 325 g per 100 m when the plants are 25 cm high.

CABBAGE: To control Cabbage loopers, use 575 g per 100 m.

OTHER COLE CROPS: To control imported Cabbage worms, Cabbage bugs, Diamondback larvae and Harlequin bugs, use 325 g per 100 m.

CUCUMBERS: To control Aphids and Spotted cucumber beetles, use 325 g per 100 m when the plants begin to send out runners.

PRECAUTIONS: KEEP OUT OF REACH OF CHILDREN. Do not use near fish ponds or aquariums. Avoid inhalation of the dust. Wash hands and other exposed skin after handling.

FIRST AID: If swallowed, call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person. **If in eyes,** hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice. **If on skin or clothing,** take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Skin contact and inhalation may be allergenic to sensitive people.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

PRO[®] Agrium Inc. / Lic. Premier Tech Brighton Ltd.