





Preparation Instructions

As powdered formula is not sterile, failure to follow preparation and storage instructions may make your baby ill. First, thoroughly wash hands and sterilize all utensils according to the instructions below. The ingredients and low-temperature processing may give the powder a characteristic colour different from other infant formulas. For better solubility, boil water to 40°C / 104°F, then add the formula powder. Strictly follow all preparation steps and hygiene measures for bottle feeding. When opening for the first time, completely remove the metal lid and always close the can tightly after each use. Do not prepare bottles in advance. After your baby's feeding, do not keep the remaining formula for more than an hour and carefully clean the containers and accessories. For making bottle feeds, it is preferable to choose a glass bottle and low-mineral water, which are particularly suitable for feeding infants. Do not heat the bottle or the water used to prepare it in the microwave. Keep the can closed tightly in a clean, cool and dry place away from light and use the powder within 3 weeks after opening. This product is not suitable for parenteral use or tube feeding.



- 1. Boil fresh tap water. Leave to cool for no longer than 30 minutes. Measure the amount of water required (see feeding guide) into a sterilized bottle
- 2. Using the scoop provided, add the correct number of levelled scoops according to the feeding guide,
- 3. Carefully close the bottle. Shake the bottle vigorously, circularly, then vertically until the powder is completely dissolved.
- 4. Before giving the bottle, be sure to check the temperature by pouring a few drops onto the inside for your wrist. The temperature should be lukewarm (approximately 37 °C / 122 °F). Close the can tightly.

Amounts and Dosage

Bébé M recommends a single hole fast flow teat. The volume and number of bottles are given as guidelines. A healthcare professional may be able to adapt these recommendations to fit your child's specific needs. 1 level scoop (4g) for 30ml of water. Only use the scoop provided in the can.

Baby's Age (In Months)		10 - 12	12 - 36
Bottle Preparation	Water (ml)	210	240
	Scoops	7	8
Number of Bottles Per 24H		3	2 - 3

Important Notes

Breast milk is best for babies. The choice of breastfeeding must be made as a priority both for your child and for your health. Bébé M Stage 3 can only be used from 10 months of age, according to the advice of your healthcare professional. It must be fed under medical supervision and only in infants with the symptoms indicated. The introduction of complementary foods, including any exceptions before 6 months of age, should only be made on the advice of a healthcare professional, based on the specific needs of each infant in terms of growth and development. Do not change infant formula without medical advice. Bébé M Stage 3 is intended for special medical purposes. It meets the nutritional needs of infants who have reached the age of at least 10 months, in case of allergies to cow's milk proteins and/or lactose intolerance and in the case of regurgitation. It can only be used as part of a diversified diet. It cannot be used as a substitute for breast milk during the first 6 months of life. It is lactose-free. It does not contain gluten in accordance with regulations. When bottle feeding, do not allow prolonged or frequent contact of milk feeds with teeth as this increases the risk of tooth decay. Make sure your baby's teeth are cleaned before the last feed at night.

Bébé M Stage 3