



Preparation Instructions

Check that the closure is intact and in perfect condition before use. Shake the still-sealed bottle before opening it for the first time. Pour the required amount of formula into a bottle and warm it up to drinking temperature (approximately 37° C / 98.6° F.). Do not heat the formula in the microwave due to the risk of scalding. Recheck the temperature before feeding. Feed the formula immediately, and do not reuse leftovers. Store the opened, unheated bottle in the refrigerator (4-6 °C / 39.2-42.8 °F) and use within 24 hours.

Amounts and Dosage

Recommended drinking quantities and the number of bottles per day are for guidance only. Your baby does not necessarily have to drink the bottle completely.

| Age | Bottles Per Day | Prepared Formula (ml) |
|---------------|-----------------|-----------------------|
| 1 - 2 weeks * | 5 - 7 | 70 ml |
| 3 - 4 weeks | 5 - 7 | 100 ml |
| 5 - 8 weeks | 5 - 6 | 135 ml |
| 3 - 4 months | 5 | 170 ml |
| From 5 months | 4 - 5 | 200 ml |
| From 6 months | 4 | 235 ml |

^{*}Unless otherwise recommended by your doctor.

Important Notes

Breastfeeding is the best thing for your baby. Please speak to your pediatrician if you would like to use formula. For your baby's dental health: Like breast milk, all formulas contain carbohydrates. They are an important part of your baby's diet. Frequent or prolonged contact of the teeth with liquids containing carbohydrates can lead to tooth decay and, thus, to health problems. Do not allow your child to continuously suck on the bottle, and switch to cup feeding as early as possible. HiPP HA Ready-to-Feed Stage PRE is suitable from birth as the sole source of nutrition or as a supplement to breast milk. Like breast milk, this formula can be given as often and as much as your baby wants to drink. HiPP HA must not be used if your baby has a diagnosed or suspected allergy to cow's milk proteins.

HiPP HA Ready-to-Feed Stage PRE