

## **SUGGESTED FORMULA**

\_\_\_\_\_

Clonazepam 5mg/mL Oral Suspension

Version number: 1.0 Volume: 100 mL

Clonazepam 10mg Vehicle for Oral Solution, NF (Ora-Sweet  $^{TM}$ ) 50mL

Vehicle for Oral Suspension, NF (Ora-Plus<sup>™</sup>) 100mL Q.S.

SUGGESTED COMPOUNDING PROCEDURES

1. Weigh Clonazepam and place in glass mortar and pestle

- 2. Dry triturate to break down any clumps in the chemical
- 3. Add enough of liquid into mortar and pestle to form a "paste"
- 4. Triturate until all Clonazepam is broken down and incorporated into liquid
- 5. Add more liquid until pourable mixture and pour into calibrated final container
- 6 Perform liquid "washes" in mortar and pestle to integrate remaining chemical
- 7. Pour each "wash" into final container taking precaution not to go over the final volume
- 8. Q.S. medication and shake well.
- 9. Suggested Quality Assessments follow pharmacy SOPs:
  - a. Weight to Volume calculation
  - b.Color
  - c.Pourability
  - d.Settling
  - e.Resuspendability

Store in air tight amber plastic containers

Store at Room Temperature or Refrigerated

No claims are made as to the safety or efficacy of this preparation. This formulation is provided solely at the unsolicited request of the pharmacist.

Beyond-Use Dates of preparations are conservative estimates by the formulator using reference books, peer-reviewed literature, intended duration of therapy, formulation from commercially available products, organoleptic observations and current USP guidelines. Compounders may have stability studies performed by a reputable laboratory if they wish to extend the Beyond-Use Date.

It is recommended that you follow USP <795> recommendations for potency testing.

**Beyond-Use Date is estimated to be <u>60 Days</u>** Per *Trissel's Stability of Compounded Formulations 5<sup>th</sup> Edition and Allen Loyd V Jr. Clonazepam Suspension. US Pharmacist 20(2);1995:84-85.* 

Precautions should be taken to prevent cross-contamination and exposure of ingredients to the compounder and contamination of the preparation by the compounder. Wear appropriate protective equipment. Use safety enclosures (hoods) when weighing and mixing.

06/15 RD

