



ARBOR ASSAYS
Interactive Assay Solutions™

OXIDATIVE
STRESS

Table of Contents

DetectX® Assay Kits	Page
Catalase Colorimetric & Fluorescent Activity Kits	4
Ceruloplasmin (Cp) Colorimetric Activity Kit	5
DNA Damage ELISA Kits	6
Formaldehyde Fluorescent Detection Kit	7
FRAP™ (Ferric Reducing Antioxidant Power) Detection Kit	8
Glutathione Colorimetric Detection Kits	9
Glutathione Fluorescent Detection Kit	10
Glutathione Reductase (GR) Fluorescent Activity Kit	11
Glutathione S-Transferase (GST) Fluorescent Activity Kit	12
Hemoglobin Colorimetric Detection Kits	13
Hydrogen Peroxide (H ₂ O ₂) Colorimetric & Fluorescent Detection Kits	14
Myeloperoxidase Human ELISA Kit	15
Nitric Oxide (NO ₂ ⁻ /NO ₃ ⁻) Colorimetric Detection Kit	16
Superoxide Dismutase (SOD) Colorimetric Activity Kit	17
TBARS / MDA Universal Colorimetric Detection Kit	18
Thiol Fluorescent Detection Kit	19



ONLINE:

www.ArborAssays.com/order-form

PHONE:

Call 734-677-1774 or Toll Free: 855-677-1774. Monday-Friday 8:30 am to 5:30 pm, EST.

FAX:

Send faxes to 734-677-6860.

E-MAIL:

Send E-mail orders to Orders@ArborAssays.com

DISTRIBUTORS:

Check our website at www.ArborAssays.com/distributors for a list of distributors.



Catalase Colorimetric & Fluorescent Activity Kits

Colorimetric: K033-H1 (2 Plate)

Fluorescent: K033-F1 (2 Plate)

FEATURES


- ▶ Use Measure Catalase Activity in Any Sample
- ▶ Time to Answer 45 Minutes
- ▶ Sensitivity Measure as little as 0.052 U/mL
- ▶ Samples/Kit 89 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Format 96-well
- ▶ Species Species Independent
- ▶ Readout Colorimetric: 560 nm
Fluorescent: 590 nm em/520 nm ex




SCIENTIFIC RELEVANCE

Hydrogen peroxide, (H₂O₂) is one of the most frequently occurring reactive oxygen species. It is formed either in the environment, as a by-product of aerobic metabolism, superoxide formation and dismutation, or as a product of oxidase activity. Both excessive hydrogen peroxide and its decomposition product hydroxyl radical, are harmful for most cell components. Its rapid removal is essential for all aerobically living prokaryotic and eukaryotic cells.

Fluorescent



Colorimetric



One of the most efficient ways of removing peroxide is through the enzyme catalase, which is encoded by a single gene and is highly conserved among species. Mammals, including humans and mice, express catalase in all tissues. A high concentration of catalase can be found in the liver, kidneys and erythrocytes. The expression is regulated at transcription, post-transcription and post-translation levels. High catalase activity is detected in peroxisomes.



Ceruloplasmin (Cp) Colorimetric Activity Kit

K035-H1 (2 Plate)


FEATURES

- ▶ Use Non-Invasive Pregnancy Marker
- ▶ Sample Urine, Serum
- ▶ Validation Humans, Felids, Polar Bear, Panda
- ▶ Species Multiple Species
- ▶ Time to Answer 60 Minutes
- ▶ Format 96-Well
- ▶ Samples/Kit 89 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Colorimetric, 560 nm



SCIENTIFIC RELEVANCE

Ceruloplasmin (Cp) is an acute phase multicopper oxidase enzyme that normally plays a protective role in responses to immune-provoking stimuli and is also associated with reproduction. Estrogens alter the subcellular distribution of copper in the liver, leading to an increase in plasma copper levels and subsequent ceruloplasmin synthesis. Serum levels of Cp have been shown to increase during normal pregnancy in some species, possibly as a protection against the oxidative costs of reproduction. In giant pandas and some felids, urinary Cp activity has shown to be elevated in pregnant vs. pseudopregnant animals beginning in the first week of gestation and continuing throughout the luteal phase.



DNA Damage ELISA Kits

K059-H1 (1 Plate) | K059-H5 (5 Plate)


FEATURES

- ▶ Use Measure oxidized guanine molecules
- ▶ Sample Serum, Plasma, Saliva, Urine, Fecal Extracts, Digested DNA, TCM
- ▶ Time to Answer 2.5 hours
- ▶ Sensitivity 51 pg/mL
- ▶ Species Species independent
- ▶ Samples/Kit 38 or 230 in Duplicate
- ▶ Readout Colorimetric, 450 nm



SCIENTIFIC RELEVANCE

Free radicals and other reactive species are constantly generated *in vivo* and cause oxidative damage to biomolecules, a process held in check only by the existence of multiple antioxidant and repair systems as well as the replacement of damaged nucleic acids, proteins and lipids. Intracellular reactive oxygen species are produced as a result of normal metabolism and extracellular forms are produced as a result of ultraviolet or ionizing radiation. Cellular function may be interrupted or halted if DNA damage corrupts the integrity of essential information contained in the genome and it is widely thought that continuous oxidative damage to DNA is a significant contributor to the age-related development of major cancers. Among numerous types of oxidative DNA damage, the formation of 8-hydroxy-2'-deoxyguanosine (8-OHdG) is a ubiquitous marker of oxidative stress. 8-OHdG is physiologically formed and enhanced by chemical carcinogens. During the repair of damaged DNA *in vivo* by exonucleases the resulting 8-OHdG is excreted without further metabolism into urine, allowing for convenient detection.



Formaldehyde Fluorescent Detection Kit

K001-F1 (2 Plate)


FEATURES

- ▶ Use Measure Formaldehyde
- ▶ Sample Urine, Water, TCM
- ▶ Convenient No Extraction, No Chemical Derivatization
- ▶ Samples/Kit 88 in Duplicate
- ▶ Sensitivity < 80 pM
- ▶ Stability Liquid 4° Stable Reagents
- ▶ Time to Answer 30 Minutes
- ▶ Readout Fluorescent, 510 nm em/450 nm ex



SCIENTIFIC RELEVANCE

Formaldehyde (methanal), $\text{H}_2\text{C}=\text{O}$, is a colorless, flammable, strong-smelling gas. It is an important industrial chemical used to manufacture building materials and to produce many household products. In the US approximately 3×10^9 Kg are produced annually. Formaldehyde is commonly used as an industrial fungicide, germicide, and disinfectant, and as a preservative in mortuaries and medical laboratories. Materials containing formaldehyde can release formaldehyde gas or vapor into the air. Formaldehyde can also be released by burning wood, kerosene, natural gas, or cigarettes, from automobile emissions, and from natural processes. Occupational exposure to formaldehyde by inhalation is mainly from three types of sources: thermal or chemical decomposition of formaldehyde-based resins, formaldehyde emission from aqueous solutions (for example, embalming fluids), and the production of formaldehyde resulting from combustion. Formaldehyde can be toxic, allergenic, and carcinogenic. Because formaldehyde resins are used in many construction materials, it is one of the more common indoor air pollutants.



FRAP™ (Ferric Reducing Antioxidant Power) Detection Kit

K043-H1 (2 Plate)


FEATURES

- ▶ Use Measure Ferric Reducing Anti-Oxidant Potential (FRAP) of Samples
- ▶ Samples Serum, Plasma, Tissue, Saliva, Cell Lysates, Urine, Food, Cosmetics, Additives
- ▶ Samples/Kit 89 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Time to Answer 30 Minutes
- ▶ Readout Colorimetric, 560 nm



SCIENTIFIC RELEVANCE

Potentially harmful reactive oxygen species (ROS) are produced as a consequence of normal aerobic metabolism. "Free Radicals" (FR) are usually removed or inactivated in vivo by a team of antioxidants. They are chemically stable atoms and molecules, which have one or more free electrons. Almost all biomolecules may be attacked by reactive free radicals. Free radicals are responsible for many pathological processes, or they can be generated as the result of the pathological stage and cause important secondary damage to biological systems and cells. Connections between free radicals and some serious diseases, including Parkinson's and Alzheimer's diseases, atherosclerosis, heart attacks, and chronic fatigue syndrome, have been demonstrated. However, short-term oxidative stress, the unbalance between the formation and scavenging of the reactive oxygen species, may be important in the prevention of aging due to triggering of the process known as mitohormesis. On average, 65 - 70% of the population is excessively impacted by oxidative stress caused by FRs.



Glutathione (GSH) Colorimetric Detection Kits

K006-H1 (4 Plate)

K006-H1C-H/L (200 Cuvette)



 MULTI
SPECIES

FEATURES

- ▶ Use Measure Total GSH and GSSG to Determine Oxidative Stress
- ▶ Samples Whole Blood, Serum, Plasma, Erythrocytes, Urine, Lysates, TCM
- ▶ Sensitivity 0.634 μM (Plate-based Format)
- ▶ Format 96-Well or Cuvette
- ▶ Species Species Independent
- ▶ Samples/Kit K006-H1: 89 (Total and GSSG) in Duplicate
K006-H1C-H/L: 43 (Total and GSSG) in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Colorimetric, 405 nm

SCIENTIFIC RELEVANCE

Glutathione (L- γ -glutamyl-L-cysteinylglycine; GSH) is the highest concentration non-protein thiol in mammalian cells and is present in concentrations of 0.5 – 10 mM. GSH is a tripeptide that contains an unusual peptide linkage between the amine group of cysteine and the carboxyl group of the glutamate side-chain. It is an antioxidant, preventing damage to important cellular components caused by reactive oxygen species such as free radicals and peroxides. Glutathione reduces disulfide bonds formed within cytoplasmic proteins to cysteines by serving as an electron donor. In the process, glutathione is converted to its oxidized form, glutathione disulfide (GSSG). Glutathione is found mostly in its reduced form since the enzyme that reverts it from its oxidized form, glutathione reductase, is constitutive and inducible upon oxidative stress. The ratio of reduced glutathione to oxidized glutathione within cells is often used as a measure of cellular toxicity.



Glutathione (GSH) Fluorescent Detection Kits

K006-F1 (1 Plate) | K006-F5 (5 Plate)
K006-F1D (384-Well Plate)

FEATURES


- ▶ Use: Measure GSH/GSSG to Determine Oxidative Stress
- ▶ Sample: Whole Blood, Serum, Plasma, Erythrocytes, Urine, Lysates, TCM
- ▶ Species: Species Independent
- ▶ Sensitivity: 45 nM Free GSH, 48 nM Total GSH
- ▶ Samples/Kit: 39 or 231 in Duplicate (K006-F1/F5)
183 in Duplicate (K006-F1D)
- ▶ Stability: Liquid 4°C Stable Reagents
- ▶ Readout: Fluorescent, 510 nm em/370-410 nm ex




SCIENTIFIC RELEVANCE

Glutathione (L-γ-glutamyl-L-cysteinylglycine; GSH) is the highest concentration non-protein thiol in mammalian cells and is present in concentrations of 0.5 – 10 mM. GSH is a tripeptide that contains an unusual peptide linkage between the amine group of cysteine and the carboxyl group of the glutamate side-chain. It is an antioxidant, preventing damage to important cellular components caused by reactive oxygen species, such as free radicals and peroxides. Glutathione reduces disulfide bonds formed within cytoplasmic proteins to cysteines by serving as an electron donor. In the process, glutathione is converted to its oxidized form, glutathione disulfide (GSSG). Glutathione is found mostly in its reduced form, since the enzyme that reverts it from its oxidized form, glutathione reductase, is constitutive and inducible upon oxidative stress. The ratio of reduced glutathione to oxidized glutathione within cells is often used as a measure of cellular toxicity.

K006-F1/F5



K006-F1D



Glutathione Reductase Fluorescent Activity Kit

K009-F1 (1 Plate)


FEATURES

- ▶ Use Measure GR activity
- ▶ Sample RBCs, Serum, Plasma, Cells, Tissues
- ▶ Convenient 20 minute End Point or Kinetic Assay
- ▶ Sensitivity 9 μ U/mL, World's Most Sensitive
- ▶ Species Species Independent
- ▶ Samples/Kit 41 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Fluorescent, 510 nm em/370-410 nm ex



SCIENTIFIC RELEVANCE

Glutathione reductase (GR) plays an indirect but essential role in the prevention of oxidative damage within the cell by helping to maintain appropriate levels of intracellular glutathione (GSH). GSH, in conjunction with the enzyme glutathione peroxidase (GP), is the acting reductant responsible for minimizing harmful hydrogen peroxide. The regeneration of GSH is catalyzed by GR. GR is a ubiquitous 100-120 kDa dimeric flavoprotein that catalyzes the reduction of oxidized glutathione (GSSG) to reduced glutathione, using β -nicotinamide dinucleotide phosphate (NADPH) as the hydrogen donor. NADPH has been suggested to also act as an indirectly operating antioxidant, given its role in the recycling of GSSG to GSH and thus maintaining the antioxidative power of glutathione.



Glutathione S-Transferase Fluorescent Activity Kit

K008-F1 (1 Plate)


FEATURES

- ▶ Use Measure GST Activity
- ▶ Sample Serum, Plasma, Urine, Cell Lysates
- ▶ Samples/Kit 40 in Duplicate
- ▶ Convenient 30 Minute End Point or Kinetic Assay
- ▶ Sensitivity < 100 μ U of GST Activity
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Fluorescent, 460 nm em/370-410 nm ex



SCIENTIFIC RELEVANCE

The Glutathione S-Transferase (GST) family of isozymes function to detoxify and neutralize a wide variety of electrophilic molecules by mediating their conjugation with reduced glutathione. Human GSTs are encoded by five gene families, expressing in almost all tissues as four cytosolic and one microsomal forms. Given its pivotal role in ameliorating oxidative stress/damage, GST activity has been repeatedly investigated as a biomarker for arthritis, asthma, COPD, and multiple forms of cancer, as well as an environmental marker. Examination of GST isoforms and activity in human cancers, tumors and tumor cell lines has revealed the predominance of the acidic pi class. Furthermore, this activity is thought to substantially contribute to the innate or acquired resistance of specific neoplasms to anticancer therapy.



Hemoglobin Colorimetric Detection Kits

Regular: K013-H1 (2 Plate)

High Sensitivity: K013-HX1 (2 Plate) | K013-HX5 (10 Plate)

FEATURES


- ▶ Sample Type K013-H: Whole Blood, RBCs
 K013-HX: Serum, Plasma
- ▶ Time to Answer 30 Minutes
- ▶ Range K013-H: 16-0.25 g/dL
 K013-HX: 20-0.313 µg/mL
- ▶ Sensitivity K013-H: 0.021 g/dL, 0.21 mg/mL
 K013-HX1: 0.053 ug/mL
- ▶ Samples/Kit 88 in Duplicate
- ▶ Stable Liquid 4°C Stable Reagents
- ▶ Readout K013-H: 560-580 nm
 K013-HX: 450 nm




SCIENTIFIC RELEVANCE

Hemoglobin (Hgb) is an erythrocyte protein complex comprised of two sets of identical pairs of subunits, each of which bind an iron-porphyrin group commonly called heme. Heme binds and releases oxygen or carbon dioxide in response to slight changes in local gas tension. Hemoglobin values are associated with a variety of conditions ranging from anemias (low Hgb), erythrocytosis (high Hgb), thalassemias (aberrant chain synthesis), and sickling disorders (abnormal complex shape).

K013-H1



K013-HX1/HX5



Hydrogen Peroxide Colorimetric & Fluorescent Detection Kits

Colorimetric: K034-H1 (2 Plate)

Fluorescent: K034-F1 (2 Plate)


FEATURES

- ▶ Use Measure H₂O₂ in Any Sample
- ▶ Sample Urine, Buffer, TCM
- ▶ Rapid 15 Minutes
- ▶ Sensitive Colorimetric: 91.3 pmol (310 ng)
Fluorescent: < 2 pmole (65 pg)
- ▶ Samples/Kit Colorimetric: 89 in Duplicate
Fluorescent: 88 in Duplicate
- ▶ Readout Colorimetric: 560 nm
Fluorescent: 590 nm em/520 nm ex


SCIENTIFIC RELEVANCE

In biological systems, incomplete reduction of O₂ during respiration produces superoxide anion (O₂⁻), which is spontaneously or enzymatically dismutated by superoxide dismutase to H₂O₂. Many cells produce low levels of O₂⁻ and H₂O₂ in response to a variety of extracellular stimuli, such as cytokines (TGF-β1, TNF-α, and various interleukins), peptide growth factors (PDGF, EGF, VEGF, bFGF, and insulin), the agonists of heterotrimeric G protein-coupled receptors (GPCR) such as angiotensin II, thrombin, lysophosphatidic acid, sphingosine 1-phosphate, histamine, and bradykinin, and by shear stress. The addition of exogenous H₂O₂, or the intracellular production in response to receptor stimulation, affects the function of various proteins including protein kinases, protein phosphatases, transcription factors, phospholipases, ion channels, and G proteins. In 1894, Fenton described the oxidation of tartaric acid by Fe²⁺ and H₂O₂. H₂O₂ and O₂ may participate in the production of singlet oxygen and peroxynitrite and the generation of these species may be concurrent with reactions involving iron, which under some circumstances might be important contributors to H₂O₂ toxicity.

Colorimetric



Fluorescent



Myeloperoxidase (MPO) Human ELISA Kit

K060-H1 (1 Plate)

FEATURES

- ▶ Use Measure Human MPO in a Variety of Matrices
- ▶ Sample Serum, Platelet-Poor Heparin Plasma, Saliva, Urine, TCM
- ▶ Time to Answer 2.5 Hours
- ▶ Sensitivity 0.068 ng/mL, 68 pg/mL
- ▶ Species Human
- ▶ Samples/Kit 40 in Duplicate
- ▶ Readout Colorimetric, 450 nm




ARBOR ASSAYS™

Developed and produced in
collaboration with **Athens Research
& Technology.**

www.athensresearch.com

SCIENTIFIC RELEVANCE

Myeloperoxidase (MPO) is a tetrameric heme-containing protein abundantly produced in neutrophil granulocytes where it plays an important anti-microbial role. During degranulation MPO is released into the extracellular space. There, as part of the neutrophils “respiratory burst”, it produces hypochlorous acid from hydrogen peroxide and Cl⁻. MPO also uses hydrogen peroxide to oxidize tyrosine to the tyrosyl radical. Both hypochlorous acid and tyrosyl are cytotoxic and when present can kill bacteria and other pathogens. Hereditary deficiency of myeloperoxidase predisposes individuals to immune deficiency. Studies have shown an association between elevated MPO levels and coronary artery disease, and in 2003 it was suggested that MPO may serve as a sensitive predictor of myocardial infarction in patients complaining of chest pain. Since that time the clinical utility of MPO testing in cardiac patients has been solidly established in the literature with well over 100 papers published. In 2010 this clinical application was further refined by additional studies which determined that measuring both MPO and C-reactive protein (CRP) provided more accurate prediction of mortality risk than measuring just CRP alone.



Nitric Oxide Colorimetric Detection Kit

K023-H1 (2 Plate)


FEATURES

- ▶ Use Measure Nitrite & Nitrate
- ▶ Sample Water, Serum, Plasma, Urine, Saliva, Lysates, Buffers, TCM
- ▶ Accurate Calibrated to NIST Standard Reference Material #3185
- ▶ Sensitivity Highest Optical Density of Any Kit
- ▶ Time to Answer 5 Minute Nitrite – 25 Minute Total NO
- ▶ Samples/Kit 88 in Duplicate
- ▶ Stability Non-Toxic, Liquid 4°C Stable Reagents
- ▶ Readout Colorimetric, 550-570 nm



SCIENTIFIC RELEVANCE

Nitric oxide (NO) is a diffusible, transient, reactive molecule that has physiological effects in the pM- μ M range. Acting through guanylate cyclase activation, NO is an important regulator of the cardiovascular, nervous, and immunological systems. NO is bio-available by two routes. It can be endogenously generated by constitutive or induced NOS enzymes, or it can be ingested as nitrates or nitrites for conversion into NO. The reactive nature of nitric oxide allows it to act as a cytotoxic factor when released during an immune response by macrophages. The reactivity also allows NO to be easily converted to a toxic radical that can produce nitrosylation damage to cells and DNA. Nitrosylation can be a regulated post-translational modification in cell signaling. The dynamics of the regulatory/damage facets of NO are major forces in mitochondrial signaling and dysfunction. NO is linked not only to coronary heart disease, endothelial dysfunctions, erectile dysfunction, and neurological disorders, but also diabetes, chronic periodontitis, autism and cancer.



Superoxide Dismutase (SOD) Activity Kit

K028-H1 (2 Plate)


FEATURES

- ▶ Use Oxidative Stress Determination
- ▶ Sample Serum, Plasma, Cells, Tissue Buffers, Erythrocytes
- ▶ Species Human and Other Mammalian Species
- ▶ Samples/Kit 88 in Duplicate
- ▶ Time to Answer 20 Minutes
- ▶ Readout Colorimetric, 450 nm



SCIENTIFIC RELEVANCE

Short-lived and highly reactive oxygen species (ROS) such as $O_2^{\cdot-}$ (superoxide), $\cdot OH$ (hydroxyl radical), and H_2O_2 (hydrogen peroxide) are continuously generated *in vivo*. The cellular levels of ROS are controlled by antioxidant enzymes and small molecule antioxidants. The major antioxidant enzymes, superoxide dismutases (SODs), including copper-zinc superoxide dismutase (Cu/ZnSOD), manganese superoxide dismutase (MnSOD), and extracellular superoxide dismutase (EC-SOD). All play a critical roles in scavenging $O_2^{\cdot-}$. Decreased SOD activity results in elevated level of superoxide which in turn leads to decreased NO and increased peroxynitrite concentrations. The major intracellular SOD is a 32-kDa copper and zinc containing homodimer (Cu/Zn SOD). The mitochondrial SOD (MnSOD) is a manganese-containing 93-kDa homotetramer that is synthesized in the cytoplasm and translocated to the inner matrix of mitochondria. EC-SOD is the primary extracellular SOD enzyme and is highly expressed in many organs. Increased SOD activity levels are seen in Downs Syndrome, while decreased activity is seen in diabetes, Alzheimer's disease, rheumatoid arthritis, Parkinson's disease, uremic anemia, atherosclerosis, some cancers, and thyroid dysfunction.



TBAR MDA Universal Colorimetric Detection Kit

K077-H1 (2 Plate)


FEATURES

- ▶ Use Assess lipid peroxidation without boiling water bath treatment
- ▶ Sample Serum, Plasma, Tissue, Cell and Food Extracts, Urine, Buffers
- ▶ Sensitivity 0.36 μ Mf
- ▶ Time to Answer 1 Hour
- ▶ Format Shaking at 37°C
- ▶ Samples/Kit 88 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Colorimetric, 535 nm



SCIENTIFIC RELEVANCE

Malondialdehyde (MDA) is a 3-carbon dialdehyde formed by lipid peroxidation of polyunsaturated fatty acids by reactive oxygen species and other oxidative mechanisms. Lipid peroxidation is a well-established mechanism of cellular injury in plants and animals and is used as an indicator of oxidative stress in cells and tissues. Lipid peroxidation products derived from polyunsaturated fatty acids decompose to form a diverse mixture of compounds, including MDA. The combined total of MDA plus other reactive substances are termed TBARS for Thiobarbituric Acid Reactive Substances. Modifications of the TBARS assay have been used to evaluate several types of samples, including mammalian tissues, serum, plasma and urine along with food samples. There is some ambiguity surrounding the use of TBARS in different sample types under different oxidative stress because of the reactivity of acidified TBA toward reactive aldehydes, however the assay is used extensively to determine lipid peroxidation. In general lipids with greater unsaturation will yield higher TBARS values.



Thiol Fluorescent Detection Kits

K005-F1 (1 Plate)


FEATURES

- ▶ Use Measure Thiol Content of Proteins and Peptides
- ▶ Adaptable Measure SH easily in 8M GuHCl Buffers
- ▶ Sensitivity 4.62 nM
- ▶ Time to Answer 30 Minutes
- ▶ Species Species Independent
- ▶ Samples/Kit 39 in Duplicate
- ▶ Stability Liquid 4°C Stable Reagents
- ▶ Readout Fluorescent, 510 nm em/370-410 nm ex



SCIENTIFIC RELEVANCE

Free thiols in biological systems have important roles. Oxidatively-modified thiol groups of cysteine residues are known to modulate the activity of a growing number of proteins. As such, it is important to be able to accurately determine the extent of modification of specific amino acids, such as cysteine residues. This is especially difficult in a complex protein sample, especially in the presence of chaotropic agents such as guanidine hydrochloride. Typical methods using Ellman’s reagent do not have sufficient sensitivity to allow economical detection of free SH groups.





STRESS

REPRODUCTIVE

OXIDATIVE
STRESS



METABOLISM

VOLUME &
NORMALIZATION



KIDNEY
INJURY

INFLAMMATION



CELL
SIGNALING



ARBOR
ASSAYS™