

60, 69, 59, 61, 68 Allergerns	26 Pages:
Histamine Immune reactions to environmental elements, inhalants, epidermal and smells	
Last Updated: 11-06-18	

No matter how many allergy reactions come up in a lab test, the main reason is the immune system is weak.

There we need to work on repair and strengthening the immune defense system and less stress in your pets environment.

#### MOST IMPORTANT

When your pet has had chemical drugs - insecticide ones, vaccines or antibiotics we need to repair organs and system from the toxic insult.

Antibiotic drugs damages, in order to strengthen the immune system we need to repair the gut flora defense system again, this will also clear current or future candida fungal infection - ears, gut or skin etc

- 1. e.g detox using **HAMPL Pet Cleanse 42 30ml** (use of any chemical drugs e.g. **antibiotics, steroids and skin drugs etc**) and activated charcoal powder (set of 2)
- 2. Repair and restore the gut flora (natural defense system) with **Probiotics and Colostrum powder** if still a young pet or if they have a stool issue (bowel issue)

e.g if you have had to give the immune suppressant healing inhibitor type of drug, they won't get better, and so need continual drugs to suppress inflammation, thus leading to many many side effects and liver failure.



#### HOW TO SUPPORT AND REPAIR THE IMMUNE SYSTEM.

# 1. Vitamin C supplement

in diet will help (as vaccines severely deplete vitamin C)

#### e.g. A whole food type of "Vitamin C" powder

(a good one is **Pure Radiance Vitamin C**)

\* order from iherb.com (1 powder from capsule = 650mg).

This type of natural vitamin c is good if it's continued long-term just for good health.

#### **DOSAGE: Organic Natural Vitamin C POWDER DOSAGE:**

Small to Large Canine: 2 powders from capsules in meals or/ and milk drinks - twice daily. \* or if sick do four times a day for a week.

Puppy or Toy Dog: 1 powder from a capsule in meals or /and milk drinks - twice daily.

#### OR

If you can't quite afford the natural form - get the "Synthetic" Vitamin C crystals (white powder)

\* only use this short-term as its synthetic .. a couple of weeks to a month.

#### Synthetic Vitamin C - SODIUM ASCORBATE POWDER

<u>Small to Large Canine:</u> Add approx. 2 teaspoons in meals and milk drinks twice to three times daily or *dilute in a little liquid an oral syringe.* 

<u>Puppy or Toy Dog:</u> Add approx. 1 teaspoon in meals and milk drinks twice to three times a day or *dilute in a little liquid an oral syringe.* 

**Vitamin C** is an anti-inflammatory that helps reduce and prevent inflammation and swelling. Research has shown that high doses of vitamin C given orally or in meals has been shown to reduce the production of pro-inflammatory cytokines. We have had great improvement seen when vitamin C is restored back into the body. Example: vaccinations will severely deplete the body of Vitamin C.

Skin Conditions, weakness in joints and spine which cause "Inflammation" and "pain in joints", spine area can be often associated with a vitamin or mineral deficiencies. Thus continued vaccinations that severely deplete vitamin c and magnesium (stress), will make your dog more susceptible to injuries and tears. Research has shown that high doses of vitamin C given orally or in meals has been shown to reduce the production of pro-inflammatory cytokines. We have had great improvement seen when vitamin C is restored back into the body.



# 2. Detoxing - from chemical drugs \* See page 7.

Using Activated Charcoal powder. This powder can detox over 4,000 toxins a day.

Research studies have shown that activated charcoal decreases the toxic load on the body by over 60% with daily use. This is due to its ability to bind and grab up chlorine, pesticides and herbicides, bacteria, viruses, yeast, parasites, toxic chemicals, and endotoxic metabolites.

PUPPY give approx: 1/4 teaspoon of the powder to meals or milk drinks.

ADULT PETS: use 1 capsule of the powder or 1/2 a teaspoon in meals and/or milk drinks.

**MULTIPLE PETS** (cats and dogs), add 2 capsules or 1 teaspoon the powder into a jar with a screw lid (one or 2 cups of filtered water) and can add the homeopathic detox remedies (along with) things like the Bioplasma drops (tissue salts) is great for growing infants and run down pets and kittens who may have had parents who were mal-nutrition or feral etc, homeopathic **ParaExpel wormer** and **DrawingOut 16 drops** if teeth or skin issues. Shake before adding to each meal or/milk drinks. pour about a tablespoon or so of the liquid each time. **TWICE a day for one month** to help detox from the heavy metals from the vaccines.

After a month, one can continue adding activated charcoal to meals daily for optimum health.

\* You should be able to order online or buy from a health food store the capsules or powder.

If previously or are on chemical drugs or vaccinations (pet was unwell from) we can "assist in detoxing" as well with the below remedies and as all drugs affect the liver, we need to repair liver:

#### **HAMPL Chemical poisoning 110 30ml**

(overdosing and use of chemical sprays and drugs e.g. heartworm, worming, flea, and tick insecticide poisons)

#### **HAMPL Pet Cleanse 42 30ml**

(use of any chemical drugs - e.g. antibiotics, steroids etc)

#### **HAMPL Vaccine Detox 30ml**

(all new customers, you will receive a sample of the Vaccine Detox pills),

#### **HAMPL Rabies Vacc Detox 30ml**

(rabies injection side effects)

#### **HAMPL Liver Detox 36 30ml drops**

(all drugs affect the liver, e.g often causes a high liver enzymes reading from using these chemical drugs)

We have an alternative, safe and effective products for all species example the link to each product below: **NATURALHEARTWORM DROPS** \* download the pdf HELP SHEET as well.

https://holisticanimalremedies.com/heartworm-homeopathic-natural-treatment-and-prevention-all-species-dog.html

#### NATURAL INTESTINAL WORMER

https://holisticanimalremedies.com/worms-paraexpel-homeopathic-intestinal-worm-in-all-species-protozoan-microscopic-parasite-tritrichomonas-foetus-nosode-t-foetus.html

#### **TICK AND FLEA SPRAY**

https://holisticanimalremedies.com/canine-natural-flea-mosquito-ticks-protection-spray-topically.html

#### **HERBAL ANTIBIOTIC**

https://holisticanimalremedies.com/herbal-antibiotic-for-a-systemic-or-localised-infection-anywhere-in-body-for-cat-dog-or-other-species.html



# 3. Gut flora - defence system

We highly recommending purchasing and using the below supplements to restore and strengthen your dog or puppies "gut flora defense system"

#### **Probiotics Powder**

a good quality one is - " **ProBiotic 12 Powder" \* 200 Billion CFU's** (brand - Seeking Health) This probiotic can be purchased from Seeking Health website or other websites also sell this. \* Mix some ¼ teaspoon of the powder into pets meals or some plain yogurt and offer or oral syringe.

\*\* It takes 2 months to fully restore gut flora \*\*

**Note:** whether your pet has an infection or not (parasite, bacteria or fungal) or past or present Antibiotic drugs use, your pet needs this foundation of healing in the gut

and

# ~Colostrum powder or chewable tablets (yummy)

Colostrumhealth 100% pure Colostrum Powder http://www.colostrum.gen.nz/

#### Order online.

Colostrum can repair <u>leaky gut syndrome</u> which can be the cause of constipation. \* if constipation is caused by a deficiency in minerals - potassium and magnesium.



# CANINE ADULT DIET Guide \*short quick guide

Incorporate less " COOKED OR RAW MEAT MEALS or \*If feeding COMMERCIAL foods" (better to omit ALL artificial food) and go to "real food" meals.

Two weeks of 50/50 Veg & Meat Ratio with fruit meals separate.

#### Incorporate mono meals.

Eventually leading to a 3 x week raw meat meals if you can. If not just alternate between fruit meals and veggie & meat meals. In general, we see dogs do better on 25-30% raw meat, 10% meaty raw bone (e.g lamb shank or chicken necks if a small dog), and the rest Veggies and Fruits.

e.g. SMOKED kangaroo, SMOKED lamb necks from fruit and veg markets if you can find someone that does this. Avoid commercially sold type of dried treats. Other: Organic Tumeric powder, Coconut oil, Dates, Raw Carrots, Dried Fruit (no sultanas or raisins) Nuts, Seeds, Squashes, Spinach, Kale, Lightly Steamed Broccoli etc. Pulverise all the veggies for optimal absorption.

**Fasting:** Fast your pet for 24 hours once a week. You can start with 12 hours and work your way up. \* can give liquids during this time.

Following the 24 hour day fast, do a fruit day. You may add almond milk, organic maple syrup, coconut shreds, raw nuts and seeds in moderation to any fruit smoothies or fruit bowls to get them to eat it.

#### PLEASE OMIT ALL

Acidic type vegetables (ie tomatoes, oranges etc)
No pork meat
wheat products or foods
Avoid ALL kibble foods

Dried pigs ears or any other toxic processed treats etc.

ALL NEW CUSTOMERS RECEIVE A DIET SUGGESTION BOOKLET WITH MORE DETAIL ABOUT DIET AND SUPPLEMENTS and a HOMEOPATHIC HEAVY METALS VACCINE DETOX PILLS.



# Vaccination Options - Titre Test

No pet should ever be re-vaccinated, not ever, without a "Titre test" to see if it's actually needed.

#### What to do?

You will need to phone around to vet clinics to ask if any a vet can order the VacciCheck in-house test kit. Which is basically only AVAILABLE by request through from a Vet Clinic. Give the contact details so they can proceed with purchasing the kit. Then once it is at the clinic they can let you know and make an appointment. Some vets are not willing to do Titers, so find a more caring Vet to help you make a safe decision on vaccinations.

#### Biogal brand - VacciCheck® Antibody Test Kit

USA - www.VacciCheck.com

Australia - Give these details to your Vet - Laboratories Diagnostics NSW.

Contact: Sonia Whittle Tel: 61-2-9668-0600 Website: www.labdiagnostics.com.au

OR Find a supplier in your Country http://www.biogal.co.il/find-distributors

If your pet has seroconverted, no matter what their age, they do not need another vaccination. Nor do they need a 'booster' at a year old, as they already have long-lasting immunity.

Dr. Edwards DVM has titre test puppies 2 weeks after initial vaccination if it's positive then they need no more vaccinations (So - any dog or cat, of any age, can be titre tested 2 weeks after any C3 vaccination. If your puppy is going to respond, it will have by then. He has titre tested several puppies after their first vaccination like this, and they've had strongly positive titre levels. Any further vaccinations for these puppies (or kittens) would have been completely unnecessary and may have caused the puppy in question harm.

The whole 'booster' thing is really a false way to describe it - nearly all the so-called 'booster' vaccinations administered to our pets are completely unnecessary. No difference what country you live in, either. There is no need for a so-called 'booster' vaccination a year after the puppy or your kitten's vaccinations.

NO PET SHOULD EVER BE RE - VACCINATED, NOT EVER, WITHOUT A TITRE TEST TO SEE IF IT'S ACTUALLY NEEDED.Dr. Ronald Schutz's research around the duration of immunity for C3 (Dog) and F3 (Cat) vaccinations has been around for ages - and is only a short google away from anyone.

THE WHOLE BODY needs detoxing, repairing and strengthening, in order for your pet to maintain a strong foundation of the defense system to maintain a natural immune system that can stand up against anything that comes there way.

Once or if we go back to putting more toxins in the body - chemicals drugs, sprays, washes, inhalant sprays in the household, insecticides wormers, flea, and heartworm drugs, this is when your pets life will start going downhill again. .. with chronic or acute illness, skin, joint, organ damages to autoimmune disorder disease. All these toxins in the commercial foods and drugs and products will shorten your pets life, no different to our own lives.



### Side Effects of the CHEMICAL DRUGS

The found of 'Homeopathic medicine" physician Samuel Hahnemann said to "remove the obstacles to the cure"

#### **HOMEOPATHIC DETOX**

#### HAMPL ChemPoision 110 30ml plus Activiated Charcoal powder (set of 2)

\* This one is good for not well since taking a drug, or insecticide products, now seizures, paralysed, High liver enzymes etc. Or we can provide the one drug antidote instead of the combination detox.

#### HAMPL Pet Cleanse 042 30ml plus Activiated Charcoal powder (set of 2)

- \* This one is good for pets that have not had alot of drugs or other dangerous chemicals but still need cleasning. High liver enzymes etc.
- \* means we have this in homeopathic form of the toxic drug is in the above formulas to aid a detox from this drug. Or on request we can provide just one drug detox specifically to the drug your pet has not been well since taking it or had acute side effects from within minutes or hours. If you dont see a drug that your pet is taking or did take, please send a tiny sample of it to our clinic and we will post out the detox homeopathic version to you.

SOME OF THE VETERINARY DRUGS GIVEN TO PETS



THE BELOW PRODUCTS, SPRAYS, DROPS, TABLETS, INJECTIONS FROM 1 TO 5 are <u>classified</u> as a toxic insecticide drug (poisons).

# **INSECTICIDE POISON - PRODUCTS**

\* RoundUp (glyphosate) "used as a week killer" (and human-pet killer) The carcongenic insecticide spray which is commonly used around parks, gardens at home worldwide. This highly toxic chemical is made by Monsanto. The Lawsuit filed against them for over 4,000 across USA diagnsoed with cancer.https://www.baumhedlundlaw.com/toxic-tort-law/monsanto-roundup-lawsuit/

#### **TEST kit**

### Australia or Canada testing yourself or pets

http://hrilabs.org/animalstudy/

**USA** <a href="https://shop.mercola.com/product/2057/1/glyphosate-environmental-exposure-test-kit">https://shop.mercola.com/product/2057/1/glyphosate-environmental-exposure-test-kit</a>

1. \* COMFORTIS Comfortis® (spinosad) flea treatment.

Spinosad is a member of the spinosyns

which is classed as a toxic insecticide poison drug.

Spinosad contains two major factors, spinosyn A and spinosyn D.

**FOR CATS:** The most common adverse reaction recorded in clinical trials was vomiting. Other adverse reactions were: lethargy, decreased appetite, weight loss, and diarrhea. Use with caution with concomitant extra-label use of ivermectin. The safe use of Comfortis in breeding, pregnant, or lactating cats has not been evaluated. Therefore avoid.

**FOR DOGS**: The most common adverse reaction reported is vomiting. Other adverse reactions reported in decreasing order of frequency are: depression/lethargy, decreased appetite, incoordination, diarrhea, itching, trembling, excessive salivation and seizures.

Serious adverse reactions have been reported following use of ivermectin with Comfortis. Post Approval Experience (June 2009): The following adverse reactions are based on post-approval adverse drug event reporting.

The adverse reactions are listed in decreasing order of frequency: **vomiting**, **lethargy**, **anorexia**, **ataxia**, **diarrhea**, **pruritus**, **trembling in pain and nerve damage**, **hypersalivation and seizures**, **(death within a day of use)**.

Use of ivermectin with Comfortis, some dogs have experienced the following clinical signs: *trembling / twitching, salivation / drooling, seizures, ataxia, mydriasis, blindness and disorientation,* incoordination, excessive dilation of pupils, blindness and disorientation.

CHEWABLE TABLETS had a much higher percentage of acute adverse side effects compared to Active Topical drug.



# 2. \* PANACUR liquid

### (worming drug) which is classed as a toxic insecticide posion drug.

Panacur Liquid for Cats and dog, horse and other animals. Fenbendazole, marketed under the brand name Panacur, is a common anthelmintic -- or dewormer -- for cats, dogs, horses and other animals. SIDE EFFECTS -The FREQUENT side effects that may occur include infrequent vomiting or diarrhea.

#### 3. \* MILBEMAX

### (wormer and heartworm drug) which is classed as a toxic insecticide posion drug.

Milbemat is a toxic insecticide drug used to control of gastrointestinal worms in pets including roundworm (*Toxocara cati, Toxascaris leonina*), hookworm (*Ancylostoma tubaeforme*) and tapeworm (*Diplylidium caninum, Taenia spp.*) and heartworm What are the active ingredients in Milbemax DRUG and how do they work? Milbemax contains praziquantel and milbemycin. Both ingredients act by <u>disrupting the nervous system</u> of the worms resulting in their paralysis and death.

SIDE EFFECTS - Systemic signs (such as lethargy), neurological signs (such as ataxia and muscle tremors) and / or gastrointestinal signs (such as emesisand diarrhoea) have been observed after administration of the veterinary medicinal product.

CUSTOMER COMMENT

Hi - I have a 14 year old cat that has had a very bad reaction to Milbemax wormer he had it adminstered on Tuesday morning and within hours developed neurological symptoms, falling over. he has gone deaf (he was not before) and will not drink or eat. Now its Friday and he still has no appetite and will not touch any food though he has been going in garden and may be drinking from bird bath. have been back to vet and she cant offer any suggestions, they have spoken to Novatis who say have not had any problems but reading this forum and other internet news these side effects are not uncommon. He is now steady on his feet but still will not eat, vet has checked throat and swallowing seems to be fine. I have managed to put a few flakes of white fish in his mouth and he will swallow. he seems interested in food but picks it up and drops it. (normally he would eat three sachets of food a day)

# 4. Trifexis<sup>®</sup> (spinosad + milbemycin oxime)

(worming, flea, heartworms) which is classed as a toxic insecticide drug.

It is the monthly, beef-flavored tablet that poisions your pets blood so it will also be that toxic in the system it will also kills fleas and prevents **heartworm** disease, and treats and adult hookworm\*, roundworm and whipworm infection.

#### WHAT ARE THE SIDE EFFECTS OF TRIFEXIS?

The most common adverse reactions based on post-approval adverse drug event reporting are .. vomiting, lethargy, itching, loss of appetite, diarrhea, trembling / shaking, incoordination, seizures, excessive salivation and redness of the skin.

#### For example let's look at "Heartworm" - A realistic Look

Heartworm is of concern to all dog owners. It is not, however, the unavoidable scourge that many vets and pharmaceutical companies would have us believe. Most people do not have any idea what is involved in a dog getting infected with the Heartworm larvae. This article should educate you on the life cycle of this larvae and ways to prevent any effects if your dog is infected.

If your dog is reasonably healthy, eating a proper real food diet, is over use "vaccine free", is not currently on any Heartworm "preventives" or monthly chemical flea and tick preventives, their immune system should be



strong enough to fight off a heartworm infestation with none of the larvae reaching adulthood. In the event that some of the larvae should manage to get past your dog's immune system defenses and survive to adulthood, it is still far from a death sentence.

They will be much weakened, and the truly healthy dog will make a less hospitable host. Even at the adult stage, a healthy dog, possibly with some veterinary assistance, should be able to fight them off with no lasting ill effects to their health.

Nor are the currently available medical cures (arsenic drugs) for Heartworm more dangerous than the monthly "preventives," as the veterinary and pharmaceutical industries would have us believe.

Mosquitoes are the only known carrier of the infective stage of the heartworm larvae. In some parts of the country mosquitoes are a problem all year round, while in others this problem is limited to the warm months. There are many sites out there on the internet with lots of "information" on heartworm. The problem with much of it is that it is either wrong or incomplete. Much of it is intended to mislead and frighten you. One site says simply that a mosquito bites an infected dog then bites another dog and infects it.

While that, to some extent, in the most general of ways, is true, it is very misleading because the entire process is much more complicated and time consuming than this "explanation " would lead one to believe. Another explanation says that the mosquito injects the heartworm into your dog. This is simply false.

Long or semi haired dogs, or dogs that sleep and live inside a house do not need to on heartworm drugs, but you can safely use alternatives if so desired.

# 5. \* Bravecto<sup>®</sup> chewable tablets

(given as a poison to kill fleas, poisons ticks, and mites - demodex, sarcoptes and ear mites)

which is classed as a toxic insecticide much more toxic that Comfortis and Capstar drugs

Bravecto is a relatively new product released in 2014 which is used for **fleas and ticks**. It contains the active ingredient fluralaner, a type of isoxazoline which maintains protection for 12 weeks, meaning you only have to use the medicine as a preventative roughly once every 3 months. Thats means it has a high amount of pesticide chemical toxins in it.

#### How does it compare to Comfortis and Capstar?

Comfortis and Capstar are two other popular flea treatments. Unlike Bravecto, Comfortis does not kill ticks and needs to be given every month, and Capstar is not effective as a flea preventative. Which medicine you choose will come down to personal preference and how well your dog tolerates each of them.

As you may have read online, many owners have seen Bravecto has severely harmed their dog. Research before giving your dogs or cats edible pesticides, once eaten there is NO ANTIDOTE.



The official numbers of the European Medicines Agency: Serious side effects and animals that died after administration of these tick and flea treatments :

\*Note: with "animal medication" only about 1 % of the side effects is actually reported to the authorities.

So the number of reported side effects you see below is probably just the tip of the iceberg...

as of May 04, 2018

BRAVECTO

**Bravecto:** 

7964 reported serious side effects

on the market since 2014

1782 deaths after administration



**Nexgard:** 

12.394 reported serious side effects

\* on the market since 2011 **745** deaths after administration

Simparica

Simparica: on the market since 2015 (simparica no update available yet)

834 reported serious side effects

118 deaths after administration

A pesticide does not belong in / on the body of our pets!

More info? Facebook: Does Bravecto Kill Dogs — Website: www.bravectotruefacts.com

These are the most recent numbers of reported side effects and deaths after treatment with the tick and flea treatments Bravecto, Nexgard and Simparica.

Please note that only about 1 % of the side effects is actually reported by veterinarians and dog / cat owners.

Most vets will simply deny that the symptoms are related to administration of these pesticides BROADCAST FROM THE NETHERLANDS: THE ANTI BRAVECTO PETITION WITH 40.000 SIGNERS HAS BEEN DELIVERED AT THE MINISTRY

- please don't forget to share, to spread the word.... The petition will continue, so you can still sign it via www.bravectopetition.org



# TICK COLLARS MANGE MITE TREATMENT

(demodex mites or skin demodex)

There are ALWAYS alternatives that are safe - TICK AND FLEA e.g https://keepthetailwagging.com/flea-and-tick-collars/

Amitraz Insecticides Poisoning in Dogs

#### **Brand Names and Other Names of Amitra**

Mitaban® (Upjohn), Preventic® (Allerderm/Virbac), Taktic® (Hoechst/Roussel) and various generic brands.

Uses of Amitraz for Dogs

- Amitraz is primarily used to treat the skin disease demodectic mange.
- It is also used for tick prevention.

#### Mechanism of action of Amitraz poison is:

The acaricidal and insecticidal activity effect on **octopamine receptors** of the nerve cells in the brain. Parasites become hyperexcited, paralyzed and eventually die. This too can cause central nervous system paralysis in your pet.

### Symptoms of Amitraz insecticide poisoning (Amitraz toxicosis) in dogs include:

- Low blood pressure.
- Low body temperature.
- Elevated levels of blood glucose.
- Ataxia.
- Decreased heart rate.
- Vomiting.
- Diarrhea.
- Seizures.

#### The most common symptoms seen in a dog include:

- Vomiting
- Diarrhea
- Lethargy
- Weakness
- StaggeringDisorientation
- Hypothermia
- Abdominal pain
- Light or severe depression

Reference: https://wagwalking.com/condition/amitraz-insecticides-poisoning



**Apoquel drug** - Used mainly for skin dermatitis issues for Canines. (atopic)

Pruritus contol (not healing) associated with allergic dermatitis of atopic dermatitis in at least 12 months of age. (too dangerous to give to younger pets or feline use)

#### **Adverse Reactions Studies**

In a masked field study to assess the effectiveness and safety of oclacitinib for the control of atopic dermatitis in dogs, 152 dogs treated with APOQUEL and 147 dogs treated with placebo (vehicle control) were evaluated for safety. The majority of dogs in the placebo group withdrew from the 112-day study by Day 16. Adverse reactions reported (and percent of dogs affected) during Days 0-16 included diarrhea (4.6% APOQUEL, 3.4% placebo), vomiting (3.9% APOQUEL, 4.1% placebo), anorexia (2.6% APOQUEL, 0% placebo), new cutaneous or subcutaneous lump (2.6% APOQUEL, 2.7% placebo), and lethargy (2.0% APOQUEL, 1.4% placebo). In most cases, diarrhea, vomiting, anorexia, and lethargy spontaneously resolved with continued dosing. Dogs on APOQUEL had decreased leukocytes (neutrophil, eosinophil, and monocyte counts) and serum globulin, and increased cholesterol and lipase compared to the placebo group but group means remained within the normal range. Mean lymphocyte counts were transiently increased at Day 14 in the APOQUEL group.

Dogs that withdrew from the masked field study could enter an unmasked study where all dogs received APOQUEL. Between the masked and unmasked study, 283 dogs received at least one dose of APOQUEL. Of these 283 dogs, two dogs were withdrawn from study due to suspected treatment-related adverse reactions: one dog that had an intense flare-up of dermatitis and severe secondary pyoderma after 19 days of APOQUEL administration, and one dog that developed generalized demodicosis after 28 days of APOQUEL administration. Two other dogs on APOQUEL were withdrawn from study due to suspected or confirmed malignant neoplasia and subsequently euthanized, including one dog that developed signs associated with a heart base mass after 21 days of APOQUEL administration, and one dog that developed a Grade III mast cell tumor after 60 days of APOQUEL administration.

One of the 147 dogs in the placebo group developed a Grade I mast cell tumor and was withdrawn from the masked study. Additional dogs receiving APOQUEL were hospitalized for diagnosis and treatment of pneumonia (one dog), transient bloody vomiting and stool (one dog), and cystitis with urolithiasis (one dog).

In the 283 dogs that received APOQUEL, the following additional clinical signs were reported after beginning APOQUEL: pyoderma (12.0%), non-specified dermal lumps (12.0%), otitis (9.9%), vomiting (9.2%), diarrhea (6.0%), histiocytoma (3.9%), cystitis (3.5%), anorexia (3.2%), lethargy (2.8%), yeast skin infections (2.5%), pododermatitis (2.5%), lipoma (2.1%), polydipsia (1.4%), lymphadenopathy (1.1%), nausea (1.1%), increased appetite (1.1%), aggression (1.1%), and weight loss (0.7).

# Side Effects of Apoquel for Dogs (as its not recommended for cats)

Just like any other drug, Apoquel comes with its own set of side effects and possible reactions. Because of its immunosuppressive properties, it can increase your dog's vulnerability to infection. This includes demodicosis infections and neoplastic health conditions.

#### Other potential side effects of a pet taking Apoquel include:

- Vomiting
- Diarrhea
- Appetite loss
- <u>Lethargy</u>



- New cutaneous or subcutaneous lumps
- Decreased leukocytes
- · Decreased globulins
- An increase of lipase and cholesterol

#### Also experienced:

- Bloodied diarrhea
- Pneumonia
- UTIs
- Skin and ear infections
- Histiocytomas
- An increase in appetite
- An increase in aggression
- And developed polydipsia

These side effects are similar to what a dog may experience when taking steroids. Note that some of the side effects from taking this drug may resolve on their own with continued use, as the body adjusts.

Oclacitinib (Apoquel™) used for:

- · Control of atopic dermatitis without cortisone side effects
- Animals at risk of <u>diabetes</u>, <u>cushings disease</u>, <u>obesity</u>, <u>pancreatitis</u>
- · Animals prone to infections
- Dogs only

#### Not good for:

- Treatment on a budget
- Cats (not suitable)

#### **CUSTOMER COMMENT**

#### Skye Leibee

My dog was on Apoquel for one week when he began having seizures. We stopped the Apoquel, treated the seizures with phenobarbital for 5 weeks, and he was seizure free. He began chewing himself again, the dr said it was safe to put him back on Apoquel, which my dog did another round of for a week. He had a seizure, out of the blue, in the early evening of February 18. He subsequently had 2 more, within 2 hours of the first one. The last seizure came at about 1:30 am, on the 19th, which lasted about half an hour by the time we got him to the emergency vet. I had to say goodbye to my spunky dog of 12 years, who suffered a horrible grand mal seizure. Now I know he was an "older" dog, but he was otherwise healthy, with no history of seizures.

Reference: https://dogshealthproblems.com/apoquel-side-effects-in-dogs/



# **ANTIBIOTICS**

#### 6. \* METRONIDAZOLE

#### (Antibiotic drug)

Metronidazole is a prescription antibiotic used in dogs and cats to treat various conditions such as inflammatory bowel disease, nonspecific diarrhea disorders, infections caused by Giardia, and periodontal disease. SIDE EFFECTS - The more common side effects that can occur include: headache, loss of appetite, nausea or vomiting. diarrhea. heartburn. cramping in your stomach area, constipation, metallic taste.

#### 7. \* DOXYCYCLINE

broad spectrum antibiotic (tablets or oral liquid )ALSO CALLED (Vibramycin®, Doxychel®, Doxy Caps®, Bio-tab®, Monodox®, Doryx®, Doxirobe®)

Doxycycline is a broad spectrum antibiotic that is a member of the tetracycline family. It is used to fight bacterial infections in dogs and cats. Doxycycline is used to treat many different bacterial infections including, leptospirosis, toxoplasmosis, mycoplasma, psittacosis, and tick borne diseases including Lyme disease, ehrlichiosis, and Rocky Mountain spotted fever. Treatments using Doxycycline are generally termed as a long term treatment. SIDE EFFECTS and Drug Reactions

Doxycycline may result in these side effects: Nausea, Vomiting, Swallowing difficulties If Doxycycline makes your pet nauseous, try administering it with food. If it makes your pet have breathing or swallowing problems, follow tablet with a small amount of water.

**Doxycycline may react with these drugs:** Antacids, Anticoagulants, Bactericidals, Barbiturates, Cathartics, Gastro-intestinal protectants, Aminophylline, Digoxin, Insulin, Iron dextran, Kaolin/Pectin, Methoxyflurane, Sodium bicarbonate, Theophylline.

USE CAUTION WHEN ADMINISTERING THIS DRUG TO YOUNG PETS, PREGNANT OR LACTATING PETS, OR PETS WITH **SEVERE LIVER DAMAGE.** (AS THIS DRUG DAMAGES THE LIVER EVEN MORE SO)

#### 8. \* CLAVAMOX

broad spectrum antibiotic for cats and dogs.

Clavamox is a mix of amoxicillin and clavulanic acid, two medications that work in combination to fight bacterial infections. Clavamox may be used to treat skin wounds and abscesses, urinary tract infections, tooth infections, pneumonia, injuries caused by bites and other infections. The medication cannot be used to treat infections caused by a virus or parasite.

### Side effects are commonly seen:

- Diarrhea or loose stool
- Allergic reaction (labored breathing, hives, etc)
- Loss of appetite.
- Vomiting/Nausea.



# 9. \* BAYTRIL (Enrofloxacin)

broad spectrum antibiotic

### Side effects are commonly seen:

- Loss of appetite.
- Vomiting.
- Diarrhea.
- Lethargy.
- Convulsions.
- · Seizures in pets with CNS disorders.
- Cataracts if given long-term.

We have safe natural medicines to treat infections. (HAMPL Infection Fighter 71 50ml)

**NOTE** Tautopathic Treatment for example, by taking Doxycycline 30C or 200C homeopathically, it will not only help detox from side effects caused by the drug, but also treat candida - yeast infections.



# FUNGAL TREATMENT DRUG

# 8. FLUCONAZOLE Drug

is an antifungal drug effective against fungus and yeast (e.g. Ringworm treatment)

It is typically used for skin infections, fungus in the nailbeds, and more intense fungal infections including Blastomycosis and Histoplasmosis. It is related to Ketoconazole, but is able to more effectively cross the brain-blood barrier, making it more successful in treating fungal infections of the central nervous system. Fluconazole is also USED against ringworm, but is typically reserved for more serious infections. .

# Side Effects and Drug Reactions

Fluconazole has been found to be very safe in pets, but not extensively researched. These side effects have been observed in humans on this drug:

- Allergic reaction (labored breathing, hives, etc)
- Vomiting
- Diarrhea
- Skin rash
- Loss of appetite
- Dark urine or jaundice due to liver failure
- Pale tongue, gums, and nose

#### Fluconazole will react with these drugs:

- Amphotericin B
- Cyclosporine
- Hydrochlorothisazide
- Rifampin

DO NOT GIVE FLUCONAZOLE TO PREGNANT PETS. USE CAUTION WHEN ADMINISTERING THIS DRUG TO DIABETIC PETS, LACTATING PETS, KIDNEY DISEASE, OR LIVER DISEASE. AS THIS DRUG AFFECTS LIVER AND KIDNEYS IN A NEGATIVE WAY, THUS FURTHER DAMAGES THE ORGANS.



# PAIN RELIEF using a anti inflammatory drug

# 9. \* METETACAM (Meloxicam)

Oral Suspension for dogs and cats - as a anti inflammatory

Each mL contains 0.5 mg meloxicam in a yellowish suspension with an odour of honey and 1.5 mg of sodium benzoate (Ph. Eur.) as the preservative. Is a nonsteroidal anti-inflammatory drug (NSAID). Meloxicam works by reducing hormones that cause inflammation and pain in the body. It is often given after surgery.

**Warnings**: Should not be administered if gastric or intestinal ulceration or bleeding is suspected; if there is evidence of cardiac, hepatic or renal disease; or if there is evidence of a haemorrhagic disorder or individual hypersensitivity to the product. Do not administer concurrently, other steroidal or nonsteroidal anti-inflammatory drugs (NSAIDs), aminoglycoside antibiotics or anticoagulant agents.

Pre-treatment with other steroidal or nonsteroidal anti-inflammatory drugs (NSAIDs) may result in additional or increased adverse reactions and accordingly a treatment-free period with such drugs should be observed for at least 24 hours before commencement of treatment depending on the pharmacokinetic properties of the products used previously. Repeat dosing in cats at doses higher than currently recommended has been associated with acute renal failure and death.

### SIDE EFFECTS

Typical adverse reactions of NSAIDs, such as loss of appetite, vomiting, diarrhea, apathy and renal failure have occasionally been reported. These adverse effects are in most cases transient and disappear following termination of the treatment but in very rare cases may be serious or fatal.

In some cases, death has been associated with some of these adverse reactions.

**Systemic**: anorexia, lethargy, death. **Gastrointestinal**: vomiting, diarrhea.

Renal and urinary: renal disorder, renal failure, urine abnormalities. Repeat dosing in cats at doses

higher than currently recommended has been associated with acute renal failure and death.

~ The most common adverse reactions reported involve the kidneys and the gastrointestinal tract.

Typical symptoms include loss of appetite, depression and vomiting. It is important in these situations to discontinue treatment immediately and contact your veterinarian. In most cases, the adverse reactions are transient and disappear after termination of treatment but in rare instances may be serious especially if treatment is not discontinued. Consult your veterinarian.



#### 10. \* TRAMADOL

#### pain relief drug

This medication is used to help relieve moderate to moderately severe pain. <u>Tramadol</u> is similar to <u>opioid</u> (narcotic) analgesics. It works in the brain to change how your body feels and responds to pain.

SIDE EFFECTS - Stop using tramadol and call your veterinarian at once if your pet has any of these serious side effects: seizure; a red, blistering, peeling skin rash; or shallow breathing.

Less serious side effects may include: drowsiness, and weakness; vomiting, constipation, loss of appetite; blurred vision; insomnia.

# PAIN AND INFLAMMATION DRUG

### 11. FIROXIB drug

#### (use for pain and inflammation associated with osteoarthritis)

Firocoxib (Previcox) is used in dogs for pain and inflammation associated with osteoarthritis. Firocoxib is also used to help control postoperative pain and inflammation associated with soft-tissue and orthopedic surgery.

Side Effects Firocoxib (Previcox) like other NSAIDs may cause some side effects. The most common side effects of Firocoxib involve digestive issues such as vomiting and decrease appetite. Other possible side effects of Firocoxib include:

- Change in bowl movements (diarrhea, or black, tarry or bloody stools)
- Change in behavior (decreased or increased activity level, incoordination, seizure, or aggression)
- <u>Jaundice</u> (yellowing of gums, skin or whites of eyes)
- Increase water consumption or urination changes (frequency, color, or smell)
- Skin irritation (redness, scabs, or scratching)
- Unexpected weight loss

It is important to stop the medication and immediately contact your veterinarian if you think your dog has any medical problems or side effects while taking Firocoxib (Previcox). Precautions - Firocoxib (Previcox) is for use in dogs only. Firocoxib should not be given to dogs that are hypersensitive to NSAIDs. Firocoxib should not be given with any other NSAIDs, including: Carprofen (Rimadyl), Meloxicam (Metacam), Etodolac (Etogesic), Deracoxib (Deramaxx), Aspirin. Dogs that are dehydrated, on concomitant diuretic therapy, or those with existing renal, cardiovascular, and/or hepatic dysfunction are at a greater risk for experiencing adverse events. Use of Firocoxib (Previcox) at doses above the recommended dose in puppies less than seven months of age has been associated with serious adverse reaction, including death. Drug Interactions Firocoxib (Previcox) should not be given with other NSAIDs or corticosteroids (e.g., prednisone, cortisone, dexamethasone or tramcinolone).



### Signs of Toxicity - Firocoxib (Previcox) can cause:

Loss of appetite, Vomiting, Diarrhea, Dark or tarry stool, Increased thirst, Increase urination, Pale gums, Jaundice, Lethargy, Fast or heavy breathing, Incoordination, Seizures, Behavior changes. If you suspect or know you dog has had an overdose, it could be fatal so please contact your veterinarian, an emergency vet clinic, or the Pet Poison Helpline at (855) 213-6680 immediately.

# SYNTHETIC STEROIDS

12. \* Prednisone (Prednisolone) Brand names: Prednisone Intensol and Rayos
Cortisone (hydrocortisone) Brand names: A-Hydrocort, Solu-CORTEE, and Cortef.

#### What this drug does

<u>Prednisone</u> and hydrocortisone are synthetic <u>corticosteroids</u> used for suppressing the immune system and inflammation. Both <u>drugs</u> are used to treat many conditions including <u>arthritis</u>, <u>colitis</u>, <u>asthma</u>, <u>bronchitis</u>, skin problems, and <u>allergies</u>.

- Brand names of prednisone include Prednisone Intensol and Rayos.
   Brand names of hydrocortisone include A-Hydrocort, Solu-CORTEE, and Cortef.
- Common side effects of both prednisone and hydrocortisone include retention of sodium (salt) and fluid, weight gain, high blood pressure (hypertension), potassium loss, headache, muscle weakness, nausea, vomiting, acne, thinning skin, restlessness, and psychiatric disturbances (depression, euphoria, insomnia, mood swings, personality changes, and psychotic behavior).
- **Prolonged use** of prednisone or hydrocortisone can depress the ability of body's adrenal glands to produce corticosteroids.
- Abruptly stopping prednisone or hydrocortisone can cause symptoms of <u>corticosteroid</u> insufficiency, such as <u>nausea</u>, <u>vomiting</u>, weakness, <u>fatigue</u>, <u>decreased appetite</u>, <u>weight loss</u>, <u>diarrhea</u>, <u>abdominal pain</u>, and <u>shock</u>.

Prednisone and prednisolone are medications acts on almost every part of the body and have a wide range of effects including reducing inflammation, suppressing the immune system, inhibiting healing, altering mood, stimulating appetite, increasing the secretion of gastric acid, weakening muscles, thinning the skin, and more. In your pet's liver, prednisone is converted to prednisolone.

Pets with liver problems are not able to make this conversion effectively, and many veterinarians believe that these pets should only be given prednisolone.

Cats also have a limited ability to convert prednisone into prednisolone, so prednisolone is the preferred medication in this species.



# Side effects - Prednisolone drug (cortisone)

Side Effects of Prednisone and Prednisolone. When pets have to be on prednisone or prednisolone at high doses and/or for long periods of time the risk of significant side effects like the following rises.

- An increased risk of infection
- · Altered behavior, including aggression
- Inhibited growth in young pets
- Development or worsening of diabetes
- Cushing's disease
- Vomiting
- Diarrhea
- · Ulceration of the digestive tract
- Lethargy
- · Delayed healing

A pet that has been on years of prednisolone will need to have his/her immune system treated and repaired from an initially low immune system (allergy reactions) and further damage from overuse of prednisolone drug injections.

# There are many many more drugs that are available.

If you dont see your drug (liquid, oral tablet, or injections, topical applications, skin treatments etc). If you need help, please save a tiny sample and post to clinic with name of drug and your full name and address so we can help -post back to a homeopathic drug detox drops from short term or long term side effects and organs damages. When we give this drug in homeopathic form in a 30Cor 200C (called tautopathic therapy SEE PAGES 23-26

#### **EXAMPLE:**

The homeopathic DOXYCYCLINE - broad spectrum antibiotic (Vibramycin®, Doxychel®, Doxy Caps®, Bio-tab®, Monodox®, Doryx®, Doxirobe®) in a 30C or 200C

Will clear side effects as well as in a this will clear symptoms of candida yeast infections or prevent it.



# Also important to use in conjunction with

### "Activated Charcoal" Health Benefits

Why "activated Charcoal powder"?

Health Benefits The American Journal of Gastroenterology published a study reporting a scientific claim that activated charcoal can reduce the build-up of **nitrogen waste** in the body; e.g. ..UREA levels (toxic waste) from kidneys. Research studies have also shown that **activated charcoal decreases the toxic load on the body by over 60% with daily use.** This is due to its ability to bind and grab up chlorine, pesticides and herbicides, bacteria, viruses, yeast, parasites, toxic chemicals and endotoxic metabolites.

Endotoxins These are substances that are produced within our body. This would include toxins produced by our gut bacteria, molecules formed by incomplete liver conjugation and <u>ammonia</u> which can arise due to inadequate kidney filtration.

Kidney Cleansing-The kidneys are critical to filtering and removing toxins through the urinary system. Additionally, it takes urea out of the body and when it is functioning poorly it can lead to elevated ammonia, which is extremely toxic to the brain.

Poorly functioning kidneys are also major contributors to blood pressure problems and cardiovascular disease. It also absorbs toxins in the kidneys and livers preventing a toxic overload, and at the same time allowing the vital organs (.. liver and kidneys) in the body to function normally. It also can help to eliminate the toxins in the body because it stimulates the bile flow

#### **SUGGESTION**

to assist your pet with – whether acute or chronic kidney weakness. Starting adding activated charcoal to your pets daily meals. e.g. get the "activated charcoal" in powder form.

**FELINE, TOY DOG to SMALL SPECIES:** Add 5 drops of the RenoAid 19-1 formula, and 1 teaspoon into a cup of filtered water in a jar with lid, shake before each use. Pour 1-2 tablespoon in meals or/and milk drink. (or build up to a tablespoon each meal or just add once a day to meals).

**SMALL to MED DOG:** Add 5 drops of the RenoAid 19-1 formula, and 2 teaspoons of AC to a cup of filtered water, stir well, then each day add one tablespoon in meals or milk drink. (or build up to a tablespoon each meal or just add once a day to meals).

**LARGE DOG:** Add 5 drops of the RenoAid 19-1 formula and double the amount of powder again into a cup of water.

Once pet has recovered, continuing using the Activiated Charcoal powder once a day for good health.

For humans a capsule in the evening before bed. Dr Robin Murphy recommends.

The found of Homeopathic medicine the physician Samuel Hahnemann said to "remove the obstacles to the cure"



**Dr Robin Murphy** .. says... Allopathic prescribed drugs kill outright 120,000 people per year: these are the reported cases only.

Indirectly they kill many more thousands of people. Modern medicine uses weird numerology to manipulate the statistics and they call it a medical science. The average American has a 9000% greater chance of dying from allopathic drugs than from a loaded gun in their own home.! A person has double the chance of dying from their doctor than from a car accident.

Classical Homeopathy is useless in cases of severe pathologies and patients on allopathic drugs. My practice consists of 60% to 70% of these types of patients. Most patients today are on 2-3 different drugs. Therefore we should be familiar with the top allopathic drugs being prescribed today. We need to know the top heart, blood pressure and diabetic remedies, etc. We need to know: drug side affects, drug overdoses, withdrawal symptoms, etc. We need to know them as potentized remedies also.

In homeopathic prescribing, I never take a patient off their drugs, therefore the allopaths don"t object. I always give low potencies of the indicated remedies along with herbal tinctures. Why? Because low potencies given daily are less prone to be antidoted by the drugs. I give higher potencies for drug overdoses, and acute withdrawal symptoms.

Once the patient is doing better, the drug doses are lowered and eventually stopped. We can get patients off their drugs by treating the underlying reason they are taking the drugs.

As the allopathic drugs are reduced, the cure rate is increased.

There are 1500 new drugs annually, these drugs are new to the human gene pool. Therefore any approved drug is still one foreign to human DNA for generations. The true toxics effects of any drug may not be known for at least 40 years or longer. Human are being chemically manipulated, poisoned and experimented on, for money and no other reason.

Chemotherapy is as toxic as getting cholera and but we have to pay the chemo poisoning and cholera is free! We are also being poisoned by Synthetic Chemicals called Xenobiotics that have been around since the early1900's. Since then there are thousands new xenobiotics every year also. Tap water has 56,000 xenobiotics, the US government tests for only 56 additions of these chemicals added to tap water.

Since 1900, 2 million pesticides, herbicides and chemicals are in the human blood stream. **Example**: In brain tumour cases, a chemical called Sodium Laurel Sulphate may be a cause of the cancer. This chemical is found in most shampoos to create foam, even the ones in the health stores. Sodium Laurel Sulphate is detected in the blood stream after one use only. We shampoo our babies hair with a cancer causing agent on a daily basis!



# The Nine Methods of Tautopathic Prescribing

A practitioner needs to be familiar with the Mims or BNF (British National Formulary) of allopathic drugs. It is important not to be predjudice about Allopathy, but to understand their drugs, as any morbific agent, such as a grief or trauma. In the treatment of patients who are on drugs, if you can use tautopathy first, you can demonstrate what homeopathy can do.

### 1. Tautopathic Prescribing for Drug Side Effects.

Drug side effects are a major obstacle to the cure. The biggest hazard lies in mixing up the case. We can get acute side effects of drugs and chronic symptoms of the patient confused. It's important to separate these two. We need to learn to distinguish between the side effects of the drugs and the symptoms of the patient when we analyze the case history. We need to learn the symptoms of allopathic drugs like we do with the homeopathic remedies.

Example: You might ask, "did you have these symptoms before or after a drug?" Then, depending on the answer, you might look at remedies like Oophorinum, Ovarian, Oestrogen etc for hormonal cases.

# 2. Tautopathic Prescribing for DRUG LAYERS

"Never been well since taking a drug or chemical poisonings" is a common phenomenon in modern practice. Drug poisonings can leave a drug layers. If a drug layer exists in the case then it must be removed first. Otherwise the patient's health will not improve. Mixing these drug layers with the general case is a big mistake. Also ignoring the top layer of the case can lead to a failure to cure or even palliate.

#### 3. Tautopathic Prescribing for DRUG WITHDRAWALS

The drug withdrawal symptoms are a whole new case. Treat all drug withdrawals as acutes, and use the tautopathic remedies in higher potencies. Flush the drugs out with water, by putting the tautopathic remedies in water.

Also use herbal tinctures to help with the detoxification process. Avena Sativa for <u>morphine</u> <u>or heroin withdrawal.</u> Chemotherapy and radiotherapy should be treated as acute poisonings. Go for the herbs that fit the organ and give the tautopathic remedies.



### 4. Tautopathic Prescribing for DRUG Addictions

Clinical homeopathy can assist greatly in the removal of cravings and addictions. Both psychological and physiological addictions can be treated effectively. Cravings for cigarettes, alcohol, coffee, marijuana, or allopathic drugs are all treatable with tautopathics. Is this being taught in the homeopathic colleges?

**Example:** for alcohol craving: Lachesis, Quercus rubens, Sulphuric acid or Amethyst, (known as the sobriety stone). However, Alcohol in potency is a great remedy. Alcohol dehydrates the brain and can lead to many chronic disorders, including epilepsy.

#### 5. Tautopathic Prescribing for DRUG and Chemical OVERDOSES

Acute drug overdoses are a common cause of death in our society. Knowing the correct tautopathic or homeopathic remedies can save someone's life.

**Example:** Barbiturate overdose symptoms are slurred speech, drowsiness, murmuring, and heavy eyes, followed by coma and death. Phenobarb. in potency is the antidote for phenobarb. The clinical picture of Gelsemium also fits.

# 6. Tautopathic Prescribing of ALLOPATHIC Drugs based on their ALLOPATHIC indictions, AS IN THE BNF / MIMS.

In spite of all the new provings going on these days, I believe learning allopathic drugs is more important. We instantly have 1,000 new chemical remedies. This involves prescribing a remedy based on its allopathic indications, because it is already a well-proven remedy.

**Example:** Ritalin is used in hyperactivity therefore it has that similar affinity, for that disease. It is a stimulant that treats over stimulated children. Therefore Ritalin in potency is a great remedy for treating hyperactivity.

**Example:** Dopamine is the drug for Parkinson's disease. Therefore Dopamine is a remedy for Parkinson's disease. These patients also have a dopamine deficiency.

**Example:** a child is a Silica constitution at birth, but after abuse goes into a Staphisagria layer, then, after difficult teenage years he gets given Prozac. These remedies have an affinity for each other, the fact that prozac treats a Staphisagria like condition means that the patient is sensitive to prozac. Drugs like prozac are new remedies: they stand with staphisagria, Arsenicum & Carcinosin.

We must be unprejudiced and treat them like another remedy.



# 7. Tautopathic Prescribing of ALLOPATHIC Drugs and Chemicals based on their HOMEOPATHIC indications.

Remedies like Lithium, Alcohol, Mercury, Lead and Adrenalin have been in the repertory for over a hundred years; via provings and case histories we have the indications for prescribing them as homeopathic remedies. That means any drug or chemical can be used as a remedy to treat things beyond it's normal allopathic indications.

#### 8. Tautopathic Prescribing of PREVENTATIVE REMEDIES

Also known as prophylactic tautopathy: If one knows a person is about to be exposed to a toxic substance then one can give a tautopathic remedy to help them through the toxic exposure.

**Example**: Before a visit to the dentist give the anaesthetic in potency, especially if the patient has a history of poor recovery from that drug. Likewise for chemotherapy you can give Cadmium sulph., on the day of the chemo treatment to lessen the side effects of the treatment.

Please let us know if you have any more questions as we would love to help.

Best Regards,

Admin Staff

HAMPL Clinic | 60 Angove Street, North Perth. 6006. WA. Australia Phone (+61)1300 132 966 | Fax (+61)1300 855 183 Visit On-Line Store | www.HolisticAnimalRemedies.com