

PRODUCT CATALOGUE



WELCOME TO THE BREMER PRODUCT CATALOGUE





Based in Warburg, Germany, **BREMER PHARMA GMBH** produces high-quality veterinary medicines in Germany since 1982 and exports to more than 50 countries worldwide. The company holds numerous product approvals in Germany and abroad.

The name BREMER PHARMA GMBH stands for highest quality and constant improvement of the products. Suppliers, raw materials and finished products are therefore subject to constant monitoring. In order to maintain this standard, BREMER PHARMA GMBH works exclusively with well-trained personnel. BREMER PHARMA GMBH is subject to the regular control of national and international authorities. Moreover, a comprehensive quality assurance system and its regular monitoring ensure the high standard of our products and services.

The Veterinary Group LLC (TVG) was born out of a desire to make a difference to the Animal Health landscape by an extremely experienced successful management team with over 60 years combined experience in the region across multiple species.

Through innovation, education, trust and perseverance TVG is passionate about driving change and delivering excellence in the Animal Health industry.



www.thevet.group



Antibiotics Hefrotrim Injection 240mg/ml 100 ml 4 Speciality Metaphosol 200mg/ml 100 ml 5 Hormones Synchromate 0.250mg/ml 20 ml 6







HEFROTRIM INJECTION 240MG/ML 100 ML INJECTION

For Cattle, Horses, Sheep, Goats, Dogs and Cats

Antibiotic against Gram-positive and Gram-negative bacteria and coccidiosis



Active ingredient: Sulfadimidine 200 mg/ml, Trimethoprim 40 mg/ml

Pack size: 100 ml

Indications for use: Therapy of diseases caused by bacteria sensitive to sulfadimidine and

trimethoprim: Infections of the respiratory tract, lungs, skin, gastrointestinal and urogenital tracts. Secondary bacterial infections during and after viral diseases, especially of the respiratory and gastrointestinal tract, wound and navel infections, infections of the

claws (e.g. paronychia).

Dosage: Horses: 1 ml per 10-15 kg b.w. (sc, im)

Cattle: 1 ml per 10-15 kg b.w. (sc, im, iv)

Sheep and Goats: 1 ml per 10-15 kg b.w. (im, iv) Dogs and Cats: 0.1 - 0.2 ml per kg b.w. (sc, im, iv)

The duration of the treatment is once-a-day between 3 to 5 days, depending on the severity of the disease. Volume at the same site

should not exceed 10 ml.

If after three days of treatment no signs of improvement are seen, the

diagnosis and the therapy should be reconsidered.

Storage: Store below 30°C. Protect from light.



METAPHOSOL 200MG/ML 100 ML SOLUTION FOR INJECTION

For Cattle, Horses, Sheep, Goats, **Dogs and Cats**

Treatment and prophylaxis of metabolic disorders.



Active ingredient: Toldimfos sodium trihydrate 200 mg/ml

Pack size: 100 ml

- Indications for use: To treat post parturient paresis, post parturient hemoglobinuria, downer cow syndrome.
 - To treat skeletal deformities such as rickets, stunted growth in young animals, osteomalacia in adult animals and fracture in conjunction with Calcium and Vit. D3.
 - As a supportive treatment in weakness and prostration after disease, dystochia, surgery.
 - Treatment and prophylaxis of tetanus and paresis due to disturbances of Calcium, Magnesium metabolism in conjunction with Calcium, magnesium infusions.
 - Treatment of infertility and improvement of conception rate.

Dosage:

For treatment of conditions resulting in acute hypophosphatemia:

Cattle and Horses: 10 to 30 ml per animal IV or IM. Repeat till the clinical effect up to 10 injections.

Sheep and Goats: 5 to 15 ml per animal IV or IM. Repeat till the clinical effect upto 10 injections.

Dogs and Cats: 0.25 to 5 ml per animal as slow IV injection over 6 hours, stop when serum phosphorus level reaches 2 mg/dl.

For treatment of conditions associated with chronic hypophosphatemia: Cattle and Horses: 10 to 25 ml per animal IM or SC repeat till the clinical effect up to 10 injections.

Sheep, Goat and Calf: 2 to 5 ml per animal IM or SC, till

the clinical effect up to 10 injections.

Dogs and Cats: 1 to 3 ml IM or SC every 2 days, repeat till clinical effect up

to 10 injections.

Storage: Store below 30°C. Protect from light.



SYNCHROMATE 0.250MG/ML 20 ML

For Cattle, Horses, Sheep and Goats

Cyclus synchronication, induction of abortion and labour.



Active ingredient: (±)Cloprostenol 0.250 mg/g (as sodium 0.263 mg/g)

Pack size: 20 ml

Indications for use: Cattle Suboestrus or non-detected (cows): For

> synchronisation of oestrus, induction of parturition, termination of normal pregnancy, termination of abnormal pregnancy, mummified foetus, hydrops of the foetal membranes, chronic endometritis

(pyometra), ovarian luteal cysts, controlled breeding.

Horses (mares): For induction of luteolysis following early foetal death and resorption, termination of persistent dioestrus, termination of pseudopregnancy, causing abortion prior to the twelfth day of gestation, treatment of lactation anoestrus, establishing oestrous cycles in barren/maiden mares, as an aid to stud management.

Sheep and Goats: For induction of parturition in the final stages of pregnancy, oestrus induction or synchronisation during the breeding season.

Dosage:

For IM injection. For deep IM injection in pigs, cattle, sheep, goats are also injectable by SC route

Cattle (cows): 2.0 ml Synchromate (equivalent to 500 µg of (±)cloprostenol) / animal and treatment.

- Suboestrus or non-detected oestrus / synchronisation of oestrus / ovarian luteal cysts / controlled breeding: After the first injection, inseminate any cows showing signs of heat. Animals that do not show signs of heat should be injected 11 days after the first injection and then inseminated 72-96 hours later.
- Chronic endometritis (Pyometra): One dose, if necessary, repeat treatment after 14 days.
- Termination of normal / abnormal pregnancy: To induce parturition use single dose.

Horses (mares): 0.5-1.0 ml Synchromate per pony, 1-2 ml Synchromate /

Sheep and Goats: 0.5 ml Synchromate per animal

- Initiation of birth: one injection
- Oestrus induction or synchronisation during the breeding season: one injection, for oestrus synchronisation animals may be treated twice 11 days apart.

Storage:

Store below 30°C. Protect from light.

Veterinary medicines for livestock





the veterinary group







THE VETERINARY GROUP LLC

THE VETERINARY GROUP LLC



The Veterinary Group LLC (TVG) Management office Detroit House, Office 410, Motor City, Dubai United Arab Emirates Phone: +971 6 539 5443 WhatsApp: +971 6 539 5443 Email: sales@thevet.group

www.thevet.group







