### What is BetaTan injection:

BetaTan is Melanotan II (also known as Bremelanotide). Melanotan II is a synthetic analogue of the peptide hormone  $\alpha$ -melanocyte-stimulating hormone ( $\alpha$ -MSH) that stimulates melanogenesis.

Melanotan II is chemically identical to a substance in our bodies, called "melanocyte-stimulating hormone."

### How does it BetaTan work?

This hormone increases the production of skindarkening pigments. Melanotan II.

### Possible side-effects:

BetaTan may cause nausea, stomach cramps, decreased appetite, flushing, tiredness, yawning, darkened skin, and other side effects. In some people, especially those with light skin, a change in the shape of moles, new moles, have all occurred in people using BetaTan.

### Pregnancy and breast-feeding:

There isn't enough clinical information available to know if BetaTan is safe to use when pregnant or breast-feeding. Consult your medical practitioner prior to use.


# How to inject BetaTan:

Administering BetaTan is a very easy and painless operation and is best performed just before you go to bed to minimize any side effects that may occur (tiredness).







Then flick to remove any air that me be in the syringe.



Now find an area of your stomach to the left or right of your bellybutton and then clean with an alcohol wipe.



Now place the index finger and thumb about 4 centimeters apart on the cleaned area of skin and simply pinch. Take the syringe in your other hand, holding it like a pen 90 degrees to your skin, needle side down. Now insert the needle with a swift motion until it is fully under the skin. Gently depress the plunger until all the solution has gone from the syringe and carefully remove it from your skin.



Next gently hold the alcohol wipe on the injected area for 1 minute.

You have now administered your first dose of BetaTan. One 3ml cartridge of BetaTan will give you 20 injections.

### Dosage:

Whilst you are going through the 'loading phase' we recommend you inject 0.15ml (15 units on an insulin syringe) of BetaTan every day until the vial is empty. During this process you will also need to have at least one sunbed session per week. Although this is not vital, it brings your tan out quicker if you do. Under normal circumstances one vial of BetaTan will get you your perfect tan. Some people, however, do take longer so you may need to use another vial during this process.

Once you have reached the desired tan you are happy with, you only need to inject 0.15ml (15 units on an insulin syringe) of BetaTan twice a week.

## How it is supplied

BetaTan is supplied in a sterile, liquid 3 mL cartridge. Each BetaTan is filled with 3 mL containing 20mg of Melanotan II, Mannitol, 85% O-phosphoric acid, L-Histidine, Poloxamer 188, Phenol, Sodium Hydroxide (for pH adjustment), and Water for Injection to deliver 20mg of Melanotan II in 3 mL.

## Storage:

BetaTan must be stored refrigerated between 2-8°C (36-46°F) before being dispensed to the user. Users should keep the cartridge refrigerated to allow the product to be used until the expiry date shown on the cartridge. BetaTan may be stored by the user for no more than 30 days at room temperature (up to 25°C (77°F) but must be used within those 30 days.