

**COPPER SOAP:** Reg No. L7064 Act 36 of 1947 **Natural organic Fungicide contains ammoniated soap of fatty acid**

**DIRECTIONS FOR USE: (use only as directed)**

Shake well before use.

This product may be used both indoors and outdoors.

Do not apply this product through any type of irrigation system.

Read and follow all applicable directions and precautions on the label of other products, before mixing them with COPPER SOAP.

Pour COPPER SOAP into spray tank at least half filled with water using adequate agitation.

COPPER SOAP is compatible with most commonly used insecticides and fungicides, however physical compatibility should first be checked by using the current proportion of products in a small jar test.

Do not mix COPPER SOAP with oil when applied to citrus.

Do not mix COPPER SOAP with chelated or liquid fertilizers.

Do not apply in temperatures above 30°C.

A spreader may be used to improve spreading on hard-to-wet plants.

COPPER SOAP can be applied up to day of harvest.

CROP	DISEASE	DOSAGE / 100ℓ water	DIRECTIONS FOR APPLICATION
Flowers	Downy Mildew Powdery Mildew	500 mL – 1ℓ per 100 ℓ water	Ensure thorough wetting of plant. Use higher rate under high infections. Apply every 7 – 10 days, depending on weather conditions favourable for the disease. Re-apply after rainfall. Established infestations will require more than 1 application.
Tomatoes	Early Blight	1ℓ - 2ℓ per 100 ℓ water	

**COPPER SOAP:** Reg No. L7064 Act 36 of 1947 **Natural organic Fungicide contains ammoniated soap of fatty acid**

**GEBRUIKSAANWYSINGS:** (gebruik slegs soos voorgeskryf)

Skud goed voor gebruik.

Moenie hierdie produk deur enige soort besproeiingssisteem toedien nie.

Lees en volg alle aanwysings en waarskuwings wat van toepassing is op ander produkte se etikette, voordat hulle met COPPER SOAP gemeng word.

Gooi COPPER SOAP in 'n spuittenk wat half vol water is en meng genoegsaam.

COPPER SOAP kan met meeste algemeen gebruikte insekdoders en swamdoders gemeng word, maar fisiese mengbaarheid behoort eers getoets te word deur die aanbevole dosisse van die produkte in 'n klein houer te meng.

Moenie COPPER SOAP met olie meng wanneer dit op sitrus toegedien word nie.

Moenie COPPER SOAP met chelaat- of vloeibare misstof gebruik nie.

Moenie in temperature bo 30° C toedien nie.

'n Benatter kan gebruik word op mocilik-benatbare plante.

COPPER SOAP kan tot op een dag voor geoes word toegedien word.

GEWAS	SIEKTE	DOSIS / 100ℓ water	AANWYSINGS VIR TOEDIENING
Blomme	Poeieragtige skimmel Donsige skimmel	500 mL – 1ℓ	Bespuit plante deeglik. Gebruik hoër dosis tydens hoog infeksie. Dien elke 7 – 10 dae toe, afhangend van weersomstandighede wat gunstig is vir die siekte. Dien na reën weer toe. Gevestigde infestasies sal meer as een aanwending nodig hé.
Tamaties	Vroeë roes	1ℓ – 2ℓ	