

**PHYTA:** Reg No. K6152 Act 36 of 1947 **Foliar feed containing Potassium Phosphite**

**DIRECTIONS FOR USE:** (Use only as directed)

**COMPATIBILITY:**

Physical compatibility should first be checked by using the current proportions of products in a small jar test. Users should then test tank-mix combinations for phytotoxicity on a sample of crops prior to use.

Read other labels carefully.

Always add Phyta to spray tank first after initial pre-mixing with water.

Always shake container before opening.

Do not spray in temperatures above 30°C.

**PHYTA CROP RECOMMENDATION:**

<b>CROP</b>	<b>TIMING</b>	<b>RATE</b>	<b>COMMENTS</b>
Onions, leeks	3 - 4 leaf	2 - 4 ℓ / Ha 500 ml / 100ℓ	At 10 - 14 day intervals
Carrots, parsnips	2 - 3 weeks after emergence	2 - 4 ℓ / Ha 500 ml / 100ℓ	At 14 day intervals
Beans, peas	2 - 3 weeks after emergence	2 - 4 ℓ / Ha 500 ml / 100ℓ	At 14 day intervals
Lettuce	Start 7 - 10 days after planting	4 ℓ / Ha 500ml / 100 ℓ	Repeat twice
Brassicas	2 - 4 leaf stage	2 - 4 ℓ / Ha 500 ml / 100ℓ	At 10 - 14 day intervals
Strawberries	Green bud	4 ℓ / Ha 500ml / 100 ℓ	At 14 day intervals
Citrus		4 ℓ / Ha 500ml / 100 ℓ	Multiple applications as required
Grapes		4 ℓ / Ha 500ml / 100 ℓ	Multiple applications as required
Potatoes	From flowering through to tuberisation	4 ℓ / Ha 500ml / 100 ℓ	Three applications
Ornamentals and turf	From spring	500ml/100ℓ	Multiple applications as required