

**Juliana Houghton**<sup>1</sup>, School of Aquatic & Fishery Sciences, University of Washington, Box 355020, Seattle, Washington 98195

**Robin W. Baird**, Cascadia Research Collective, 218½ West 4th Avenue, Olympia, Washington 98501

**Candice K. Emmons**, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Northwest Fisheries Science Center, 2725 Montlake Boulevard East, Seattle, Washington 98112

and

**M. Bradley Hanson**, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Northwest Fisheries Science Center, 2725 Montlake Boulevard East, Seattle, Washington 98112

## Changes in the Occurrence and Behavior of Mammal-eating Killer Whales in Southern British Columbia and Washington State, 1987–2010

### Abstract

The primary prey species of mammal-eating killer whales in the Salish Sea, the inland waters of southern British Columbia and Washington state, have experienced dramatic increases in population abundances in the last 25 years. It is possible that changes in prey abundance over time have resulted in changes in predator spatial use, occurrence and group size. Focused studies of mammal-eating killer whale behavior in the area were undertaken from 1987–1993, and an extensive record of sightings with confirmed identifications was available from 2004–2010. Changes in occurrence across years, months, and subareas of the Salish Sea were examined as well as changes in group size and in the identity of specific matriline using the area. Occurrence of mammal-eating whales increased significantly from 2004–2010 with different seasonal peaks compared to 1987–1993. Different matriline occurred in different seasons, time periods, and subareas. Group size was larger in 2004–2010 than in 1987–1993. The whales may be increasing use of the area due to increasing prey abundance or an overall increase in the whale population size. Changes in seasonal patterns of occurrence and the increase in group size between the two periods could be due to increased prey diversity.

**Key words:** predator-prey interactions, spatial use, occurrence, group size, *Orcinus orca*

### Introduction

Top predators are key components of ecosystems around the globe. Their removal, and the consequent loss of ecological interactions they facilitate, may be detrimental to ecosystems (Janzen 1974; reviewed in Estes et al. 2011). The marine environment is particularly vulnerable to the loss of top predators due to human interactions such as fishing, whaling, bycatch, boat collisions, pollution, and destruction of habitat. Previously, research focus has been placed on the consumptive effects of predation on prey populations (Estes and Palmisano 1974, Lima 1998, Ainley et al. 2006, London et al. 2012), although increasing emphasis in examining non-consumptive effects

(e.g., changes in prey behavior due to perceived risk of predation) across multiple systems and taxa has recently developed in ecological studies, with an indication that these effects may be universal throughout ecosystems (Wirsing et al. 2008, Sheriff et al. 2009, Wirsing and Ripple 2010). For example, temporal variation in occurrence of sharks in Shark Bay, Australia, significantly alters the behavior of their prey (Heithaus et al. 2007, Wirsing et al. 2007), likely impacting the seagrass ecosystem as a whole. In Bermuda, where numbers of predatory sharks have decreased, an increase in herbivorous sea turtle populations have coincided with declines in seagrasses vital to the ecosystems of this area (Baum et al. 2003, Murdoch et al. 2007, Heithaus et al. 2007). Losing top predators can lead to ecosystem changes stemming from both consumptive and non-consumptive effects on

<sup>1</sup> Author to whom correspondence should be addressed.  
Email: stephj5@uw.edu

prey populations. An examination of occurrence patterns can provide insights into the potential effects of a loss, or of an increase, of predators.

Predator and prey populations can be tightly linked in marine ecosystems (May et al. 1979, McLaren and Peterson 1994, Whipple et al. 2000). As more prey become available through increased population abundance or increased occurrence in a given area (e.g., changes in distribution), predator abundance and occurrence may also increase according to numerical response theory (Soloman 1949).

Mammal-eating killer whales (*Orcinus orca*) in the Gulf of Alaska have been suggested to influence a variety of prey populations (Estes et al. 1998, Springer et al. 2003), although there is considerable controversy about their precise role in the Gulf of Alaska ecosystem (Demaster et al. 2006, Wade et al. 2007). Mammal-eating killer whales are frequently found in the Salish Sea, the inland waters of Washington state and British Columbia, and are classified as part of the west coast stock of 'transient' killer whales (Bigg et al. 1990, Ford et al. 1998, Allen and Angliss 2011). These killer whales are known to prey upon seals, sea lions, porpoises and occasionally other cetaceans throughout their range (Baird et al. 1992, Baird and Dill 1995, Ford et al. 2006, Dahlheim and White 2010). Several of the most common prey species consumed by mammal-eating killer whales in the Salish Sea have undergone dramatic changes in the last 30 years. Harbor seals (*Phoca vitulina*), resident to the area year-round, tripled in abundance between 1978 and 1999, reaching carrying capacity (Jeffries et al. 2003, Fisheries and Oceans Canada 2010, Carretta et al. 2011). California sea lion (*Zalophus californianus*) and Steller sea lion (*Eumetopias jubatus*) numbers in the area vary seasonally, with peak occurrence in April and May (Pat Gearin, NOAA National Marine Mammal Laboratory, personal communication), but both species have increased in abundance within the Salish Sea. The U.S. stock of California sea lions in the North Pacific increased in abundance at an annual rate of 5–6% between 1975 and 2005, suggesting that the population more than tripled throughout the 1987–2010 study period (Lowry

and Maravilla-Chavez 2005, Puget Sound Action Team 2007, Carretta et al. 2011). The eastern North Pacific Steller sea lion population (ranging from California to southeast Alaska) has also increased in abundance by an average of about 9% per year between 1987 and 2010, reaching the highest breeding population level in recorded history (Pitcher et al. 2007, Steven Jeffries, Washington Department of Fish and Wildlife, personal communication). This population is therefore four to five times larger in 2010 than it was in 1987. The California breeding stock of northern elephant seals (*Mirounga angustirostris*) has doubled in abundance from 1987 to 2006 and individuals are frequently observed in the Salish Sea year-round (Carretta et al. 2011). The population of harbor porpoise (*Phocoena phocoena*) in the Strait of Juan de Fuca and San Juan Islands has tripled since the early 1990s (M.B. Hanson, unpublished data), with a recolonization of this species to Puget Sound (Carretta et al. 2011). There is no trend information available on Dall's porpoise (*Phocoenoides dalli*) abundance, which also are found year-round in the Salish Sea. Although most of these prey species of mammal-eating killer whales have increased in abundance and occurrence, the potential effects on their predator are undocumented.

Current ecosystem-based models in the Salish Sea attempt to incorporate marine mammals as predators, but do not include mammal-eating killer whales due to a lack of current information on spatial use and occurrence (Harvey et al. 2010). The purpose of this study was to determine if there have been changes in mammal-eating killer whale occurrence, spatial use, and/or group size over the period 1987–2010. We hypothesized that there is a change in mammal-eating killer whale occurrence and group size over the 1987–2010 time period. The potential change could be due to an increase in prey abundance and occurrence in the Salish Sea from 1987–2010, or an overall increase in the size of the mammal-eating killer whale population (Ford et al. 2007, Towers et al. 2012). However, a causal relationship between increased prey and changes in killer whale occurrence and behavior cannot be identified from the sighting data. Our findings will inform ecosystem models and ecological studies of predator-prey

dynamics in the Salish Sea. This study will also provide insights for other ecosystems where large-bodied predators have the capacity to alter their movement and behavior on a large scale to take advantage of seasonally or locally abundant prey.

## Methods

Using existing long-term catalogs, individual killer whales were identified by unique markings (Bigg et al. 1987, Ford and Ellis 1999, Towers et al. 2012). Mammal-eating killer whales travel in matrilineal groups with dispersal occurring for some adult males and for females that have offspring of their own (Bigg et al. 1987, Ford and Ellis 1999, Baird and Whitehead 2000). For this study, a matriline was defined as a female killer whale and any offspring not known to have dispersed from the maternal group. Between 1987 and 1993, focused studies of mammal-eating killer whale behavior were undertaken in the Salish Sea from a small research vessel (Baird and Dill 1995, 1996). The objectives of the previous study were to document mammal-eating killer whale occurrence, seasonality, group size, and foraging behavior, and the identity of all whales present in each encounter was documented (Baird and Dill 1995, 1996). Documenting whale presence in that effort involved extensive small-boat surveys as well as responding to sightings reported by whale-watch operators, the general public, and other researchers working in the area. Effort from 1987–1993 was highest in the months of May to September (Baird and Dill 1995). From 2004–2010, killer whale sighting networks (i.e. organizations that collect and record sightings of killer whales from multiple sources, including whale-watching boats) created an extensive sighting record of identifications in the area, with attempts to document all occurrences in the area. Orca Network was the largest resource of identification data in the 2004–2010 dataset (Orca Network 2011). Within the 2004–2010 period, the number of commercial whale-watching boats was relatively constant (Giles and Koski 2012). With commercial whale-watching trends as a proxy for sighting effort of whales in the Salish Sea, effort is estimated to have been consistent from 2004–2010.

Prior to 2004, whale-watching effort and sighting network activity were inconsistent. Therefore the years 2004–2010 were chosen as a comparable number of years to the 1987–1993 dataset (Baird and Dill 1995, 1996). The spatial distribution of effort between the two time periods likely varied substantially, but this potential limitation was addressed in the data analysis.

The 2004–2010 sightings were collected mostly through citizen science and therefore reliability of the data may be variable. As a caveat of citizen science, it is possible that misidentifications may have occurred due to similar markings on different individuals. It is also possible that there was a failure to identify individuals or matriline actually present within aggregations or that group composition may have been based on one or two well-marked individuals. However, only confirmed identifications of mammal-eating killer whales were used in analyses. Identifications were confirmed if either a known, reputable source recorded the sighting (i.e., a researcher or whale-watch operator with several years of experience) or if multiple sources independently recorded the sighting. There are occasionally sightings reported to the sighting network with questionable identifications (e.g., “maybe T101”). Any such identification was not included in this study. Despite the limitations of citizen science, a previous study evaluated the reliability of sightings reported largely by commercial whale-watch operators in the same region (Hauser et al. 2006). It was determined that sighting data can be effective in determining the location and identification of killer whales in the Salish Sea in order to evaluate patterns of spatial use (Hauser et al. 2006, Hauser et al. 2007). The findings from this very similar dataset have validated the use of sighting data for our analyses, as long as the limitations of the method are acknowledged (e.g. only including identifications from experienced observers).

With increased effort from 2004–2010, there were often multiple sightings of the same group when it was encountered in the Salish Sea. To minimize repeated detections of the same group of killer whales when they were in the area, and reduce the bias associated with increased effort

TABLE 1. The number of concurrent days any matriline was encountered in the Salish Sea, restricted to matrilines seen on more than 20 occasions. There were no cases in which one of these matrilines was encountered more than 6 days concurrently.

Number of concurrent days encountered	Frequency
1 <sup>a</sup>	280
2	40
3	15
4	4
5	0
6	1

<sup>a</sup> No concurrent days after the first day of encounter

in the 2004–2010 period, encounters from both periods were reduced to a measure of occurrences. We accomplished this by examining the number of concurrent days that groups of whales were encountered in the Salish Sea (Table 1). From this analysis, we were able to determine that if a group was not encountered again within six days, it had likely left the Salish Sea and any encounters greater than six days before or after the original encounter should be treated as a separate occurrence of that group. Therefore, an occurrence was defined as an encounter of a unique group of mammal-eating killer whales when there was not an encounter of that group in the Salish Sea within six days before or after the encounter.

For geographic analyses the study area was split into five subareas: 1) central Strait of Juan de Fuca; 2) Gulf Islands and southern Strait of Georgia; 3) San Juan Islands (including eastern Strait of Juan de Fuca); 4) Puget Sound; and 5) Hood Canal (Figure 1). Hood Canal was separated from Puget Sound analyses due to an anomalous event of an extended stay by mammal-eating killer whales in 2005 (London et al. 2012). Research effort from 1987–1993 was concentrated in the central Strait of Juan de Fuca, San Juan Islands, and Gulf Islands and Strait of Georgia, with no effort in Puget Sound or Hood Canal. Any comparisons of area use and occurrence between the 1987–1993 and 2004–2010 time periods therefore did not include any occurrences in Puget Sound and Hood Canal.

Cumulative occurrence of mammal-eating killer whale encounters was determined by month, year, and subarea to examine patterns of seasonal/annual occurrence and spatial use. Nonparametric Kruskal Wallis tests in the R programming environment (R Core Team 2013) were used to test for differences in the number of occurrences as a function of month, year, and subarea within each period. Minimum group size was determined from sighting records based on the total number of whales with confirmed identifications. The Wilcoxon rank-sum test was used to compare mean and modal group sizes between the two time periods.

Matrilineal groups were classified based on identifications (Towers et al. 2012) in order to examine differences in the groups of whales most commonly occurring in the Salish Sea. There were 28 matrilines identified and defined for this study that were documented more than once in the Salish Sea in the two periods (Appendix 1). For the 1987–1993 period, the most commonly encountered matrilines were classified as nearshore foragers ( $n = 3$ ) or as non-nearshore foragers ( $n = 3$ ), with patterns in seasonal occurrence being one of the differences between the two classifications (Baird and Dill 1995). Changes in seasonal occurrence for these six specific matrilines were compared between the two periods. The patterns of behavior of nearshore versus non-nearshore foragers may be specific to the Salish Sea region, but as this study is a fine-scale analysis of the region the classifications are relevant. For the purposes of our analyses, different matrilines occurring in a given area can be considered analogous to different species present in a community, thus analytical techniques appropriate to community ecology were used in multivariate analyses of matrilineal patterns. The cumulative number of occurrences of each matriline was determined for each season, period, and subarea. Seasons were broken into ecologically relevant periods: August–September (harbor seal pupping–weaning season), October–March, April–May (peak sea lion occurrence season), and June–July. Time periods were broken into 4 sections: 1987–1990, 1991–1993, 2004–2007, and 2008–2010. Season, time period, and subarea were the categorical explanatory variables. Therefore, the multivariate






Figure 1. Subareas of the Salish Sea used for analyses of spatial use. The international boundary is shown. The northernmost area of the Gulf Islands and Strait of Georgia and the westernmost area of the Strait of Juan de Fuca had less small boat and sighting effort, thus were excluded from our analyses.

data set included 28 response variables (number of occurrences for each matriline) within each unique combination of season, time period, and subarea (28 x 48 matrix).

A permutational multivariate analysis of variance test (PERMANOVA) was conducted to examine temporal and spatial patterns in matriline occurrence (Anderson 2001). This test was chosen because it does not have distributional assumptions as does a multivariate analysis of variance test. PERMANOVA is a powerful permutation test that can be used to analyze the effect of categorical explanatory variables on multivariate data. The matriline occurrence distance matrix (based on Bray-Curtis distance measures as is appropriate for zero-inflated data; Bray and Curtis 1957, Faith et al. 1987) was compared to the explanatory variables using PERMANOVA to test for a significant effect of season, time and subarea on the occurrence of the most abundant matriline. Two-way interactions between each explanatory variable were also examined because all were ecologically plausible (e.g., season may have an effect on matrilineal patterns of occurrence in the older period but may not in the more recent period). PERMANOVA was conducted using the *adonis* function in the *vegan* (Oksanen et al. 2008) package of R with 10,000 permutations. Statistical significance of an environmental factor (e.g. subarea) would indicate that the matriline most commonly occurring across different levels of the factor (e.g. the three subareas) were significantly different. The PERMANOVA addresses potential spatial and temporal heterogeneity in the dataset as subareas and time (both seasonally and longitudinally) are explanatory variables in which variability is partitioned. Specifically, this accounts for variable survey and spatial effort between the two time periods for the multivariate analyses.

## Results

Mammal-eating killer whales were encountered more frequently in the 2004–2010 period (339 occurrences, excluding Puget Sound and Hood Canal) than in the 1987–1993 period (97 occurrences; Table 2). Between 1987 and 1993, unique groups of mammal-eating killer whales occurred between

9 and 19 times per year (Figure 2). The number of occurrences did not change by year between 1987 and 1993 (Kruskal Wallis  $\chi^2 = 2.504$ ,  $df = 6$ ,  $P = 0.868$ ). Between 2004 and 2010, unique groups of mammal-eating killer whales occurred in the Salish Sea between 26 and 79 times per year (excluding Puget Sound and Hood Canal), with an increasing number of occurrences in later years (Figure 2). The total number of occurrences in the Salish Sea significantly differed by year between 2004 and 2010 (Kruskal Wallis  $\chi^2 = 18.996$ ,  $df = 6$ ,  $P = 0.004$ ).

The number of occurrences of mammal-eating killer whales was significantly different by subarea for both periods; 1987–1993 (excluding Puget Sound and Hood Canal; Kruskal Wallis  $\chi^2 = 13.632$ ,  $df = 4$ ,  $P = 0.001$ ) and 2004–2010 (including Puget Sound and Hood Canal; Kruskal Wallis  $\chi^2 = 24.693$ ,  $df = 4$ ,  $P < 0.001$ ). Mammal-eating killer whales were more frequently observed in the Strait of Juan de Fuca than in any other subarea of the Salish Sea throughout this study. Other subareas of the Salish Sea (e.g., Puget Sound) had a slight increasing trend in annual occurrence of mammal-eating killer whales from 2004 to 2010 (Figure 2).

Occurrence of mammal-eating killer whales differed significantly by month in 1987–1993 (Kruskal Wallis  $\chi^2 = 42.671$ ,  $df = 11$ ,  $P < 0.001$ ) and in 2004–2010 (Kruskal Wallis  $\chi^2 = 23.795$ ,  $df = 11$ ,  $P = 0.014$ ). From 1987–1993, whales occurred most frequently in August–September (Figure 3). From 2004–2010, whales also occurred most frequently in August–September, but there was also a strong second peak in April–May (Figure 3).

The most commonly observed group size increased between the 1987–1993 (mode = 3; see Figure 3 in Baird and Dill 1996) and 2004–2010 periods (mode = 4, Figure 4), although the difference was not statistically significant ( $W = 18885.5$ ,  $P = 0.231$ ). The mean group size in the 2004–2010 period (mean  $\pm$  SD:  $5.17 \pm 4.36$ , range: 1 to 36) was greater than the mean group size in the 1987–1993 period (mean  $\pm$  SD:  $4.40 \pm 2.82$ , range: 1 to 15).

TABLE 2. Number of occurrences of mammal-eating killer whales in the San Juan Islands, Gulf Islands and Strait of Georgia, and Strait of Juan de Fuca subareas by year and by month from 1987–2010. Number of occurrences in the Puget Sound and Hood Canal subareas are noted in parentheses.

Month	Year												Total																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																						
	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2423	2424	2425	2426	2427	2428	2429	2430	2431	2432	2433	2434	2435	2436	2437	2438	2439	2440	2441	2442	2443	2444	2445	2446	2447	2448	2449	2450	2451	2452	2453	2454	2455	2456	2457	2458	2459	2460	2461	2462	2463	2464	2465	2466	2467	2468	2469	2470	2471	2472	2473	2474	2475	2476	2477	2478	2479	2480	2481	2482	2483	2484	2485	2486	2487	2488	2489	2490	2491	2492	2493	2494	2495	2496	2497	2498	2499	2500	2501	2502	2503	2504	2505	2506	2507	2508	2509	2510	2511	2512	2513	2514	2515	2516	2517	2518	2519	2520	2521	2522	2523	2524	2525	2526	2527	2528	2529	2530	2531	2532	2533	2534	2535	2536	2537	2538	2539	2540	2541	2542	2543	2544	2545	2546	2547	2548	2549	2550	2551	2552	2553	2554	2555	2556	2557	2558	2559	2560	2561	2562	2563	2564	2565	2566	2567	2568	2569	2570	2571	2572	2573	2574	2575	2576	2577	2578	2579	2580	2581	2582	2583	2584	2585	2586	2587	2588	2589	2590	2591	2592	2593	2594	2595	2596	2597	2598	2599	2600	2601	2602	2603	2604	2605	2606	2607	2608	2609	2610	2611	2612	2613	2614	2615	2616	2617	2618	2619	2620	2621	2622	2623	2624	2625	2626	2627	2628	2629	2630	2631	2632	2633	2634	2635	2636	2637	2638	2639	2640	2641	2642	2643	2644	2645	2646	2647	2648	2649	2650	2651	2652	2653	2654	2655	2656	2657	2658	2659	2660	2661	2662	2663	2664	2665	2666	2667	2668	2669	2670	2671	2672	2673	2674	2675	2676	2677	2678	2679	2680	2681	2682	2683	2684	2685	2686	2687	2688	2689	2690	2691	2692	2693	2694	2695	2696	2697	2698	2699	2700	2701	2702	2703	2704	2705	2706	2707	2708	2709	2710	2711	2712	2713	2714	2715	2716	2717	2718	2719	2720	2721	2722	2723	2724	2725	2726	2727	2728	2729	2730	2731	2732	2733	2734	2735	2736	2737	2738	2739	2740	2741	2742	2743	2744	2745	2746	2747	2748	2749	2750	2751	2752	2753	2754	2755	2756	2757	2758	2759	2760	2761	2762	2763	2764	2765	2766	2767	2768	2769	2770	2771	2772	2773	2774	2775	2776	2777	2778	2779	2780	2781	2782	2783	2784	2785	2786	2787	2788	2789	2790	2791	2792	2793	2794	2795	2796	2797	2798	2799	2800	2801	2802	2803	2804	2805	2806	2807	2808	2809	2810	2811	2812	2813	2814	2815	2816	2817	2818	2819	2820	2821	2822	2823	2824	2825	2826	2827	2828	2829	2830	2831	2832	2833	2834	2835	2836	2837	2838	2839	2840	2841	2842	2843	2844	2845	2846	2847	2848	2849	2850	2851	2852	2853	2854	2855	2856	2857	2858	2859	2860	2861	2862	2863	2864	2865	2866	2867	2868	2869	2870	2871	2872	2873	2874	2875	2876	2877	2878	2879	2880	2881	2882	2883	2884	2885	2886	2887	2888	2889	2890	2891	2892	2893	2894	2895	2896	2897	2898	2899	2900	2901	2902	2903	2904	2905	2906	2907	2908	2909	2910	2911	2912	2913	2914	2915	2916	2917	2918	2919	2920	2921	2922	2923	2924	2925	2926	2927	2928	2929	2930	2931	2932	2933	2934	2935	2936	2937	2938	2939	2940	2941	2942	2943	2944	2945	2946	2947	2948	2949	2950	2951	2952	2953	2954	2955	2956	2957	2958	2959	2960	2961	2962	2963	2964	2965	2966	2967	2968	2969	2970	2971	2972	2973	2974	2975	2976	2977	2978	2979	2980	2981	2982	2983	2984	2985	2986	2987	2988	2989	2990	2991	2992	2993	2994	2995	2996	2997	2998	2999	3000	3001	3002	3003	3004	3005	3006	3007	3008	3009	3010	3011	3012	3013	3014	3015	3016	3017	3018	3019	3020	3021	3022	3023	3024	3025	3026	3027	3028	3029	3030	3031	3032	3033	3034	3035	3036	3037	3038	3039	3040	3041	3042	3043	3044	3045	3046	3047	3048	3049	3050	3051	3052	3053	3054	3055	3056	3057	3058	3059	3060	3061	3062	3063	3064	3065	3066	3067	3068	3069	3070	3071	3072	3073	3074	3075	3076	3077	3078	3079	3080	3081	3082	3083	3084	3085	3086	3087	3088	3089	3090	3091	3092	3093	3094	3095	3096	3097	3098	3099	3100	3101	3102	3103	3104	3105	3106	3107	3108	3109	3110	3111	3112	3113	3114	3115	3116	3117	3118	3119	3120	3121	3122	3123	3124	3125	3126	3127	3128	3129	3130	3131	3132	3133	3134	3135	3136	3137	3138	3139	3140	3141	3142	3143	3144	3145	3146	3147	3148	3149	3150	3151	3152	3153	3154	3155	3156	3157	3158	3159	3160	3161	3162	3163	3164	3165	3166	3167	3168	3169	3170	3171	3172	3173	3174	3175	3176	3177	3178	3179	3180	3181	3182	3183	3184	3185	3186	3187	3188	3189	3190	3191	3192	3193	3194	3195	3196	3197	3198	3199	3200	3201	3202	3203	3204	3205	3206	3207	3208	3209	3210	3211	3212	3213	3214	3215	3216	3217	3218	3219	3220	3221	3222	3223	3224	3225	3226	3227	3228	3229	3230	3231	3232	3233	3234	3235	3236	3237	3238	3239	3240	3241	3242	3243	3244	3245	3246	3247	3248	3249	3250	3251	3252	3253	3254	3255	3256	3257	3258	3259	3260	3261	3262	3263	3264	3265	3266	3267	3268	3269	3270	3271	3272	3273	3274	3275	3276	3277	3278	3279	3280	3281	3282	3283	3284	3285	3286	3287	3288	3289	3290	3291	3292	3293	3294	3295	3296	3297	3298	3299	3300	3301	3302	3303	3304	3305	3306	3307	3308	3309	3310	3311	3312	3313	3314	3315	3316	3317	3318	3319	3320	3321	3322	3323	3324	3325	3326	3327	3328	3329	3330	3331	3332	33




Figure 2. Cumulative annual occurrence of mammal-eating killer whales in the Salish Sea by subarea from 1987–1993 and 2004–2010.




Figure 3. Cumulative monthly occurrence of mammal-eating killer whales in the Salish Sea by subarea from 1987–1993 and 2004–2010.






Figure 4. Percent of occurrences for each observed minimum group size of mammal-eating killer whales in the Salish Sea from 1987–1993 and 2004–2010. From 1987–1993, modal group size was 3 whales (mean  $\pm$  SD:  $4.40 \pm 2.82$ ). From 2004–2010, modal group size was 4 whales ( $5.17 \pm 4.36$ ).

The most commonly occurring matriline from the 1987–1993 period also occurred more frequently in the 2004–2010 period (Table 3). However, there were several other matrilines that were not documented in the study in 1987–1993 that were common from 2004–2010 (Table 3). Differences in seasonal occurrence between matrilines existed in both periods. For the 1987–1993 period, the three matrilines classified as nearshore foragers had a more pronounced August–September peak, while the three matrilines classified as non-nearshore foragers were encountered relatively consistently throughout the year (Figure 5; see Baird and Dill 1995). For the 2004–2010 period, the same matrilines of nearshore foragers still had a peak in occurrence from August–September, but the matrilines of non-nearshore foragers had peaks in occurrence in April–May and August–September (Figure 5). The non-nearshore foragers increased in overall occurrence between the two time periods to a greater extent than the nearshore foragers increased.

The matrilines occurring in each season ( $F = 2.300$ ,  $df_1 = 3$ ,  $df_2 = 18$ ,  $P < 0.001$ ), time period ( $F = 4.281$ ,  $df_1 = 3$ ,  $df_2 = 18$ ,  $P < 0.001$ ), and subarea of the Salish Sea ( $F = 2.796$ ,  $df_1 = 2$ ,  $df_2 = 18$ ,  $P = 0.002$ ) differed significantly (PERMANOVA  $R^2 = 0.74$ ). Matrilineal occurrence in each area differed across time periods with marginal significance ( $F = 1.336$ ,  $df_1 = 6$ ,  $df_2 = 18$ ,  $P = 0.089$ ). However, the

matrilines that commonly occurred in each season did not differ across time periods or across areas.

## Discussion

Some of the increase in mammal-eating killer whales' use of the Salish Sea between 1987 and 2010 is likely due to an increase in sighting effort between the two periods. However, we attempted to control for differences in effort by looking at occurrences of groups in the Salish Sea (i.e., taking into account repeated sightings over six days). We also only included the three subareas that had sighting effort in both time periods when comparing 1987–1993 to 2004–2010. The number of occurrences of mammal-eating killer whales also increased significantly within the 2004–2010 period itself, when sighting effort was relatively consistent. It is relevant to note that the measure of occurrences may affect spatial analyses by underrepresenting certain subareas (e.g., those that are utilized after arrival in the Salish Sea via one of the straits). Although mammal-eating killer whales most commonly occurred in the Strait of Juan de Fuca, they also increased in occurrence in other subareas (e.g., Puget Sound) in recent years. We suggest this apparent increase in occurrence over time in the Salish Sea is due to an increase in the abundance of mammal-eating killer whale primary prey, as seals, sea lions, and porpoises have increased substantially in abundance and

TABLE 3. Number of occurrences of at least one member of each matriline in each period (see Appendix 1 for matriline designation codes). The total number of occurrences does not include occurrence in the Puget Sound or Hood Canal (noted separately in 2004–2010 column). Six of the commonly occurring matriline in 1987–1993 were classified as either nearshore foragers or non-nearshore foragers by Baird and Dill (1995).

Matriline	1987–1993 # of Occurrences	2004–2010 # of Occurrences (Puget Sound and Hood Canal Occurrences)	Total # of Occurrences	Classification (Baird and Dill 1995)
T21	17	47 (3)	64	Non-nearshore forager
T13	14	40	54	Non-nearshore forager
T41	16	27	43	Nearshore forager
T100	0	43 (1)	43	
T2	19	20 (1)	39	Non-nearshore forager
T88	3	34 (6)	37	
T124	0	35 (3)	35	
T10	15	20	35	Nearshore forager
T32	24	10	34	Nearshore forager
T30	0	33 (2)	33	
T101	0	31 (6)	31	
T19	0	24 (2)	24	
T49	0	22	22	
T46	0	22 (1)	22	
T11	12	8	20	
T109	12	5	17	
T104	0	16	16	
T36	0	12 (2)	12	
T12	10	2	12	
T65	0	11 (2)	11	
T123	0	6	6	
T124A	0	5 (6)	5	
T99	0	5	5	
T7	2	3	5	
T75	4	1	5	
T137	0	4 (3)	4	
T71	0	3 (6)	3	
T185	0	3	3	

occurrence in the area (Jeffries et al. 2003, Lowry and Maravilla-Chavez 2005, Pitcher et al. 2007, Fisheries and Oceans Canada 2010, Carretta et al. 2011). The increased local occurrence in the Salish Sea could also be due to an increase in abundance of the greater population of west coast ‘transient’ killer whales throughout the study period (Ford et al. 2007, Towers et al. 2012). These causative

links are not mutually exclusive, and it is likely that both prey abundance and overall population growth have contributed to the changes in occurrence documented in the Salish Sea.

Seasonal patterns of occurrence differed between the two periods with a single peak in occurrence of mammal-eating killer whales in August–September from 1987–1993 and two




Figure 5. Cumulative monthly occurrence of non-nearshore and nearshore foragers from 1987–1993 and 2004–2010.

peaks in 2004–2010, one in April–May and one in August–September. We suspect these changes in seasonal occurrence could be a consequence of increased prey diversity. From 1987–1993, half of the most commonly observed mammal-eating killer whale matriline were found to be nearshore foragers, specializing on harbor seal prey near haul-out sites during the pupping-weaning season in inland waters (Bigg 1969, Temte et al. 1991, Baird and Dill 1995). It is likely that the increased use of the Salish Sea was by non-nearshore foragers that did not focus hunting effort near harbor seal pupping and haul-out sites, but still foraged on harbor seals throughout the year. Other prey species, including porpoises, elephant seals, and sea lions, are all found in areas away from harbor seal haul

outs. Although sea lions and occasionally elephant seals also utilize haul out sites, killer whales seem less likely to attack sea lions at haul outs, as attacks are often prolonged and sea lions may be able to reach refuge sites in shallow water or on shore if attacked nearshore (Baird and Dill 1995). The second peak in mammal-eating killer whale occurrence in April–May during 2004–2010 corresponds with the increased occurrence of California sea lions and Steller sea lions in inland waters in those months, suggesting the whales may be taking advantage of the increased occurrence of these pinnipeds. The lower occurrence of mammal-eating killer whales in the Salish Sea in June–July may be associated with seasonal fluctuations in California sea lions and Steller sea lions as many of these pinnipeds migrate to

the outer coast to return to natal rookeries at that time. Peaks in occurrence in cetacean prey species (Dahlheim et al. 2009) and differences in timing of the peak harbor seal pupping period (Temte et al. 1991) elsewhere throughout the range of mammal-eating killer whales likely also influence killer whale movements to areas outside of the Salish Sea in winter and in the months June–July.

The most commonly occurring group size of mammal-eating killer whales in the Salish Sea had an increasing trend from 1987 to 2010, but this was not statistically significant. This pattern could be due to increased prey diversity as optimum foraging group size changes by prey species, depending in part on the difficulty of capture, risk of injury, and the energetic value of the prey (Baird and Dill 1996). When feeding primarily on harbor seals, the optimum foraging group size was three individuals, the most frequently observed group size in the 1987–1993 period (Baird and Dill 1996). Although we have not been able to quantify it, there is anecdotal evidence of increased predation on faster and more difficult to catch prey species, such as harbor porpoise, and larger prey species, such as sea lions, in our study area in recent years; from observations both in southeast Alaska and in our study area faster and larger prey species are targeted by larger groups of mammal-eating killer whales (Baird and Dill 1995, Dahlheim and White 2010). Mammal-eating killer whales are cooperative hunters with the cost of prey capture varying with group size. Higher encounter rates with increasingly abundant sea lions and porpoises in the Salish Sea facilitate the larger groups recently observed (Baird 2002). Increased prey abundance in the Salish Sea could relax the need to forage in the optimum group size as efficiency becomes less important when there are more foraging opportunities available. Increased prey abundance also likely leads to increased consumption, growth, and reproduction by mammal-eating killer whales. Increased reproduction in this population is consistent with the observation of several commonly occurring matrilineal lines having many young, undispersed offspring in recent years (Towers et al. 2012; e.g., T100, T101, and T124 matrilineal lines; Table 3). Therefore, increased reproductive output due to increased

prey abundance could also lead to the larger group sizes recently observed in the Salish Sea.

Matrilineal differences in annual occurrence contributed to the pattern of increasing use of the Salish Sea over time. Many of the commonly observed matrilineal lines from 1987–1993 increased their use of the area in 2004–2010, and several matrilineal lines previously not encountered in the Salish Sea were also observed frequently between 2004 and 2010. Our results suggest that the changes in annual occurrence are explained by different matrilineal lines utilizing the area in recent years compared to those that were common in 1987–1993. The matrilineal lines that occurred commonly in 2004–2010 but were not encountered in 1987–1993 may have increased their use of the Salish Sea where prey abundance has increased, in addition to utilizing prey resources elsewhere. Several of these matrilineal lines were encountered in southeast Alaska frequently within the 1987–1993 period (Dahlheim and White 2010). In the inland waters of southeast Alaska, harbor porpoise populations have declined significantly from 1991 to 2011 (Marilyn Dahlheim, NOAA National Marine Mammal Laboratory, personal communication) and one of several stocks of harbor seals has also declined from 1992 to 2008 (Womble et al. 2010). The recent changes in mammal-eating killer whale occurrence in the Salish Sea may indicate that these whales are utilizing a broader range of their distribution to take advantage of areas exhibiting increased prey abundance.

Matrilineal differences can explain the changes in seasonal occurrence between the two time periods. Non-nearshore foragers, which did not focus foraging effort on harbor seal pupping and haul-out sites in 1987–1993, increased in overall occurrence to a greater extent than did nearshore foragers. The divergent pattern between the two foraging types contributed to the greater seasonal occurrence of killer whales in times of the year that do not coincide with harbor seal pupping–weaning time. The PERMANOVA results indicated that seasonal differences in occurrence could be attributed to the occurrence of different matrilineal lines. It is possible that the more recently common matrilineal lines in the Salish Sea in 2004–2010 exhibit the same seasonal patterns as non-nearshore foragers, contributing to

the changes in seasonal occurrence between 1987 and 2010. However, the classification of nearshore and non-nearshore foragers may be specific to the Salish Sea region and may differ from broad-scale patterns of occurrence and behavior.

These new findings update our previous understanding of mammal-eating killer whale spatial use, occurrence and group size in the Salish Sea. Mammal-eating killer whales are more common at different times of the year and have also increased in their occurrence and group size. Current information on top predator geospatial ecology will better inform ecosystem-based models in the Salish Sea, and provide insights for marine populations in other systems. Increasing knowledge of predator-prey relationships in the Salish Sea allows for a greater understanding of the role of mammal-eating killer whales in this ecosystem. Findings from this study provide an example of how predators might be altering their spatial use and seasonal occurrence in order to take advantage of increasing prey abundance and occurrence. Other studies have shown that an increase in prey abundance alters the diet and abundance of marine predators (Womble and Sigler 2006), but none have examined changes in aspects of behavior such as occurrence and group size. It is likely that mammal-eating killer whales are also having non-consumptive effects on their prey populations due to the changes in whale behavior and increased use of the Salish

Sea in recent years. For example, prey populations could increase predator vigilance in months where mammal-eating killer whales are increasing seasonal use, which would increase the total effect of the predator in this system. Intraspecific variation in occurrence and spatial use by different matriline is also a unique aspect of this study and will be important to consider in future studies in order to determine how individuals or matriline may differ in other behavioral traits or patterns.

## Acknowledgments

Funding for analyses of the 2004–2010 data was provided by the Northwest Fisheries Science Center (NWFSC) through the Undergraduate Research Program at the University of Washington. The map of the Salish Sea was produced and provided by D. Holzer, NWFSC. The authors would like to thank Orca Network, The Whale Museum, members of the public, whale-watch operators, and researchers, who contributed sightings of killer whales in the Salish Sea. RWB would like to thank P. Stacey and T. Guenther for assistance with data collection in 1987–1993. M. Dahlheim provided early feedback on the 2004–2010 analyses. J. Bakker and K. MacIntyre provided guidance on analytical methods for examining matrilineal patterns of occurrence. D. Noren, A. Wirsing and four anonymous reviewers provided comments that greatly improved this manuscript.

## Literature Cited

- Ainley, D. G., G. Ballard, and K. M. Dugger. 2006. Competition among penguins and cetaceans reveals trophic cascades in the western Ross Sea, Antarctica. *Ecology* 87:2080–2093.
- Allen, B. M., and R. P. Angliss. 2011. Alaska marine mammal stock assessments, 2010. U.S. Department of Commerce, NOAA Technical Memorandum. NMFS AFSC-223.
- Anderson, M. J. 2001. A new method for non-parametric multivariate analysis of variance. *Austral Ecology* 26:32–46.
- Baird, R. W., P. A. Abrams, and L. M. Dill. 1992. Possible indirect interactions between transient and resident killer whales: implications for the evolution of foraging specializations in the genus *Orcinus*. *Oecologia* 89:125–132.
- Baird, R. W., and L. M. Dill. 1995. Occurrence and behavior of transient killer whales: seasonal and pod-specific variability, foraging behavior and prey handling. *Canadian Journal of Zoology* 73:1300–1311.
- Baird, R. W., and L. M. Dill. 1996. Ecological and social determinants of group size in transient killer whales. *Behavioral Ecology* 7:408–416.
- Baird, R. W., and H. Whitehead. 2000. Social organization of mammal-eating killer whales: group stability and dispersal patterns. *Canadian Journal of Zoology* 78:2096–2105.
- Baird, R. W. 2002. *Killer Whales of the World: Natural History and Conservation*. Voyageur Press, Stillwater, MN.
- Baum, J. K., R. A. Myers, D. G. Kehler, B. Worm, S. J. Harley, and P. A. Doherty. 2003. Collapse and conservation of shark populations in the Northwest Atlantic. *Science* 299:389–392.
- Bigg, M. A. 1969. Clines in the pupping season of the harbour seal, *Phoca vitulina*. *Journal of Fisheries Research Board of Canada* 26:449–455.



- Bigg, M. A., G. M. Ellis, J. K. B. Ford, and K. C. Balcomb. 1987. Killer Whales—A Study of Their Identification, Genealogy and Natural History in British Columbia and Washington State. Phantom Press, Nanaimo, BC.
- Bigg, M. A., P. F. Olesiuk, G. M. Ellis, J. K. B. Ford, and K. C. Balcomb. 1990. Social organization and genealogy of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. Report of the International Whaling Commission Special Issue 12:386-406.
- Bray, J. R., and J. T. Curtis. 1957. An ordination of upland forest communities in southern Wisconsin. Ecological Monographs 27:325-349.
- Carretta, J. V., K. A. Forney, E. Oleson, K. Martien, M. M. Muto, M. S. Lowry, J. Barlow, J. Baker, B. Hanson, D. Lynch, L. Carswell, R. L. Brownell Jr., J. Robbins, D. K. Mattila, K. Ralls, and M. C. Hill. 2011. U.S. Pacific Marine Mammal Stock Assessments: 2010. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-SWFSC-476.
- Dahlheim, M. E., P. A. White, and J. M. Waite. 2009. Cetaceans of Southeast Alaska: distribution and seasonal occurrence. Journal of Biogeography 36:410-426.
- Dahlheim, M. E., and P. A. White. 2010. Ecological aspects of transient killer whales *Orcinus orca* as predators in southeastern Alaska. Wildlife Biology 16:308-322.
- Demaster, D. P., A. W. Trites, P. Clapham, S. Mizroch, P. Wade, R. J. Small, and J. Ver Hoef. 2006. The sequential megafaunal collapse hypothesis: Testing with existing data. Progress in Oceanography 68:329-342.
- Estes, J. A., and J. F. Palmisano. 1974. Sea otters: their role in structuring nearshore communities. Science 185:1058-1060.
- Estes, J. A., M. T. Tinker, T. M. Williams, and D. F. Doak. 1998. Killer whale predation on sea otters linking oceanic and nearshore ecosystems. Science 282:473-476.
- Estes, J. A., J. Terborgh, J. S. Brashares, M. E. Power, J. Berger, W. J. Bond, S. R. Carpenter, T. E. Essington, R. D. Holt, J. B. C. Jackson, R. J. Marquis, L. Oksanen, R. T. Paine, E. K. Pikitch, W. J. Ripple, S. A. Sandin, M. Scheffer, T. W. Schoener, J. B. Shurin, A. R. E. Sinclair, M. E. Soulé, R. Virtanen, and D. A. Wardle. 2011. Trophic downgrading of planet Earth. Science 333:301-306.
- Faith, D. P., P. R. Minchin, and L. Belbin. 1987. Compositional dissimilarity as a robust measure of ecological distance. Vegetation 69:57-68.
- Fisheries and Oceans Canada. 2010. Population Assessment Pacific harbour seal (*Phoca vitulina richardsi*). DFO Canadian Science Advisory Secretariat Science Advisory Report 2009/011.
- Ford, J. K. B., and G. M. Ellis. 1999. Transients: Mammal-hunting Killer Whales. UBC Press, Vancouver, BC.
- Ford, J. K. B., G. M. Ellis, L. Barrett-Lennard, A. B. Morton, R. S. Palm, and K. C. Balcomb. 1998. Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in coastal British Columbia and adjacent waters. Canadian Journal of Zoology 76:1456-1471.
- Ford, J. K. B., G. M. Ellis, D. R. Matkin, K. C. Balcomb, D. Briggs, and A. B. Morton. 2006. Killer whale attacks on minke whales: prey capture and antipredator tactics. Marine Mammal Science 21:603-618.
- Ford, J. K. B., G. M. Ellis, and J. W. Durban. 2007. An assessment of the potential for recovery of west coast transient killer whales using coastal waters of British Columbia. Fisheries and Oceans Canada: Research Document 2007/088.
- Giles, D. A., and K. L. Koski. 2012. Managing vessel-based killer whale watching: a critical assessment of the evolution from voluntary guidelines to regulations in the Salish Sea. Journal of International Wildlife Law & Policy 15:125-151.
- Harvey, C. J., K. K. Bartz, J. R. Davies, T. B. Francis, T. P. Good, A. D. Guerry, M. B. Hanson, K. K. Holsman, J. Miller, M. Plummer, J. C. Reum, L. D. Rhodes, C. A. Rice, J. F. Samhour, G. D. Williams, N. J. Yoder, P. S. Levin, and M. H. Ruckelshaus. 2010. A mass-balance model for evaluating food web structure and community-scale indicators in the central basin of Puget Sound. U.S. Department of Commerce, NOAA Technical Memorandum NMFS-NWFSC-106.
- Hauser, D. D. W., G. R. VanBlaricom, E. E. Holmes, and R. W. Osborne. 2006. Evaluating the use of whale-watch data in determining killer whale (*Orcinus orca*) distribution patterns. Journal of Cetacean Research and Management 8:273-281.
- Hauser, D. D. W., M. G. Logsdon, E. E. Holmes, G. R. VanBlaricom, and R. W. Osborne. 2007. Summer distribution patterns of southern resident killer whales *Orcinus orca*: core areas and spatial segregation of social groups. Marine Ecology Progress Series 351:301-310.
- Heithaus, M. R., A. Frid, A. J. Wirsing, L. M. Dill, J. W. Fouquerean, D. Burkholder, J. Thomson, and L. Bejder. 2007. State-dependent risk-taking by green sea turtles mediates top-down effects of tiger shark intimidation in a marine ecosystem. Journal of Animal Ecology 76:837-844.
- Janzen, D. H. 1974. Deflowering of Central-America. Natural History 83:48-53.
- Jeffries, S. J., H. R. Huber, J. Calambokidis, and J. Laake. 2003. Trends and status of harbor seals in Washington state: 1978–1999. Journal of Wildlife Management 67:207-218.
- Lima, S. L. 1998. Nonlethal effects in the ecology of predator-prey interactions. Bioscience 48:25-34.
- London, J. M., J. M. Ver Hoef, S. J. Jeffries, M. M. Lance, and P. L. Boveng. 2012. Haul-out behavior of harbor

- seals (*Phoca vitulina*) in Hood Canal, Washington. PLoS One 7(6): e38180.
- Lowry, M. S., and O. Maravilla-Chavez. 2005. Recent abundance of California sea lions in western Baja California, Mexico and the United States. In D. K. Garcelon and C. A. Schwemm (editors), National Park Service Technical Publication CHIS-05-01. Institute for Wildlife Studies, Arcata, CA. Pp. 485-497.
- May, R. M., J. R. Beddington, C. W. Clark, S. J. Holt, and R. M. Laws. 1979. Management of multispecies fisheries. Science 205:267-277.
- McLaren, B. E., and R. O. Peterson. 1994. Wolves, Moose, and Tree Rings on Isle Royale. Science 266:1555-1558.
- Murdoch, T. J. T., A. F. Glasspool, M. Outerbridge, J. Ward, S. Manuel, J. Gray, A. Nash, K. A. Coates, J. Pitt, J. W. Fourqurean, P. A. Barnes, M. Vierros, K. Holzer, and S. R. Smith. 2007. Large-scale decline in offshore seagrass meadows in Bermuda. Marine Ecology Progress Series 339:123-130.
- Oksanen, J., R. Kindt, P. Legendre, B. O'Hara, G. L. Simpson, P. Solymos, M. H. M. Stevens, and H. Wagner. 2008. Vegan: Community Ecology Package. R Package Version 1, 15.11. Available from <http://vegan.r-forge.r-project.org>.
- Orca Network. 2011. Orca Network sightings archives. Orca Network, Whale Sighting Network Project, Freeland, WA. Available online at [http://www.orcanetwork.org/Archives/index.php?categories\\_file=Sightings%20Archives%20Home](http://www.orcanetwork.org/Archives/index.php?categories_file=Sightings%20Archives%20Home) (accessed 1 February 2011).
- Pitcher, K. W., P. F. Olesiuk, R. F. Brown, M. S. Lowry, S. J. Jeffries, J. L. Sease, W. L. Perryman, C. E. Stinchcomb, and L. F. Lowry. 2007. Abundance and distribution of the eastern North Pacific Steller sea lion (*Eumetopias jubatus*) population. Fisheries Bulletin 107:102-115.
- Puget Sound Action Team. 2007. 2007 Puget Sound Update: Ninth Report of the Puget Sound Assessment and Monitoring Program. Puget Sound Action Team. Olympia, WA.
- R Core Team. 2013. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.
- Sheriff, M. J., C. J. Krebs, and R. Boonstra. 2009. The sensitive hare: sublethal effects of predator stress on reproduction in snowshoe hares. Journal of Animal Ecology 78:1249-1258.
- Soloman, M. E. 1949. The natural control of animal populations. Journal of Animal Ecology 19:1-35.
- Springer, A. M., J. A. Estes, G. B. van Vliet, T. M. Williams, D. F. Doak, E. M. Danner, K. A. Forney, and B. Pfister. 2003. Sequential megafaunal collapse in the North Pacific Ocean: An ongoing legacy of industrial whaling? Proceedings of the National Academy of Science 100:12223-12228.
- Temte, J. L., M. A. Bigg, and Ø. Wiig. 1991. Clines revisited: The timing of pupping in the harbour seal (*Phoca vitulina*). Journal of Zoology 224:617-632.
- Towers, J. R., G. M. Ellis, and J. K. B. Ford. 2012. Photo-identification catalogue of Bigg's (transient) killer whales from coastal waters of British Columbia, northern Washington, and southeastern Alaska. Canadian Data Report of Fisheries and Aquatic Sciences 1241: v + 127 pp.
- Wade, P. R., V. N. Burkanov, M. E. Dahlheim, N. A. Friday, L. W. Fritz, T. R. Loughlin, S. A. Mizroch, M. M. Muto, D. W. Rice, L. G. Barrett-Lennard, N. A. Black, A. M. Burdin, J. Calambokidis, S. Cerchio, J. K. B. Ford, J. K. Jacobsen, C. O. Matkin, D. R. Matkin, A. V. Mehta, R. J. Small, J. M. Straley, S. M. McCluskey, G. R. VanBlaricom, and P. J. Clapham. 2007. Killer whales and marine mammal trends in the North Pacific—a re-examination of evidence for sequential megafauna collapse and the prey-switching hypothesis. Marine Mammal Science 23:766-802.
- Whipple, S. J., J. S. Link, L. P. Garrison, and M. J. Fogarty. 2000. Models of predation and fishing mortality in aquatic ecosystems. Fish and Fisheries 1:22-40.
- Wirsing, A. J., M. R. Heithaus, and L. M. Dill. 2007. Fear factor: do dugongs (*Dugong dugon*) trade food for safety from tiger sharks (*Galeocerdo cuvier*)? Oecologia 153:1031-1040.
- Wirsing, A. J., M. R. Heithaus, A. Frid, and L. M. Dill. 2008. Seascapes of fear: evaluating sublethal predator effects experienced and generated by marine mammals. Marine Mammal Science 24:1-15.
- Wirsing, A. J., and W. J. Ripple. 2010. A comparison of shark and wolf research reveals similar behavioral responses by prey. Frontiers in Ecology and the Environment 9:335-341.
- Womble, J. N., and M. F. Sigler. 2006. Seasonal availability of abundant, energy-rich prey influences the abundance and diet of a marine predator, the Steller sea lion *Eumetopias jubatus*. Marine Ecology Progress Series 325:281-293.
- Womble, J. N., G. W. Pendleton, E. A. Mathews, G. M. Blundell, N. M. Bool, and S. M. Gende. 2010. Harbor seal (*Phoca vitulina richardii*) decline continues in the rapidly changing landscape of Glacier Bay National Park, Alaska 1992–2008. Marine Mammal Science 26:686-697.

Received 13 July 2013

Accepted for publication 25 January 2015

## Appendix 1

List of matriline codes and matriline membership used in this study

T2	T2 (female), T1, T2A, T2B, T2C
T7	T7 (female), T7A, T7B
T10	T10 (female), T10A, T10B, T10C
T11	T11 (female), T11A
T12	T12 (female), T12A, T12B, T12C
T13	T13 (female), T14
T19	T19 (female), T19B, T19C
T21	T21 (female), T20, T22
T30	T30 (female), T30A, T30B, T30C
T32	T32 (female), T31
T36	T36 (female), T36A, T36B, T36C
T41	T41 (female), T41A, T44
T46	T46 (female), T46A, T46B, T46C, T46D, T46E
T49	T49 (female), T49A, T49B
T65	T65 (female), T63, T65A, T65B
T71	T71 (female), T71A, T71B, T71C
T75	T75 (female), T77, T78, T75A, T75B, T75C
T88	T88 (female), T87, T90
T99	T99 (female), T99A, T99B
T100	T100 (female), T100A, T100B, T100C, T100D
T101	T101 (female), T101A, T101B, T102
T104	T104 (female), T103
T109	T109 (female), T109A, T109B, T109C, T109D
T123	T123 (female), T123A
T124	T124 (female), T124C, T124D, T124E
T124A	T124A (female), T124A1, T124A2, T124A3
T137	T137 (female), T137A, T137B
T185	T185 (female), T186, T187, T185A