

HistoPlex®

The Immune System

The immune system normally discriminates between “self” and “non-self” foreign substances. Thus, it represents an elaborate, finely tuned defense system to destroy microorganisms, cancer cells, or other invasive materials. However, an imbalanced immune system can attack the body’s own tissues (auto-immune conditions) and lead to cumulative oxidative stress.

The immune system consists of two branches, cellular immunity and humoral immunity. The first depends upon the active participation of multiple cell types, including a variety of phagocytic cells, to engulf invaders. Lymphocytes include highly specialized T cells, which proliferate and differentiate in the thymus gland. In a typical scenario, macrophages process ingested antigens and display recognition fragments on their surfaces. T helper cells acting as generals, read these antigens and in turn stimulate the production of T killer cells via lymphokines such as IL-2 and gamma interferon. Mast cells are a type of T cell embedded in tissues to fight localized infection. When they contact foreign materials, mast cells destroy them. In the process, they release histamine and other agents that cause edema, increase blood vessel permeability and recruit neutrophils to the site of inflammation.

Humoral immunity pertains to proteins of the blood and lymph. This aspect of the immune system relies on B cells which mature to plasma cells that secrete specific antibodies and complement proteins to fight infections. The complement system also triggers localized inflammation. T helper cells regulate the production and maturation of B cells through lymphokines. T suppressor cells gear down the immune response and thus are critical in allergy attacks and autoimmune conditions.

Immunity is a hallmark of the immune system. Recovery from an invader renders the individual immune to a subsequent attack. The underlying mechanism involves memory T cells and memory B cells. These memory cells multiply rapidly when they reencounter the stimulating antigen.

Allergies

An allergic response involves both cellular and humoral mechanisms. In contrast, other types of sensitivities to foods, such as lactose intolerance or sulfite sensitivity, do not employ the immune system. Allergic responses can be generally classified as immediate or delayed hypersensitivities. Immediate hypersensitivity represents a rapid over-reaction of the immune system, often within minutes after exposure to the problem antigen. Particularly mast cells release histamine, leukotrienes and other stimuli that evoke swelling, hives, itchiness, copious mucus secretion and muscle spasm of the GI tract and bronchioles. The reaction is mediated by IgE antibodies. Anaphylactic shock results from an immediate hypersensitive allergic reaction affecting the whole body. Depending on how food allergy is defined, estimates of the prevalence range from 2% to 25% of the US population. Immediate

allergic reactions account for only a small fraction of food allergies. Typical food allergies frequently involve IgG antibodies, and symptoms develop within hours or days after contact with a problematic food. Faulty digestion and increased gut permeability are often a root cause of multiple food allergies, because food allergens that are normally excluded, cross the intestinal barrier and enter the bloodstream. The resulting immune complexes, if not cleared rapidly, can lodge in various organs and tissues where they can trigger inflammatory conditions. Food allergies have been linked to a range of autoimmune conditions, rheumatoid arthritis, for example.⁽¹⁾

Detecting food allergies can be perplexing. The delay between ingesting a specific food and the development of sometimes vague symptoms often masks food allergy reactions. The most frequent symptom is fatigue. Other symptoms include delayed hypersensitivity, irritability, achy joints, blood sugar dysregulation, PMS-like symptoms, puffy eyes with dark circles and postnasal drip. Food allergies can cause asthma when they affect the respiratory tract; headache and insomnia when they affect the nervous system; and indigestion, irritable colon, constipation or diarrhea when they affect the GI tract. A variety of laboratory tests to detect IgE and IgG subclasses responsive to particular antigens have been developed. These have varying degrees of reliability.

Botanical Support of the Immune System

A variety of botanical preparations have been reported as useful nutritional adjuncts for protocols designed to help normalize immune processes:

Juglans nigra. Originating in Asia, *Juglans nigra* has a history of use in Greek and Roman traditions. It is said to be bitter, pungent, warm with cooling potential and is used with intestinal Qi stagnation.⁽²⁾ Black walnut leaf supports the immune system and normal tissue repair processes.

Olea europaea. Buds and leaves of olive trees contain oleuropein, a phenolic glycoside, and rutin, luteolin and hesperidin - flavonoid glycosides.⁽³⁾ The latter flavonoids possess well-defined antioxidant properties and affect aspects of the vascular system as well as the immune system. Oleuropein is the bitter principle of green olives. It can effectively prevent the oxidation of LDL *in vitro*.⁽⁴⁾ Antimicrobial properties of oleuropein have been reported, perhaps explaining the resistance of olive tree to microbial overgrowth.⁽⁵⁾



(800) 231-5777

6801 Biotics Research Drive • Rosenberg, TX 77471
biotics@bioticsresearch.com • www.bioticsresearch.com

Hydrangea arborescens. Contains thunberginols and hydrangenols which have been extensively studied by Japanese investigators. This family of compounds was shown to inhibit the release of histamine from rat mast cells and histamine-induced contraction of isolated guinea pig trachea.⁽⁶⁾ Thunberginol A administered to rats and mice normalized various aspects of immune function.⁽⁷⁾

Hesperidin. The rutinoid of hesperitin. Hesperidin is the predominant flavonoid in lemons and sweet oranges. In orange, lemon, and grapefruit skins, hesperidin and hesperitin are mainly responsible for the antioxidant activity of methanolic extracts.⁽⁸⁾ It seems also to have a membrane-stabilizing effect.⁽⁹⁾ Effects in supporting the immune system have been reported.⁽¹⁰⁾

Andrographis paniculata (Kalmegh). Folk traditions in India use this herb to support functions of the gastrointestinal tract, reduce flatulence, improve appetite and normalize liver function. Pretreatment of rats with extracts of this plant prevented carbontetrachloride induced liver damage.⁽¹¹⁾ Leaf extracts and isolated andrographolide inhibited lipid peroxidation due to exposure to carbon tetrachloride.⁽¹²⁾

Coriolus versicolor. Polysaccharide extracts of this mushroom are composed mainly of beta 1,4 glucan, together with some protein. This material has been in use in Japan, where it has been reported to enhance normal functioning of the immune system. For example, it improved T cell response in gut associated lymphoid tissue in mice.⁽¹³⁾ In rats, this extract increased B cells. The production of T cells increased in spleen and Peyer's patches, but not thymus.⁽¹⁴⁾ These mushroom polysaccharide extracts are associated with superoxide dismutase activity.⁽¹⁵⁾

Picrorrhiza kurroa. Roots and rhizomes contain a mixture of flavonoids, including glycosides, piroside and kutkoside, picrorrhizin, and the phenolic ether, apocynin. Apocynin was shown to inhibit the neutrophil oxidative burst *in vitro*.⁽¹⁶⁾ Apocynin inhibited the formation of thromboxane A2, while stimulating the release of prostaglandins by isolated guinea pig pulmonary macrophages.⁽¹⁷⁾ Picrorrhiza abstracts were found to support normal of T and B cell function using rodent model systems.⁽¹⁸⁾

Ligusticum porteri. The Ligusticum family contains a number of compounds with aromatic character. Osha has long been used in folk traditions in Southwestern United States and Northern and Central Mexico.^(19,20)

Other Nutritional Support

Dimethylglycine. Methyl donor. When lymphocytes were isolated from rabbits fed dimethylglycine and inoculated with viral or bacterial vaccines, they revealed higher proliferation rates than obtained from control animals. In addition, lymphocytes from immunized animals fed dimethylglycine showed a increased rate of DNA synthesis than controls.⁽²¹⁾

Calcium borogluconate. Boron has a role in calcium and magnesium metabolism, and possibly vitamin D.⁽²²⁾

Boronate - containing polymer stimulated T cells *in vitro*.⁽²³⁾

References

1. Kjeldsen-Kragh J et al. Controlled trial of fasting and one year vegetarian diet in rheumatoid arthritis. *Lancet* 1991; 338: 899- 902.
2. Holmes, Peter. *The Energetics of Western Herbs. Vol. I*, Artemis press, 1989; pp 415-417.
3. Ficcaro P et al. HPLC analysis of oleuropein and some flavonoids in leaf and bud of Olea Europaea L. *II Farmaco* 1991; 46:803-815.
4. Visioli R, Galli C. Oleuropein protects low density lipoprotein from oxidation. *Life Sciences* 1994; 55: 1965-1971.
5. Tassou CC, Nychas CJL. Inhibition of Staphylococcus aureus by olive phenolics in broth and in a model food system. *J Food Protection* 1994; 57: 120-124, 132.
6. Yamahara J, Murakami N. Development of bioactive functions in hydrangea ducki folium V. On the anti-allergenic and antimicrobial principles of hydrangea. *Chem Pharam Bulletin* 1996; 44: 1440-1447.
7. Yamahara J et al. Effects of thunberginol A continued in Hydrangea dulcis forium on type I-IV allergies. *Nippon Jajurigaku Zasshi* 1995; 105: 365-379.
8. Kroyer G. Uber dei antioxidative Aktivitatat von Zitrusfurchschalen (The antioxidant activity of citrus fruit peels). *Z Ernahrungswiss* 1986; 25: 63-69.
9. Santus R et al. Daflon as a cellular antioxidant and a membrane-stabilizing agent in human fibroblasts irradiated by ultraviolet A radiation. *Photodermatol hotoimmunol Photo Med* 1991; 8: 200-205.
10. Galati EM et al. Biological effects of hesperidin, a citrus flavonoid: anti-inflammatory and analgesic activity. *Farmaco* 1994; 40: 709-712.
11. Handa SS and Sharma A. Evidence of hepatoprotective activity of Andrographis panurata needs due to the diterpene lactone and andrographolide. *Research and Development of Indigenous Drugs* PC Dandiy and SB Vohora, eds. Institute of History of Medicine and Medical Research, New Delhi, 1989; 147-152.
12. Handa SS, Sharma A. Hepatoprotective activity of andrographolide against galactoseamine and paracetamol intoxication in rats. *Indian J Med Res* 1990; 92: 284-292.
13. Harada M et al. Oral administration of PSK can improve impaired antitumor CO4+ t cell responses in gut-associated lymphoid tissue of pathogen-free mice. *Int J Cancer*, 1997; 70: 362-372.
14. Nakasaki H et al. Effect of the proteoglycan (PCK) on lymphocyte subsets in normal rats. *Tokai J Exp Clin Med*, 1995; 20: 37-44.
15. Lin F et al. Free radical scavenger activities of mushroom polysaccharide extracts. *Life Sci* 1997; 60: 763-771.
16. Simons J et al. Metabolic activation of natural phenols into selective oxidative burst agonists by activated human neutrophils. *ree Radio Biol Med* 1990; 8: 251-258
17. Engels R et al. Effects of apocynin on arachidonic acid metabolism. *FEBS Letters* 1992; 305: 254-256.
18. Sharma ML et al. Immunostimulatory activity of Picrorrhiza kurroa leaf extract. *J Ethnopharmacol* 1994; 41: 185-192.
19. Appelt GD. Pharmacologic aspects of selected herbs employed in Hispanic folk medicine in the San Luis Valley of Colorado. *Ligusticum porteri* (osha) and *Matricaria Chamomilla* (Manzanilla). *J Ethnopharmacol* 1985; 13: 51-55.
20. Lihares E, Bye RA Jr. A study of four medicinal plant complexes of Mexico and adjacent United States. *J. Ethnopharmacol* 1987; 19: 153-183.
21. Reap EA, Lawson JW. Stimulation of the immune response by dimethylglycine, a nontoxic metabolite. *J Lab Clin Med* 1990; 115: 481-486.
22. Meacham SL et al. Effect of boron supplementation on blood and urinary calcium, magnesium and phosphorus, and urinary boron in athletic and sedentary women. *Am J Clin Nutr* 1995; 61: 341-345.
23. Miyazaki H et al. Boronate-containing polymer as a novel mitogen for lymphocytes. *Biochem Biophys Res Commun* 1993;195: 829-836. October 31, 1997.

To place your order for **HistoPlex®** or for additional information please contact us below.



(800) 231-5777

6801 Biotics Research Drive • Rosenberg, TX 77471
biotics@bioticsresearch.com • www.bioticsresearch.com