| ACCEPTED VIA NOTIFICATION LABEL NOT REVIEWED

Classified for "RESTRICTED USE" in New York State under 6NYCRR Part 326

SEPT 17 2020

New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration Doc ID: 569404

FLUMIOXAZIN GROUP

14

HERBICIDE

SCHOONER[™]
51.0% WDG

Non-Crop Herbicide.

For The Management of Undesirable Aquatic Vegetation in Slow Moving or Quiescent Waters

ACTIVE INGREDIENT: (% by weight)
Flumioszin' 510%
OTHER INGREDIENTS: 48.0%

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.) See inside label booklet for Precautionary Statements and Directions for Use.

FIRST AID - If on skin or clothing: *Take off contaminated clothing. *Rinse skin immediately with planty of water for 15 - 20 minutes. *Call a poison control center or doctor for restance advice. If Inflated: *Nove person for fest are 'I here says not breathing, call 91 or an ambulance, then give a striftical respiration, preferably mouth-to-mouth, if possible. *Call a poison control center or doctor for further treatment advice. If in eyes: *Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. *Remove contact lenses, if present, after the first 5 minutes, then continue missing eye. *Call a poison control center or doctor for treatment advice. If swallowed* Call a poison control center or doctor immediately for treatment advice. A twee person sip a glass of water if able to swallow. *Do not induce vomiting unless told to do so by the poison control center or doctor. *Do not give anything by mouth to an unconscious person. HOT LINE NUMBER* — Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also control statement information.

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

EPA Reg. No.: 91234-129
EPA Est. No.: 11773-IA-1(V); 62171-MS-3(OP); 39578-TX-001(M)
First letter (s) in lot number correspond to letter (s) following the EPA Est. No.
910977902

Net Weight: 5 Pounds



Manufactured for:
Atticus, LLC
5000 CentreGreen Way, Suite 100
Cary, NC 27513

Job 164913

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CALITION

Harmful if absorbed through the skin or inhaled. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Avoid breathing dust and spray mist.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below

Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride
- · Shoes and socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside
 of gloves before removing. As soon as possible, wash thoroughly and
 change into clean clothing.

ENVIRONMENTAL HAZARDS

SCHOONER 51.0% WDG is toxic to non-target plants and aquatic invertebrates. Do not apply to water except as specified on the label. Drift and runary be hazardous to non-target plants and aquatic organisms in neighboring areas, if not used in accordance with the label directions. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This pesticide is toxic to plants and must be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix or allow coming in contact with an oxidizing agent. Hazardous chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAU-Tionary Statements and directions, and with applicable State and Federal Regulations.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

RESISTANCE MANAGEMENT

SCHOONER 51.0% WDG is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to SCHOONER 51.0% WDG and other Group 14 herbicides. Weed species with acquired resistance to Group 14 herbicides may eventually dominate the weed population if Group 14 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by SCHOONER 51.0% WDG or other Group 14 herbicides

To delay herbicide resistance:

- Do not use SCHOONER 51.0% WDG or other target site of action Group 14 herbicides that might have a similar target site of action, on the same weed species.
- Use tank mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Base use on a comprehensive Integrated Pest Management (IPM) program.
 Monitor treated weed populations for loss of field efficacy.
- Contact your local extension specialist, certified crop advisors and/or manufacturer for herbicide resistance management and/or integrated weed management measures for specific crops and resistant weed biotypes.

2

TANK MIXES NOTICE

Tank mixing and/or use of this product with another product that is not specifically and expressly authorized by the label shall be at the exclusive risk of user, applicator, and/or application advisor to the extent allowed by applicable law.

Read and follow the entire label of each product to be used in the tank mix with this product.

PRODUCT USE INFORMATION

This product is a fast-acting contact herbicide that controls selected submersed, emergent, and floating aquatic weeds. It is most effective when applied to young, actively growing weeds in water with a pH of less than 8.5.

This product may be applied to the following quiescent or slow-moving bodies of water:

- Bayous
- Marshes
- Canals
 Ponds (including golf course ponds)
- Drainage ditches
- Reservoirs

Lakes

Application of this product to public aquatic areas may require special approval and/or permits. Consult with local state agencies, if required.

USE PRECAUTIONS

- There is no post-application holding restriction against use of treated water for drinking or recreational purposes (e.g. swimming, fishing).
- In areas with dense weed vegetation only treat 1/2 the water body at one time and wait 10 - 14 days before treating the remaining area. Do not retreat the same section of water within 28 days of application.
- Treated water may be used for irrigation purposes on turf and landscape ornamentals as outlined in the IRRIGATION RESTRICTIONS FOLLOWING APPLICATION table.

USE RESTRICTIONS

- Do not apply to intertidal or estuarine areas.
- Do not use treated water irrigation purposes on food crops until at least five (5) days after application.
- Do not use in water utilized for crawfish farming.
- Do not re-treat the same section of water with this product more than 6 times per year.
- \bullet Do not exceed 400 ppb of this product during any one application.

ADDITIVES

When applying this product to the foliage of floating or emerged aquatic weeds, mix with an adjuvant approved for use in aquatic sites. Mix this product with a non-ionic surfactant containing at least 80% active ingredient. Follow adjuvant manufacturer's label rates. Verify mixing compatibility by a jar test before using.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND SCHOONER 51 0% WDG

Perform a jar test before mixing commercial quantities of this product, when using this product for the first time, when using new adjuvants or when a new water source is being used.

- Add 1 pt. of water to a quart jar. The water must be from the same source and have the same temperature as the water used in the spray tank mixing operation.
- Add 3 grams (approximately 1 level tsp.) of SCHOONER 51.0% WDG for the 8 oz./A rate or 4 grams (approximately 1 1/2 tsp.) for 12 oz./A rate to the jar. Gently mix until product disperses.
- 3. Add 60 mL (4 Tbsp. or 2 fl. oz.) of additive to the quart jar and gently mix.
- 4. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
- 5. An ideal tank mix combination will be uniform and free of suspended particles. If any of the following conditions are observed do not use the adjuvant:
- Layer of oil or globules on the solution surface.
- b. Flocculation: Fine particles in suspension or as a layer on the bottom of the irr
- c. Clabbering: Thickening texture (coagulated) like gelatin.

MIXING INSTRUCTIONS

- Mix with water having pH of 5 7. If pH is higher than 7, use an appropriate buffer to reduce pH to desirable range.
- Fill clean spray tank 1/2 full of desired level with water and add buffering agent if necessary.
- Add the required amount of this product to the spray tank while agitating.
- Fill spray tank to desired level with water. Ensure that this product is thoroughly mixed before making applications. Continue agitation until spray solution has been applied.
- Mix only the amount of spray solution that can be applied the day of mixing. Apply this product within 12 hours of mixing.

SPRAYER CLEANUP

If spray equipment is dedicated to application of aquatic herbicides, completely drain the spray tank and rinse the application equipment thoroughly, including the inside and outside of the tank and all in-line screens. If spray equipment will be used for purposes other than applying aquatic herbicides, it must be thoroughly cleaned following application of SCHOOMER 51.0% WDG. The following steps must be used to clean the spray equipment:

- Completely drain the spray tank and rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
- 2. Fill the tank with clean water and flush all hoses, booms, screens, and nozzles.
- 3. Top off tank with clean water.
- 4. Circulate through sprayer for 5 minutes.
- 5. Then flush all hoses, booms, screens, and nozzles for a minimum of 15 minutes.
- 6. Drain tank completely.
- 7. Remove all nozzles and screens and rinse them with clean water.

AFRIAL APPLICATION

To obtain satisfactory weed control with aerial application of **SCHOONER 51.0% WDG**, coverage must be uniform. When applied by air,

this product may not provide adequate control of some submersed weeds. Do not spray when dirft is possible or when wind velocity is more than 10 mph. Do not spray **SCHOONER** 51.096 **WDG** within 200 feet of dwellings, adjacent sensitive crops or environmentally sensitive areas. To obtain satisfactory apolication and dirft, the following directions must be observed:

Volume Pressure

Apply SCHOONER 51.0% WDG in 5 - 10 gals, of water per acre, with a maximum spray pressure of 40 PSI. Application at less than 5 gals, per acre may not provide adequate weed control. Higher gallonage applications generally provide more consistent weed control.

Nozzles and Nozzle Operation

Use nozzles that produce flat or hollow cone spray patterns. Use non-drip type nozzles such as diaphragm type nozzles to avoid unwanted discharge of spray solution. The nozzle must be directed toward the rear of the aircraft, at an angle between 0° and 15° downward. Do not place nozzles on the outer 25% of the winos or rotors.

Adjuvants

Refer to the additive section or the tank mix partners label for adjuvant recommendation.

IRRIGATION RESTRICTIONS FOLLOWING APPLICATION

Application Method	Application Rate	Average Water Depth	Turf and Landscape Ornamentals	Ornamentals Grown for Production in Greenhouse and Nursery
Surface Spray	6-12 oz per surface acre	Greater than 3 feet	None	5 days
		Less than 3 feet	12 hours	5 days
Subsurface	Less than 200 ppb	N/A	1 day	5 days
	200-300 ppb	N/A	2 days	5 days
	300-400 ppb	N/A	3 days	5 days

SPRAY DRIFT REDUCTION

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions.

Do not apply under circumstances where possible drift to unprotected persons or to food, forage or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption can occur. These precautions are not applicable for subsurface injection by closed

- Use the largest droplet size consistent with acceptable efficacy. Formation
 of very small droplets may be minimized by appropriate nozzle selection,
 by orienting nozzles away from the air stream as much as possible and by
 avoiding excessive spray boom pressure. For ground boom and aerial
 applications, use medium or coarser spray nozzles according to ASAE 572
 definition for standard nozzles or a volume mean diameter (VMD) of
 300 microns or oreater for spinning atomizer nozzles.
- Make aerial, ground, or watercraft-based surface applications when the
 wind velocity favors on-target product deposition. Apply only when the
 wind speed is less than or equal to 10 mph. For all non-aerial applications,
 wind speed must be measured adjacent to the application site on the
 upwind side, immediately prior to application.
- Do not make aerial or ground applications into areas of temperature inversions. Inversions are characterized by stable air and increasing temperatures with increasing distance above the ground. Mist or fog may indicate the presence of an inversion in humid areas. Where permissible by local regulations, the applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the oround surface.
- Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift. Avoid spraying during conditions of low humidity and/or high temperatures.

Properly maintain and calibrate all aerial, ground, and water-based application equipment.

Where states have more stringent regulations, they should be observed.

DIRECTIONS FOR USE TO CONTROL FLOATING AND EMERGED WEEDS USING SURFACE APPLICATION

SCHOONER 51.0% WDG will control weeds and algae listed in Table 1 when applied as a broadcast spray with appropriate equipment. For best results, apply SCHOONER 51.0% WDG to the foliage of actively growing weeds.

Table 1. Floating and Emerged Weeds

Table 11 Touting and Emerged Treese				
COMMON NAME	SCIENTIFIC NAME			
Alligator Weed	Alternanthera philoxeroides			
Duckweed*	Lemna spp.			
Frog's-bit	Limnobium spongia			
Water Fern	Salvinia spp.			
Water Lettuce	Pistia stratiotes			
Watermeal*	Wolffia spp.			
Water Pennywort	Hydrocotyle spp.			
Filamentous Algae	Pithophora			
Filamentous Algae	Cladophora			

^{*200} ppb water concentration rate may be required to treat duckweed and watermeal-see DIRECTIONS FOR USE TO CONTROL SUBMERSED AND FLOATING WEEDS USING SUBSURFACE APPLICATIONS section for additional application information.

SURFACE APPLICATION

SCHOONER 51.0% WDG product as a broadcast spray at 6 - 12 ounces (0.19 - 0.38 lb ai/A) of formulated product per acre plus an adjuvant approved for use in aquatics.

SCHOONER 51.0% WDG is a contact herbicide that quickly degrades in the water column so plants that do not initially come in contact with the herbicide will not be controlled. Apply SCHOONER 51.0% WDG in a minimum of 30 gallons of water per acre to all areas of the water body where weeds exist. Coverage is essential for effective control as all floating weeds need to be exposed to lethal concentrations in all parts of the water body. Any untreated escapes or re-introductions of plants that were not treated will reestablish in areas where surface weeds had previously been controlled. Make a second

application if required to provide control once the return of these weeds is first observed, but no sooner than 28 days after the last treatment.

Apply SCHOONER 51.0% WDG during early morning hours to enhance weed control. When applying to densely packed actively growing surface weeds, ensure adequate coverage. Rapid decomposition of vegetation resulting from herbicide treatment can result in loss of oxygen in water. A sudden decrease in dissolved oxygen can result in fish suffocation. If aquatic vegetation is dense, treat floating surface weeds in sections to avoid a rapid decrease in dissolved oxygen.

SCHOONER 51.096 WDG may be tank mixed with 2,4-0, diquat, glyphosate or other registered foliar applied herbicides for enhanced control of floating and emergent weeds. Consult a manufacturer's label for specific rate restrictions and weeds controlled. Always follow the most restrictive label restrictions and precautions for all products used when making an applications involving tank mixes.

APPLICATION EQUIPMENT

Apply SCHOONER 51.0% WDG with sprayers equipped with nozzles designed to deliver the desired spray pressure and spray volume. Apply by backpack or handgun sprayer, airboat, helicopter, airplane or other application equipment that will ensure thorough coverage of target plant foliage.

DIRECTIONS FOR USE TO CONTROL SUBMERSED AND FLOATING WEEDS USING SUBSURFACE APPLICATIONS

This product controls submersed and floating weeds listed in **Table 2**.

Submersed and Floating Weeds Controlled by Subsurface Application, when applied subsurface with appropriate equipment.

Table 2. Submersed and Floating Weeds Controlled by Subsurface Application

a) ourser race repriserior			
COMMON NAME	SCIENTIFIC NAME		
Coontail	Ceratophyllum demersum		
Duckweed	Lemna spp.		
Fanwort	Cabomba caroliniana		
Hydrilla	Hydrilla verticillata		
Hygrophila	Hygrophila polysperma		
Naiad, Southern	Najas guadalupensis		

(continued)

Table 2. Submersed and Floating Weeds Controlled by Subsurface Application (continued)

SCIENTIFIC NAME	
Potamogeton crispus	
Potamogeton pectinatus	
Potamogeton diversifolius	
Salvinia spp.	
Pistia stratiotes	
Wolffia spp.	
Myriophyllum spicatum	
Myriophyllum heterophyllum	

SUBSURFACE APPLICATION

Apply this product at a rate that will produce an initial concentration of 200 to 400 ppb (of active ingredient flumioxazin) in the water column.

This product is rapidly absorbed by target plants, but also breaks down quickly in water with a pH greater than 8.5. The pH of water surrounding mats of submersed vegetation can exceed 8.5 by early to mid-day, due to photosynthetic processes. Application of this product under these conditions may provide only partial weed control, and regrowth is likely. For best control, apply this product in a minimum of 30 gallons of water per acre in the early morning to actively growing weeds and early in the season before surface matting occurs. Complete coverage and sufficient contact time of submersed weeds with this product is required for optimal performance. Application of this product with subsurface trailing hoses designed to distribute the herbicide within the plant stand will provide more effective and longer-term control of submersed weeds. Use Table 3. Subsurface Application Rates to determine the amount of this product needed to achieve desired concentration at different water depths. Use higher concentrations when weed biomass is heavy and/or weeds are more mature and topped out. Any untreated plants that are left in the water column can reinfest treated areas that had previously been controlled. Make a second application to provide control once the return of these weeds is first observed, but no sooner than 28 days after the last treatment.

When applying this product to densely packed actively growing submersed weeds, a rapid decomposition of vegetation resulting from herbicide treatment can result in loss of oxygen in water. A sudden decrease in dissolved oxygen can result in fish suffocation. If aquatic vegetation is dense, treat submersed weeds in sections to avoid a rapid decrease in dissolved oxygen. This product may be tank mixed with other registered submersed applied herbicides for enhanced control of submersed and floating weeds.

APPLICATION EQUIPMENT FOR WATER COLUMN TREATMENT

To improve distribution in the water column and ensure adequate coverage, when possible apply this product with subsurface trailing hoses in order to place the herbicide under the surface and throughout the biomass of aquatic vegetation. Keep swath width to a minimum in order to maximize contact with submersed aquatic vegetation. In small shallow water bodies, surface sprays are required to apply this product. Apply by backpack or handgun sprayer or other application equipment that will ensure adequate coverage of target plant.

INFORMATION ON HYDRILLA CONTROL IN FLORIDA

Apply this product as a subsurface treatment for Hydrilla control. For best control of Hydrilla apply during the late Winter/early Spring and/or early to late Fall. Efficacy of this product will be enhanced at these timings due to lower potential biomass present and lower pH of the water. If applied to mature topped out Hydrilla, this product will cause some discoloration and loss of growing tips, but regrowth will be rapid.

Tank mixing this product with other registered herbicides, especially if Hydrilla is approaching maturity or biomass is heavy.

Table 3. Subsurface Application Rates

Do not exceed 400 ppb of this product during any one application.

Water Depth (feet)	Pounds of SCHOONER 51.0% WDG Required Per Surface Acre to Achieve Desired Water Concentration			
	200 ppb	300 ppb	400 ppb	
1	1.1	1.6	2.1	
2	2.1	3.2	4.2	
3	3.2	4.8	6.4	
4	4.2	6.4	8.5	
5	5.3	8.0	10.6	
6	6.4	9.5	12.7	
7	7.4	11.1	14.8	

Example: To achieve an initial concentration of 200 ppb of flumioxazin in a 4-foot-deep water column, apply 4.2 lbs. of this product per surface acre.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for quidance.

CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Nonrefillable containers < 50 pounds: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available.

Nonrefillable containers > 50 pounds: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling if available.

LIMITATION OF WARRANTY AND LIABILITY

IMPORTANT: READ BEFORE USE. Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Inteffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, Ltc. All such risks shall be assumed by the user or buyer. DISCLAIMER OF WARRANTIES: To the extent consistent with applicable law, ATTICUS, Ltc makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise. that extend

LIMITATIONS OF LIABILITY: To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, nendigence, strict liability or otherwise, shall not exceed the purchase price

SCHOONER 51.0% WDG is a trademark of Atticus, LLC.

beyond statements on this label.



For The Management of Undesirable Aquatic Vegetation in Slow Moving or Quiescent Waters

ACTIVE INGREDIENT:	(% by weight)
Flumioxazin*	51.0%
OTHER INGREDIENTS:	49.0%
TOTAL:	100.0%
+0.57.0 0.4.15. 1 0 4.70 15.07/4.4.1-	

2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2*H*-1,4-benzoxazin-6-yl]-4.5.6.7-tetrahydro-1H-isoindole- 1.3(2H)-dione

KEEP OUT OF REACH OF CHILDREN CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If vou do not understand the label, find someone to explain it to you in detail.)

FIRST AID - If on skin or clothing: . Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.
 Call a poison control center or doctor for treatment advice. If inhaled: • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. . Call a poison control center or doctor for further treatment advice. If in eves: • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If swallowed: Call a poison control center or doctor immediately for treatment advice.

· Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. . Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER - Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-844-685-9173 for emergency medical treatment information.

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident, Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

PRECAUTIONARY STATEMENTS - HAZARDS TO HUMANS & DOMESTIC ANIMALS - CAUTION: Harmful if absorbed through the skin or inhaled. Avoid contact with skin, eves. or clothing. Causes moderate eve irritation. Avoid breathing dust and spray mist. ENVIRONMENTAL HAZARDS: SCHOONER 51.0% WDG is toxic to non-target plants and aquatic invertebrates. Do not apply to water except as specified on the label, to areas where surface water is present or to intertidal areas below the mean high-water mark. Drift and runoff may be hazardous to non-target plants and aquatic organisms in neighboring areas, if not used in accordance with the label directions. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate. This pesticide is toxic to plants and must be used strictly in accordance with the drift and run-off precautions on this label in order to minimize off-site exposures. Under some conditions SCHOONER 51.0% WDG may have a potential to run-off to surface water or adjacent land. Where possible, use methods which reduce soil erosion, such as no-till, limited-till and contour plowing; these methods also reduce pesticide run-off. Use vegetation filter strips along rivers, creeks, streams, wetlands, or on the downhill side of fields, where run-off could occur to minimize water run-off. PHYSICAL OR CHEMICAL HAZARDS: Do not mix or allow coming in contact with an oxidizing agent. Hazardous chemical reaction may occur.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal, PESTICIDE STORAGE: Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area, PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance, CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Offer for recycling if available or discard in a sanitary landfill. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. Nonrefillable containers ≤ 50 pounds: Nonrefillable Container: Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling if available, Nonrefillable containers > 50 pounds; Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple Rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer for recycling if available.

See inside label booklet for additional Precautionary Statements and Directions

EPA Reg. No.: 91234-129

EPA Est. No.: 11773-IA-1(V); 62171-MS-3(OP); 39578-TX-001(M) First letter(s) in lot number correspond to letter(s) following the EPA Est. No.

Manufactured for: Atticus, LLC, 5000 CentreGreen Way, Suite 100, Cary, NC 27513

PF 164913

PROOF THIS PROOF IS TO BE

CHECKED FOR ACCURACY

Please review and approve Text, Spelling, Copy Placement, Size, Shape, Colors and Dieline.

Authorized signature accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.

printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors.

THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH. Dieline does not print.



Experts in Extended Text Labeling

4201 North Westport Ave. • Sioux Falls, SD 57107 Phone: (605) 978-0451 • Fax: (605) 978-0463

DATE **JOB NUMBER CUSTOMER** 3/3/20 164913 Atticus LABEL SIZE **BOOKLET SIZE** 5" x 5 5" 4.75" x 4.5" LABEL COLORS **BOOKLET OUTSIDE COLORS BOOKLET INSIDE COLORS** PLEASE NOTE: Due to color variance between BLK BLK CYAN 342 647 BLK PATTERN VARNISH: X YES Form: CS 006B - 3/29/2017 **ARTWORK IS APPROVED** REVISED PROOF NEEDED

WE CANNOT PROCESS

THIS ORDER WITHOUT AN **AUTHORIZED SIGNATURE**

Sianed. Date