

## Cold remedy

keeps it with this medicine, and, please read on the occasion of remedy by all means.

### Common cold agent 3 A

It is a medicine improving symptoms for cold.

The acetaminophen ethenzamide is alleviation of fever, analgesic effect

The methylephedrine hydrochloride chlorpheniramine maleate phosphoric acid dihydrocodeine noscapine for mucus, nasal congestion, a cough, the symptom of the sputum

The Glycyrrhiza powdered, bellflower powdered improves for swelling and the pain of the throat.

### Must not do

(a current symptom worsens when we do not protect it, and a side effect, an accident becomes easy to occur)

1. The next person, please do not take it

(1) The person who has had a hypersensitivity symptom (rash, redness, itch, edema) with this medicine.

(2) The person who we took this medicine or other cold remedies, antipyretic analgesic, and has had asthma.

2. While you take this medicine, please do not take neither next pharmaceuticals

The oral medicine which contains other cold remedies, antipyretic analgesic, a sedative, an antitussive expectorant, an antihistamine (medicine for oral medicine for rhinitis, motion sickness medicine, allergies)

3. After remedy, please do not do the control of a vehicle or the instrumentation (sleepiness may appear.)

4. Please do not drink at the remedy

5. Please do not do long-term therapy

### Caution

1. The next person, please talk with a physician or a pharmacist before remedy

(1) The person who is treated by a physician or the dentist.

(2) A pregnant woman or the person who seems to be pregnant.

(3) The person whom we are nursing.

(4) Milk, 幼, children (under 15 years old) whom or there is the suspicion who have varicella (chicken pox) or influenza.

(5) Elderly people.

(6) The person himself or a family is the person of the allergic constitution.

(7) The person who has had an allergic symptom by medicine.

(8) The person with the next.

High fever, dysuria

(9) The person who received the next diagnosis.

Dysfunction thyroid, diabetes, cardiac disease, hypertension, liver disease, renal disease, gastroduodenal ulcer, glaucoma

(10) The person who is treated for a diabetes remedy or anticoagulant.

2. In the case of next, we cancel remedy promptly, and, please talk with a physician or a pharmacist with this document

(1) When, after remedy, the next symptom appears

Site concerned	Symptom
Skin	Rash, redness, itch
Digestive organ	Nausea, vomiting, anorexia
Mental nervous system	Dizziness
Others	Dysuria

The following serious symptom may rarely occur. In that case, please receive the practice of the physician promptly.

Name of the symptom	Symptom
Shock (anaphylaxis)	The complexion becomes pale, and hands and feet become cold, and a cold sweat, suffocation appear with a nettle-rash, edema, anguish after remedy immediately.
Oculomucocutaneous syndrome (Stevens-Johnson syndrome), toxic epidermal necrolysis (Lyell's syndrome)	With high fever, the intense symptoms such as a rash, redness, the burn-like blister appear in the mucosa of systemic skin, mouth and eyes.
Liver dysfunction	Systemic tiredness, jaundice (skin and white of eyes yellow) appear.
Interstitial pneumonia	With hacking cough (cough without the sputum), shortness of breath, dyspnea, fever appear. (when symptoms and distinction of a cold may be difficult, and symptoms such as hacking cough, the fever worsen, these symptoms canceling remedy, and receiving the practice of the physician)
Asthma	

(2) When 5-6 times do not come to have good symptom even if they take it

3. When the continuation of these symptoms or the reinforcement is found because the next symptom may appear, we discontinue taking, and, please talk with a physician or a pharmacist

Constipation, a thing of mouth or the side

### Indication

Palliation of symptoms (mucus, nose jamming, sneezing, sore throat, cough, sputum, chills, fever, headache, articular pain, pain of the muscle) of a cold

### An ingredient and effects

We include the next ingredient in 7.5 g (adults daily dose).

Ingredient	In 2.5g/pack	In 7.5 g/3pack	Effects
------------	--------------	----------------	---------

dl-methylephedrine hydrochloride powder	0.2g	0.6g	Stop cough and cut a sputum.
Chlorpheniramine maleate	0.25mg	0.0075g	suppress the allergic symptoms such as the sneezing, mucus, nose jamming.
100 phosphorus acid dihydrocodeine trituration	0.8g	2.4g	stop cough and cut a sputum.
Narcotine	0.016g	0.048g	stop cough.
Acetaminophen	0.15g	0.45g	down fever and release pain.
Ethenzamide	0.25g	0.75g	Down fever and release pain.
Caffeine	0.025g	0.075g	help with the function of alleviation of fever, analgesia ingredient.
Licorice powdered	0.267g	0.8g	It relieves the pain of the muscle and relieves the gastric mucosa irritant action of alleviation of fever, the analgesia ingredient.
Bellflower powdered	0.533g	1.6g	cut a sputum.
Starch, lactose or these mixtures	Appropriate amount	Appropriate amount	A diluting agent.

#### Dosage regimen

We do a single dose as follows and take it after a meal three times a day.

We take a single dose as follows and take it after a meal three times a day.		
Age	Single dose	The number of daily remedy
Adults (15 years old or older)	2.5 g of one pack	Three times
We are younger than 15 years old 11 years old or older	Two-thirds of adults	
We are younger than 11 years old 7 years old or older	Half of adults	
We are younger than 7 years old 3 years old or older	One-third of adults	
We are younger than 3 years old 1 year old or older	Quarter of adults	
Infants younger than one year	Do not take it	

Attention> associated with the <dosage regimen

- (1) Please adhere rigidly to a dosage regimen.
- (2) When you are given to children, please give me it with the instruction supervision of the protector.
- (3) We give priority to that you make infants younger than two years receive the practice of the physician, and, please give me it only when you are unavoidable.

#### Safekeeping and attention in the handling

- (1) Please keep it in the shaded dry cool place.
- (2) Please keep it out of reach of children.
- (3) Please do not replace it to other containers. (it causes the misuse, and a quality changes.)
- (4) When you take one remainder which you divided, we describe a residual quantity and keep it, and, please take it within two days.

#### ■ Reference

Seven project 0742-71-9501 Ltd.

Production sales agency

Drugstore seven pharmacy head office 0742-71-4783

[an external container or the matter which you should describe in external encapsulation]

#### Attention

1. The next person, please do not take it.
  - (1) The person who has had a hypersensitivity symptom (rash, redness, itch, edema) with this medicine.
  - (2) The person who we took this medicine or other cold remedies, antipyretic analgesic, and has had asthma.
2. After remedy, please do not do the control of a vehicle or the instrumentation.
3. On the occasion of remedy, please read explanation documents carefully.
4. Please keep it in the shaded dry cool place.
5. Please keep it out of reach of children.

We give priority to that you make infants younger than 6.2 years receive the practice of the physician, and, please give me it only when you are unavoidable.

This contents are not certificated.

Before using, please read check original document in Japanese.

If you were under any medication, please ask to us before taking.

+81-742-71-4783 (9:00~20:00, Sat 9:00~15:00, Sunday, national holiday is off)

<https://www.seven-ph.co.jp/asking>



<https://www.facebook.com/sevenpharmacy/>



Official LINE :