

FREY'S HATCHERY LTD.

80 Northside Drive, St. Jacobs, Ontario, N0B 2N0 • Tel (519) 664-2291 • Fax (519) 664-3491 • e-mail: orders@freyshatchery.com

2024 PRICE LIST – DAY OLD TRUCK DELIVERY

Inspected and Tested Breeder Flocks

DAY OLD MINIMUM TOTAL ORDER IS 5 – minimums are required to keep poultry comfortable in transit

SMALL ORDER CHARGES: - add \$8.00 for each AGE GROUP with a quantity of less than 10
2 Age Groups: - 1) Day Old 2) Ready-to-Lay
Quantity for Pricing: - total birds of one BREED (e.g., If ordering 50 WR pullets + 50 WR cockerels, use price for 100)
Shipping Charges: - are extra and subject to HST

CHICKS	PULLETS (Female)				NON-SEXED AS HATCHED				COCKERELS (Male)			
	1-19	20-49	50-99	100	1-19	20-49	50-99	100	1-19	20-49	50-99	100
Brown Egg Layer Breeds												
RSL Red Sex-Link	4.50	4.00	3.50	3.00	3.70	3.20	2.70	2.20	3.50	3.00	2.50	2.00
BSL Black Sex-Link	4.50	4.00	3.50	3.00	3.70	3.20	2.70	2.20	3.50	3.00	2.50	2.00
CRX Columbian Rock X	4.50	4.00	3.50	3.00	3.70	3.20	2.70	2.20	3.50	3.00	2.50	2.00
RB Rhodebar	4.50	4.00	3.50	3.00	3.70	3.20	2.70	2.20	3.50	3.00	2.50	2.00
Heritage Breeds												
RIR Rhode Island Red *	4.75	4.25	3.75	3.25	4.05	3.55	3.05	2.55	3.75	3.25	2.75	2.25
BR Barred Plymouth Rock *	4.75	4.25	3.75	3.25	4.05	3.55	3.05	2.55	3.75	3.25	2.75	2.25
Sasso Dual Purpose Breeds												
IA Irona	4.75	4.25	3.75	3.25	4.05	3.55	3.05	2.55	3.75	3.25	2.75	2.25
ST Scarlet	4.75	4.25	3.75	3.25	4.05	3.55	3.05	2.55	3.75	3.25	2.75	2.25
CI Ciara	4.75	4.25	3.75	3.25	4.05	3.55	3.05	2.55	3.75	3.25	2.75	2.25
SV Silver	4.75	4.25	3.75	3.25	4.05	3.55	3.05	2.55	3.75	3.25	2.75	2.25
VY Ivory	4.75	4.25	3.75	3.25	4.05	3.55	3.05	2.55	3.75	3.25	2.75	2.25
Other Egg Layer Breeds												
LEG White Leghorn (white)	4.50	4.00	3.50	3.00	3.70	3.20	2.70	2.20	3.50	3.00	2.50	2.00
AZ Azure Blue (blue)	6.45	5.95	5.45	4.95	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
NN Noirsans (dark brown)	6.45	5.95	5.45	4.95	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
OV Olive (green)	6.45	5.95	5.45	4.95	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
LV Layer Variety (our choice of 3 breeds)	4.65	4.15	3.65	3.15	3.65	3.15	2.65	2.15	3.35	2.85	2.35	1.85
HDC Hatch Day Choice (our choice 1 breed)	4.10	3.60	3.10	2.60	3.20	2.70	2.20	1.70	2.95	2.45	1.95	1.45

*90% sexing accuracy guarantee on Barred Rock & Rhode Island Red, all other breeds 96% (on orders over 30). No sexing accuracy guaranteed for orders less than 30 chicks. DO NOT order Barred Rock, Rhode Island Red or Columbian Rock X if you absolutely cannot have any cockerels.

Slow Growth Meat Breed												
SD Frey's Special Dual Purpose	3.65	3.15	2.65	2.15	3.40	2.90	2.40	1.90	3.30	2.80	2.30	1.80
Alternative Growth Meat Breed												
RA Rustic Ranger	3.70	3.20	2.70	2.20	3.70	3.20	2.70	2.20	4.00	3.50	3.00	2.50
Fast Growth Meat Breed												
WR White Rock X (Broiler)	3.45	2.95	2.45	1.95	3.45	2.95	2.45	1.95	3.75	3.25	2.75	2.25

Day Old continued on back

ALL PRICES SUBJECT TO CHANGE WITHOUT NOTICE.

Check our website for more information – www.freyshatchery.com

DAY OLD Continued

TURKEY POULTS		Available non-sexed only		
		1-19	20-49	50 (MAX.)
LW	Large White	7.55	7.05	6.55
MW	MiniCLASSIC White	8.10	7.60	7.10
OB	Orlopp Bronze	9.15	8.65	8.15
AG	Artisan Gold	9.65	9.15	8.65

DUCKS		Available non-sexed only			
		1-19	20-49	50-99	100
MUS	White Muscovy Ducklings	5.80	5.30	4.80	4.30

PHEASANTS <small>Minimum order is 10</small>		Available non-sexed only			
		10-19	20-49	50-99	100
PH	Ringneck Pheasants *	5.80	5.30	4.80	4.30

* There is a licence from MNR for keeping pheasants in captivity

READY-TO-LAY PULLETS (18 to 20 week old)

“READY-TO-LAY” refers to pullets that are at the maturing stage (age and weight), and will usually start laying within 2 to 4 weeks if proper management and environmental conditions are adhered to.

READY-TO-LAY PULLETS	PULLETS				COCKERELS (Limited availability)	NOTE
	2-19	20-49	50-99	100		
RTL-RSL Red Sex-Link	17.10	16.60	16.10	15.60	16.10	<p>Minimum quantity total per order is 2.</p> <p>Quantity for pricing is total birds of one breed.</p> <p>Small Order Charges: Add \$8.00 for orders less than 10 of any one age group.</p> <p>You must provide your own containers.</p>
RTL-BR Barred Plymouth Rock	19.60	19.10	18.60	18.10	18.60	
RTL-CRX Columbian Rock X	19.60	19.10	18.60	18.10	18.60	
RTL-BSL Black Sex-Link	19.60	19.10	18.60	18.10	18.60	
RTL-RIR Rhode Island Red	19.60	19.10	18.60	18.10	18.60	
RTL-LEG Leghorn	19.60	19.10	18.60	18.10	18.60	
RTL-RB Rhodebar	19.60	19.10	18.60	18.10	18.60	
RTL-IA Irona	19.60	19.10	18.60	18.10	18.60	
RTL-ST Scarlet	19.60	19.10	18.60	18.10	18.60	
RTL-CI Ciara	19.60	19.10	18.60	18.10	18.60	
RTL-SV Silver	19.60	19.10	18.60	18.10	18.60	
RTL-VY Ivory	19.60	19.10	18.60	18.10	18.60	
RTL-BF Buff Orpington	23.10	22.60	22.10	21.60	22.10	
RTL-LO Lavender Orpington	23.10	22.60	22.10	21.60	22.10	
RTL-SS Speckled Sussex	23.10	22.60	22.10	21.60	22.10	
RTL-AZ Azure Blue (blue eggs)	23.10	22.60	22.10	21.60	N/A	
RTL-NN Noirans (dark brown eggs)	23.10	22.60	22.10	21.60	N/A	
RTL-OV Olive (green eggs)	23.10	22.60	22.10	21.60	N/A	

Please note that other breeds mature several weeks slower than the Red Sex-Link.

All RTL are beak treated at day old to help prevent picking and egg breakage. Beak treatment still allows the hens to “scratch” feed or range. Pullets will have been on 12 hours of light per day when you receive them. Increase ½ to 1 hour per week (in the morning) until you reach 16 hours of light per day. Total hours of light per day and proper nutrition (no scratch or “green scraps”) will bring the pullets into lay. Greens and scratch may be given at a later age, however, if birds have access to these feed sources we recommend a vitamin supplement in the water to prevent a nutritional deficiency. As well, if you plan to allow your pullets access to the outdoors, consider putting a “wormer” medication in the water (1 or 2 application days – available from your feed supplier).

New and used crates are available for purchase. **WE DO NOT RENT OR LEND CRATES.** Green crates are an exclusive colour to Frey’s Hatchery and are NOT SOLD under any circumstances. Frey’s Hatchery cannot allow any of our crates to be used for any other purpose than our Ready-to-Lay deliveries.

Proper Handling Procedures to Prevent Contamination: Please be aware that all poultry, even chicks, are livestock and should be treated as such. Proper handling procedures will reduce or eliminate the chance of bacterial contamination. A human hand can have hundreds of species and millions of individual bacteria on it, as can poultry. **Wash your hands with soap and water after handling poultry or any equipment that comes in contact with poultry.**

Frey’s Guarantee: We guarantee 100% live arrival of the quantity ordered. Extras may be included in your order. Orders are accepted subject to conditions incidental to the hatchery business. Our responsibility for loss or damage arising out of incorrect shipment of breeds, sex or numbers, delay or failure of shipment or any other cause is limited to the original price paid. Frey’s Hatchery Ltd. operates under Federal and Provincial government regulations in regards to the operation of a licensed hatchery, animal husbandry practices, and disease control standards. It is understood and agreed that no representation or warranty as to livability, production, freedom from disease, fitness for use, or otherwise, is made by Frey’s Hatchery Ltd.



White Rock Weights

Approximate White Rock Live Weights						
Age in weeks	Females			Males		
	kgs	lbs	Approx feed kgs	kgs	lbs	Approx feed kgs
3	.7	1.5	1.1	.8	1.75	1.3
4	1.2	2.6	2	1.4	3	2.3
5	1.7	3.7	3	2.0	4.4	3.6
6	2.2	4.8	4.5	2.7	6	5
7	2.7	5.9	6	3.4	7.4	6.6
8	3.2	7	7.5	3.9	8.5	8.2
9				4.3	9.4	9.5

Note: Eviscerated weights are generally 75%+ of live weight

These are the typical ages that White Rocks are processed in the commercial chicken industry. If you grow them longer and larger, then extra care is needed to minimize losses. You can try to slow growth through feed restriction or other methods to get them to do something other than eating! Vitamins in the water are great any time (feed restriction, hot/cold weather, or periods of stress).

Rustic Ranger Weights

Approximate Rustic Ranger Live Weights						
Age in weeks	Females			Males		
	kgs	lbs	Approx feed kgs	kgs	lbs	Approx feed kgs
6	1.55	3.4	3.1	1.66	3.7	3.5
7	1.96	4.3	4.0	2.1	6.9	4.5
8	2.36	5.2	4.9	2.53	5.6	5.4
9	2.75	6.1	5.8	2.94	6.5	6.4
10	3.09	6.8	6.7	3.31	7.3	7.4
11	3.39	7.5	7.7	3.66	8.1	8.5
12	3.69	8.1	8.9	3.95	8.7	9.8

Note: Eviscerated weights are generally 75%+ of live weight

The heaviest colour broiler. Perfect compromise between rusticity, feed conversion rate and yield. They are a great pasture bird and excellent forager. Produces a table bird in just 12 weeks.

Special Dual Purpose

Females will be 5-6 lbs and males 7-8 lbs live weight in 15 weeks. These birds can take up to 13 kgs of feed to reach 15 weeks old.



SPECIALTY TURKEYS

Artisan Gold Turkey

The Artisan Gold is a rare breed that is smaller in size and slow growing. It has black feathers, unique colouring in the face and feet, long black legs, and a pronounced beak. It has natural tendencies such as roaming and roosting, and offers a robust, gamey taste that is exotically different from traditional turkeys.

MiniCLASSIC White Turkey

This turkey is small in stature and has been bred to develop and mature like natural bred turkeys of the past. This means that the Toms don't grow as big and can be harvested more efficiently.

Orlopp Bronze Turkey


This premium broad-breasted turkey has excellent conformation, high meat quality, natural fat layering, and beautiful feathering. Weights and feed conversion will rival traditional Large White turkeys.

Approximate Live Weights (lbs)								
Age in weeks	Artisan Gold		MiniCLASSIC		Orlopp Bronze		Large White	
	Hens	Toms	Hens	Toms	Hens	Toms	Hens	Toms
8	4.75	5.93	6.52	8.13	8.14	9.79	8.65	10.85
9	5.75	7.35	8.04	10.08	9.81	11.99	10.61	13.49
10	6.75	8.84	9.60	12.11	11.51	14.33	12.69	16.47
11	7.72	10.34	11.12	14.18	13.22	16.78	14.70	19.46
12	8.63	11.84	12.52	16.24	14.91	19.31	16.62	22.46
13	9.45	13.29	13.73	18.25	16.53	21.89	18.45	25.46
14	10.16	14.65	14.73	20.15	18.05	24.50	20.12	28.45
15	10.73	15.9	15.47	21.92	19.45	27.10	21.66	31.43
16	11.15	16.99	15.99	23.49	20.67	29.68	23.04	34.31
18	11.45	18.57	16.61	25.89	22.48	34.65	25.37	39.82
20					23.22	39.18	27.28	44.91

Note: Eviscerated weights are generally 75-80% of live weight

Please note that the figures above are based on trials recorded but your results may differ depending on individual circumstances.

For more information see – www.freyshatchery.com



Day-old Chicks: Transport in the boxes at 75°F/24°C and DO NOT put them in direct sunlight or cool drafts.

- Temperature 92°F/33°C at level of chicks (2 in/5 cm) from the top of the bedding.
- A thermometer is a **MUST!** A hi/lo thermometer is best.
- 3 to 4 in/8 to 10 cm of bedding (preferably shavings).
- You can put a sheet of styrofoam under the shavings.
- No drafts in the brooding area (18 in/46 cm high guard should be used to stop drafts). Keep brooding area “rounded” so chicks don't crowd into a corner and smother. Enlarge brooding area as they grow and temperatures are lowered.
- Lower temperature 4°F/2°C per week until six weeks old.
- Keep shavings dry (clean up water spills or leaks and replace with dry shavings).
- Keep water and feed very close to chicks.
- Use lukewarm water in drinkers at first and dip the chick's beak in the water when you take them out of the box.

White Rocks:

Same as above along with these extra tips

- White Rocks are a **VERY** picky breed and need their conditions to be perfect.
- They require a consistent temperature of 92F/33C degrees for their first week and then following the guidelines above for decreasing heat. We recommend raising your chicks in an inside environment where you have better control of regulating heat. A thermometer is a must to monitor this.
- Please note that our chicks go home to various environments so we cannot control your success once they leave our facility.
- More information can be found at www.freyshatchery.com under *CHICKENS, White Rock Care*.

Turkeys and Pheasants:

Same as above EXCEPT

- Temperature at their level 95-99°F/35-37°C.
- Put a cloth, towel or coarse permeable covering on top of the shavings in the brooding area and scatter feed all over it. (This will allow them to learn what feed is more easily and prevent them from slipping and causing irreversible leg damage.)
- Remove the cloth after a day or two or if it gets too damp or wet.
- Always have lots of feed trays and drinkers very close to the turkeys.

Ready-to-lays:

- Ready-to-lays can handle lower temperatures during transit but do not let them get overheated...keep air moving around them if it is a warm day.
- Pullets have been on 12 hours of light per day when you receive them. Increase ½ to 1 hour per week in the morning until you reach 16 hours of light per day. Total hours of light per day and proper nutrition (no scratch or “green scraps”) will bring the pullets into lay. Greens and scratch may be given at a later age, however, if birds have access to these feed sources we recommend a vitamin supplement in the water to prevent a nutritional deficiency.
- If you plan to allow your pullets access to the outdoors, consider putting a “wormer” medication in the water (1 or 2 application days – available from your feed supplier).

What is infectious laryngotracheitis (ILT)?

ILT is a viral respiratory disease of poultry, fowl and other birds. Clinical signs range from mild disease including discharge from the eyes and nose, sneezing and a drop in egg production to more severe signs including coughing up blood, bloody discharge from the nose and mouth, gasping for air, and death. It takes anywhere from 6 to 14 days from when the bird becomes infected to when it shows clinical signs allowing for asymptomatic spread during this time. Five to 100% of the flock may become sick and 0 to 20% of the flock may die.



Infectious Laryngotracheitis in Poultry - Poultry - Merck Veterinary Manual (merckvetmanual.com)

What causes ILT?

ILT is caused by a herpes virus which causes a lifelong infection in the bird. Herpes viruses have the ability to hide from the infected bird's immune system until times of stress when the virus "awakens" and causes disease. Stressful events which can cause this reactivation include; mixing of new birds, sudden weather changes, transportation, vaccination for other diseases and illness. The ILT virus can survive outside of the bird too, including 3 months on wood surfaces protected from light, 3 weeks in buried carcasses, 3 days in droppings, 4 days in wet litter and 20 days in dry litter.

How does ILT spread?

ILT is most commonly spread to susceptible birds through direct contact with infected birds. The virus can be spread between barns and farms via inanimate objects however, such as contaminated equipment, vehicles, clothing and footwear. Winds can also carry the virus downwind of infected barns and manure. Enhanced biosecurity is therefore critical to prevent spread.



Blood and exudate in the trachea of a bird with ILT (photo credit: Dr. M Petrik)

How is ILT diagnosed?

Your veterinarian may submit sick or dead birds for post-mortem and tissue examinations. Findings consistent with ILT will be confirmed with testing for viral DNA using PCR (polymerase chain reaction) testing. To be considered a case of ILT, 2 of the following 3 criteria must be met; clinical signs consistent with ILT, evidence of the virus on tissue examination (viral inclusion bodies), PCR positive test.

How is ILT treated and managed?

There is no treatment for ILT, however your veterinarian may prescribe antibiotics for secondary bacterial infections. Key management practices for controlling the spread of ILT within and between flocks include:

Vaccination

- If your previous flock was infected or if you are in close proximity to an infected premises, it is recommended that broiler chicks be vaccinated for ILT for two successive crops.
- If you are a producer in the FBCC biosecurity advisory area, discuss with your veterinarian about vaccinating your flock for ILT

Manure management

- Heat-treat litter at 38° C (100° F) for 100 hours prior to thorough cleanout and disinfection.
- When possible, wait for 8 weeks before manure is moved off the premises. On-site manure must be covered to reduce spread. If manure must be moved, ensure the manure hauler is aware of ILT on your farm and they employ appropriate biosecurity practices including securely covering manure in transit.

Delivery Dates	Week #	Day Old Chicks	Day Old Turkeys	Day Old Ducks	Day Old Pheasants	Ready-to-lay (18-20 wks)				
						RSL	Variety AZ/BSL/BR/ CRX/RIR	Fancy AZ/BF/LO/ NN/OV/SS	Sasso IA/ST/ VY/SV/CI	Col. Egg AZ/LEG/ NN/OV/RB
March 20	12 Wed	Yes				Yes	Yes			
April 17	16 Wed	Yes	Yes	Yes		Yes		AZ/NN only		
May 15	20 Wed	Yes	Yes	Yes	Yes	Yes				
June 12	24 Wed	Yes	Yes	Yes	Yes			Yes		
July 10	28 Wed	Yes	Yes	Yes	Yes	Yes			Yes	
Aug. 14	33 Wed	Yes	Yes			Yes				Yes
Sept. 11	37 Wed	Yes	Yes LW only			Yes	Yes			

B & BR (24)

Place order 4 weeks in advance for day olds.

Ready-to-lays are available on a first-come first-serve basis.

Please note that ready-to-lay delivery days of the week are subject to change dependent on outside air temperatures and the logistics of arranging deliveries.

Frey's Hatchery
 Email to info@cleary.shop

**** NEW ORDERS ONLY ** / Do not use for changes**

To avoid duplicate orders,
 please initial when order faxed.

For revisions (dates, breed, sex, services) and cancellations, **PLEASE** send sales order with changes indicated.

CUSTOMER: _____

DEALER NO.: A9926

DEALER: CLEARY FEED & SEED

ADDRESS: _____

Please check your delivery dates!

PHONE: (H) _____ (W) _____

DATE ORDERED: _____

1st CHOICE DATE: _____

ORDER TAKEN BY: _____

2nd CHOICE DATE: _____

Office Use Only	Customer Email	Family Food # (Form 300)	Office Use Only
-----------------	----------------	--------------------------	-----------------

DAY-OLD BIRDS *Check price list for small order charge					READY-TO-LAY PULLETS *Check price list for minimum & small order charge				
Breed	Qty.	Sex P / C / NS	Unit Pr.	Total	Breed	Qty.	Sex P / C	Unit Pr.	Total

Check if okay to substitute similar breed(s) if first choice breed not available or list alternate breed choices in NOTES section below

MISCELLANEOUS CHARGES (Used crates, eggs, services, etc.)				
Description	Qty.	Price	H.S.T.	Total

NOTES:

Bird Cost (No HST) _____
 Small Order Charge(s) _____
 Other Charges _____
 Delivery + HST _____
 (Less Deposit) _____
TOTAL OWING _____