

DC Brunnenstraße 61 53347 Alfter Germany

Phone: +49 - (0)2222 / 9108 - 10 +49 - (0)2222 / 9108 - 40 Fax: E-mail: info@dermaconsult.com Web: www.dermaconsult.com

23.06.2020 Date:

Expertise

Examination of the Product

"BIOVOLEN® Harzsalbe"

F-505 Charge: HT-0520-245-09 (cosmetic)

Concentration: undiluted

by Human Patch Test (Cosmetic Trial)

Sponsor

Evertz Pharma GmbH

Speicherstr. 20 60327 Frankfurt a.M. Germany

Performing Laboratory

Derma Consult GmbH

Brunnenstr. 61 53347 Alfter Germany

Study Details

Type of study....: Determination of irritating effects to the skin with an occlusive patch test.

Study Period: June 2020

Study Director: Dr. med. H. Prieur

Test subjects: 50 (18-60 years; sex distribution non-standardized)

30 normal healthy, 3 eczema, 1 allergy and 16 subjects with sensitive skin

Test site..... Back

Concentration....: Undiluted

Controls...... SDS (1% in water), water

Summary Results

All participants completed the study. Under the test conditions, SDS (1% in water) caused positive reactions in 12 subjects. The negative control water showed no reactions. None of the subjects showed any reaction to the test product. On the basis of the test results and under the test conditions, the product

"BIOVOLEN® Harzsalbe"

is to be classified as 'harmless' as regards the possibility of skin irritation.

Signature:

Dr. med. H. Prieur Dermatologist - Allergist

Signature:

Dr. J. Nissen Pharmacist - M.D.R.A.

Page 1 of 3

Dr. Hans-Peter Nissen Dr. Jana Nissen VAT-REG.No. DE 123309725

district court Bonn HRB 5272 Tax No. 222/5703/0374

manager:

Methodology

Introduction

The epicutaneous patch test allows us to assess the primary skin irritation potential of cosmetic-finished products and raw materials.

Description

All the work described in this expertise was conducted considering the guidelines by COLIPA (Walker A.P. et al: Test Guidelines for Assessment of Skin Compatibility of Cosmetic Finished Products in Man. Food and Chemical Toxicology 34, 1996, 651-660; COSMETICS EUROPE: Product test guidelines for the assessment of human skin compatibility 1997).

Because it was a study with humans, it was carried out taking into account the principle requirements of the Declaration of Helsinki (1964) and subsequent revisions.

Experiments were carried out on 50 volunteers (30 normal healthy subjects, 3 eczema patients, 1 allergy patients, 16 subjects with sensitive skin) between the ages of 18 to 60. Sex distribution was not standardized. The volunteers were clearly informed, verbally and in writing, regarding the nature of the study, the timetable, constraints and possible risks. They gave their written informed consent before participating in the study.

Participants could withdraw from the study at any time without giving reason. During the test period, the subjects refrained from using other substances on the test areas.

Inclusion criteria

- informed volunteers
- age \geq 18 years

Exclusion criteria

- pregnant or lactating women
- blemishes or marks (tattoos, sunburn) which interfere with scoring
- any skin disease that may interfere with the aim of the study

Procedure

The product was applied in a concentration as outlined above in square test-chambers (allergEAZE® clear Patch Test Chambers; SmartPractice®, Phoenix, AZ) to the backs of the panellists for a period of 48 hours. Proper adherence of the test patches was assured by the inclusion of sodium dodecyl sulphate (SDS) in one concentration (1%) as positive control. Water was used as a negative control.

The treatment sites were assessed for the presence of irritation by a trained evaluator using a 5 point visual scoring scale at 48 h (30 min after patch removal) and 72 h after patch application.

Scoring scale

Erythema 0: no E., 1: slight E., 2: significant E., 3: pronounced E., 4: strong E.

Fissure 0: no F., 1: minimal F., 2: significantly perceptible F., 3: pronounced F., 4: ulceration

Scaling0: no Sc., 1: minimal Sc., 2: moderate Sc., 3: significant Sc., 4: closed scale crust

Results

The test results outlining the data for erythema, scaling and fissure formation on a per subject base for the test product are attached in tabulated form.

Literature

Magnus Lindberg and Mihaly Matura:

"Patch Testing" in

J. D. Johansen, P.J. Frosch, J.-P. Lepoittevin (Eds.),

Contact Dermatitis 5th Edition

Springer-Verlag, Berlin Heidelberg, Germany (2011), pp. 439-464

J-M. Lachapelle, H. I. Maibach:

"Patch Testing and Prick Testing - A practical Guide

Official Publication of the ICDRG"

Third Edition

Springer-Verlag, Berlin Heidelberg, Germany (2012)

Appendix: test protocol

Product: BIOVOLEN® Harzsalbe

F-505 Charge:, HT-0520-245-09 (cosmetic)

| No. | Туре | after 48 h | | | after 72 h | | | |
|-----|--|------------|-----|-----|------------|-----|-----|--|
| | | E | F | S | Е | F | S | |
| 1 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 2 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 3 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 4 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 5 | S | Ō | 0 | Ō | 0 | Ō | 0 | |
| 6 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 7 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 8 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 9 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 10 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 11 | ľΕ | 0 | 0 | 0 | 0 | 0 | 0 | |
| 12 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 13 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 14 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 15 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 16 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 17 | s | 0 | 0 | 0 | 0 | 0 | 0 | |
| 18 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 19 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 20 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 21 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 22 | | 0 | 0 | 0 | 0 | 0 | ō | |
| 23 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 24 | E | 0 | 0 | 0 | 0 | 0 | 0 | |
| 25 | + | 0 | 0 | 0 | 0 | 0 | 0 | |
| 26 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 27 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 28 | E | 0 | 0 | 0 | 0 | 0 | 0 | |
| 29 | - - | 0 | 0 | 0 | 0 | 0 | 0 | |
| 30 | 1 | 0 | 0 | 0 | 0 | 0 | ō | |
| 31 | 1 | 0 | 0 | 0 | 0 | 0 | ō | |
| 32 | A | 0 | 0 | 0 | 0 | 0 | 0 | |
| 33 | ^ | 0 | 0 | 0 | 0 | 0 | 0 | |
| 34 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 35 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 36 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 37 | 1 3 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 38 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 39 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 40 | - | 0 | 0 | 0 | 0 | 0 | 0 | |
| 41 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 42 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 43 | 1 3 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 44 | - | 0 | 0 | 0 | 0 | 0 | 0 | |
| 45 | S | 0 | 0 | 0 | 0 | 0 | 0 | |
| 46 | + 3 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 47 | S | | | 0 | 0 | 0 | 0 | |
| 48 | + 3 | 0 | 0 | 0 | 0 | 0 | 0 | |
| 48 | | 0 | 0 | 0 | 0 | 0 | 0 | |
| 50 | | 0 | 0 | | 0 | 0 | 0 | |
| | 4 | | | 0 | | | _ | |
| SUM | | 0,0 | 0,0 | 0,0 | 0,0 | 0,0 | 0,0 | |

Erythema (E): no E.: 0, slight E.: 1, clear E.: 2, severe E.: 3, very severe E.: 4

Fissures (F): no F.: 0, minimal F.: 1, clearly visible F.: 2, distinct F.: 3, ulceration: 4

Scales (S): no S.: 0, minimal S.: 1, clearly visible S.: 2, moderate S.: 3, distinct S.: 4

S: subjects with sensitive skin

E: patients with eczema

A: patients with allergy

| No. | Туре | after 48 h | | | after 72 h | | |
|----------|----------------------------|------------|-----|-----|------------|-----|-----|
| . As | , , | Е | F | S | Е | F | S |
| 1 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 2 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 3 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 4 | THE REPORT OF THE PARTY OF | 0 | 0 | 0 | 0 | 0 | 0 |
| 5 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 6 | - | 0 | 0 | 0 | 0 | 0 | 0 |
| 7 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 8 | - ŭ | 0 | 0 | 0 | 0 | 0 | 0 |
| 9 | | 0 | 0 | 0 | 0 | 0 | ō |
| 10 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 11 | E | 0 | 0 | 0 | 0 | 0 | 0 |
| 12 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 13 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 14 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 15 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 16 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 17 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 18 | 3 | 0 | 0 | 0 | 0 | 0 | 0 |
| 19 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 20 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 21 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 22 | 3 | 0 | 0 | 0 | 0 | 0 | 0 |
| | | | | | | | |
| 23 | - | 0 | 0 | 0 | 0 | 0 | 0 |
| 24 25 | E | 0 | 0 | 0 | 0 | 0 | 0 |
| | | 0 | | 0 | 0 | 0 | 0 |
| 26 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 27 | _ | 0 | 0 | 0 | 0 | 0 | 0 |
| 28 | E | 0 | 0 | 0 | 0 | 0 | 0 |
| 29 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 30 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 31 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 32 | Α | 0 | 0 | 0 | 0 | 0 | 0 |
| 33 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 34 | _ | 0 | 0 | 0 | 0 | 0 | 0 |
| 35 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 36 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 37 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 38 | | 0 | 0 | 0 | 0 | 0 0 | 0 |
| 39 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 41 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 42 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 43 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 44 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 45 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 46 | - | 0 | 0 | 0 | 0 | 0 | 0 |
| 47 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 48 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 49 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 50 | | 0 | 0 | 0 | 0 | 0 | 0 |
| SUN | VI | 0,0 | 0,0 | 0,0 | 0,0 | 0,0 | 0,0 |

Erythema (E): no E.: 0, slight E.: 1, clear E.: 2, severe E.: 3, very severe E.: 4
Fissures (F): no F.: 0, minimal F.: 1, clearly visible F.: 2, distinct F.: 3, ulceration: 4
Scales (S): no S.: 0, minimal S.: 1, clearly visible S.: 2, moderate S.: 3, distinct S.: 4

S: subjects with sensitive skin E: patients with eczema A: patients with allergy

| No. | Туре | after 48 h | | | after 72 h | | |
|-----|-------|------------|-----|------|------------|-----|------|
| , | .,,,- | E | F | S | Е | F | S |
| 1 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 2 | S | 1 | 0 | 1 | 2 | 0 | 1 |
| 3 | S | 1 | 0 | Ö | 0 | 0 | 0 |
| 4 | | 0 | 0 | 0 | ō | 0 | 0 |
| 5 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 6 | | 0 | 0 | 0 | Ö | 0 | 0 |
| 7 | S | 1 | 0 | 0 | 0 | 0 | 0 |
| 8 | | Ö | ō | 0 | 0 | 0 | ō |
| 9 | | ō | 0 | 0 | 0 | 0 | 0 |
| 10 | | ō | 0 | Ō | 0 | 0 | 0 |
| 11 | E | 0 | ō | 0 | Ö | 0 | 0 |
| 12 | | 0 | Ō | 0 | 0 | 0 | 0 |
| 13 | S | 0 | 0 | 0 | ō | 0 | 0 |
| 14 | S | 1 | Ō | 0 | 0 | 0 | 0 |
| 15 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 16 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 17 | S | 1 | 0 | 0 | 1 | 0 | 0 |
| 18 | -ŭ- | 0 | 0 | 0 | 0 | 0 | 0 |
| 19 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 20 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 21 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 22 | Ů | 0 | 0 | 0 | 0 | 0 | 0 |
| 23 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 24 | E | 2 | 0 | 2 | 2 | 0 | 2 |
| 25 | - | 0 | 0 | 0 | 0 | 0 | 0 |
| 26 | - | 0 | 0 | 0 | 0 | 0 | 0 |
| 27 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 28 | E | 0 | 0 | 0 | 0 | 0 | 0 |
| 29 | | 1 | 0 | 0 | 0 | 0 | 0 |
| 30 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 31 | - | 0 | 0 | 0 | 0 | 0 | 0 |
| 32 | _ | 1 | 0 | 0 | 1 | 0 | 0 |
| 33 | A | 0 | 0 | 0 | 0 | 0 | 0 |
| 34 | - | 0 | 0 | 0 | 0 | 0 | 0 |
| 35 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 36 | S | 1 | 0 | 0 | 1 | 0 | 0 |
| 37 | 3 | 0 | 0 | 0 | 0 | 0 | 0 |
| | | _ | _ | 0 | 0 | 0 | 0 |
| 38 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 41 | S | 1 | 0 | 0 | 0 | 0 | 0 |
| 42 | S | 1 | 0 | 1 | 1 | 0 | 1 |
| 43 | - | 0 | 0 | 0 | 0 | 0 | 0 |
| 43 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 45 | S | 0 | 0 | 0 | 0 | 0 | 0 |
| 46 | 3 | 0 | 0 | 0 | 0 | 0 | 0 |
| 47 | S | 1 | 0 | 0 | 0 | 0 | 0 |
| 48 | - | 0 | 0 | 0 | 0 | 0 | 0 |
| 49 | | 0 | 0 | 0 | 0 | 0 | 0 |
| 50 | | 0 | 0 | 0 | 0 | 0 | 0 |
| | | | | == | | | |
| SUM | | 0,26 | 0,0 | 0,08 | 0,16 | 0,0 | 0,08 |

Erythema (E): no E.: 0, slight E.: 1, clear E.: 2, severe E.: 3, very severe E.: 4
Fissures (F): no F.: 0, minimal F.: 1, clearly visible F.: 2, distinct F.: 3, ulceration: 4
Scales (S): no S.: 0, minimal S.: 1, clearly visible S.: 2, moderate S.: 3, distinct S.: 4

S: subjects with sensitive skin

E: patients with eczema A: patients with allergy