

AUSTRALIAN MANUKA FACT SHEET



KEY POINTS:

Manuka honey is produced by bees from the flowers (nectar) of *Leptospermum* species plants, **native to Australia and New Zealand.**



Australia is home to over 80 species of *Leptospermum*, while New Zealand has only one species (*Leptospermum scoparium*), which is believed to have originated from Tasmania, Australia.

Australia has had **a long history of producing Manuka honey**, since beekeeping was introduced.

The special properties of Manuka honey with **exceptionally high levels of activity** were first identified by Professor Peter Molan in the 1980s while he was studying *L. scoparium* honey from New Zealand, and we have since discovered many sources of Australian Manuka.

Manuka honey is **one of the most famous honeys in the world** because of its medicinal properties, including:

- Antimicrobial activity – that is, the ability to kill superbugs (antibiotic-resistant bacteria and other difficult to treat microbes) that cause serious infections
- Wound healing and anti-inflammatory activity.

THE SCIENCE BEHIND THE MEDICINAL PROPERTIES OF AUSTRALIAN MANUKA HONEY:

Humans have used honey for its healing properties for thousands of years, and now with on-going research scientists are able to explain the reasons behind its medicinal power. We are also starting to understand that some honeys are hundreds of times more powerful than others.

The medicinal power of honey all comes down to which plant flowers bees visit to collect the nectar that they turn into honey.

Manuka honey (from *Leptospermum* species plants) is one of the most famous honeys in


the world because there have been numerous scientific studies showing that this honey kills superbugs and stimulates wound healing.

The special properties of Manuka honey were first found in New Zealand in the 1980s (Molan & Russell, 1988) and since then we have discovered many sources of active Manuka honey in Australia stemming from the Leptospermum species (Cokcetin et al. 2016). This isn't surprising as Australia is home to the largest diversity of Leptospermum plants in the world, and we have

more than 80 species compared to New Zealand's one! Our native Leptospermum plants are found throughout Australia, and grow in some of the most pristine natural environments in the world.



WHAT MAKES MANUKA HONEY SO SPECIAL?



A natural compound called **methylglyoxal (MGO)** is responsible for much of the unique activity of Manuka honey. The MGO comes from a component, called dihydroxyacetone (DHA), which occurs organically in the nectar of flowers of some Leptospermum plants.

Various species of the Manuka plant produce different levels of DHA that give rise to the varying antibacterial potencies of this special honey. Scientists looking at Australian honeys

have found a number of sources of Manuka with exceptionally high levels of activity. (Cokcetin et al. 2016; Windsor et al. 2012).

One of the most amazing things about the antibacterial activity of Manuka honey is that although bacteria that cause serious infections can develop resistance to our modern antibiotics (which is how they become superbugs), they cannot develop resistance to the activity of Manuka honey (Blair et al. 2009).

SOME MORE FACTS ABOUT AUSTRALIA'S MANUKA HONEY:

- Manuka honey is produced from the Leptospermum species of plants.  Leptospermum scoparium is one of more than 80 species found in Australia, but the only one found in New Zealand, with the NZ species originating in Australia (Fleming 1975, Thompson 1989, Van Eaton 2014).

- The New Zealand Maori have at least six terms for NZ's Leptospermum scoparium, not just the word Manuka (Van Eaton, 2014).
- Manuka honey was not produced in Australia or New Zealand until the European honey bee (*Apis mellifera*) was introduced. This happened in 1822 for the Australian mainland, 1831 for Tasmania

(Watson), and in 1839 for New Zealand (Mary Bumby).

- Australia has been using the word Manuka since European settlement for naming places, property and of course the plant. In 1880, there is documented evidence of the term Manuka being used in reference to the plant, it is included in maps held in the Tasmanian archives, and there have been many other references in newspapers, titles and other documents since then.
- Australia also has a long history of calling Leptospermum honey Manuka. For example, in 1935 in the South Australian Chronicle there is mention of the production of Manuka honey, including it being described as “the repulsive honey”

by a beekeeper from Border Town, and discussing how to avoid ending up with Manuka honey in hives, its strong flavour and the effects it would have on the flavour of other Australian premium honey, such as eucalypt.

- Sometimes the term ‘tea tree’ and Australian Manuka are interchanged as a consequence of the similar appearance of the plants, but tea trees (famous for their oils) are from the Melaleuca genus within the Myrtaceae family, while Manuka is from the Leptospermum genus within this family.
- The suggestion that only honey from New Zealand can be described as Manuka is clearly ridiculous, and strongly refuted by the Australian honey industry.

ANTIBACTERIAL POTENCY TESTING OF MANUKA HONEY EXPLAINED:

For commercially competitive reasons, there have been a number of symbols introduced into the market aiming to represent the antibacterial strength of active honey. These include **NPA** (Non-Peroxide Activity), **ULF** (Unique Leptospermum Factor) and **MGO** (methylglyoxal).

The antibacterial strength can be measured and is often shown as a number on the labels (i.e. **NPA 10+**; **ULF 10+**; **MGO 263+**). The higher the activity rating the higher the antibacterial properties and strength of the honey.

There is currently no widely accepted gold standard for measuring the activity of honey, making deciding on a particular honey complex for consumers and the marketplace. However, we will explain the varying techniques:

1. **Non-peroxide activity (NPA):** In 1981, Professor Peter Molan (MBE), University of Waikato in New Zealand, was researching the hydrogen peroxide activity in honeys from around the world when he found that when he deliberately destroyed this activity only one honey type still showed strong antibacterial activity. This activity became known as “non-peroxide activity” (NPA). Professor Molan went on to research this activity over many years and found that it was very stable and unique to Manuka honey. This Manuka NPA could withstand both heat and light exposure and still remain highly active and effective.



Further research by Professor Molan, his team, and others has revealed that Manuka honey is very effective against many strains of bacteria, including deadly strains of *Staphylococcus aureus*, such as the superbug MRSA (a.k.a. 'golden staph'). Professor Molan's team also found that Manuka honeys had different levels of NPA, so a test was developed to measure the strength of this activity in each batch. The rating for NPA is measured by comparing the activity of the honey to a known antimicrobial, phenol. So, for example, a NPA 5+ Manuka honey has the same non-peroxide antibacterial activity as 5% phenol.

Note – Since Professor Molan's pioneering work, Manuka honey containing the unique non-peroxide activity (NPA) has been found to occur naturally in both New Zealand and Australia


2. ULF (Unique Leptospermum Factor) measures Non Peroxide Activity (NPA). ULF, or Unique Leptospermum Factor, serves as a pivotal marker in assessing the distinctive fingerprint of Manuka Honey, particularly in terms of its Non Peroxide Activity (NPA). This specialized measurement system, unique to Manuka Honey, is designed to quantify the bioactive components that contribute to its renowned therapeutic properties. ULF essentially gauges the presence and potency of non-peroxide compounds, such as methylglyoxal, which are responsible for the honey's exceptional antibacterial and anti-inflammatory qualities. As an indicator of the specific attributes that set Manuka Honey apart from other honey varieties, ULF plays a crucial role in providing consumers with a standardized metric to identify and appreciate the genuine, high-quality nature of this extraordinary honey.

Note – NPA and ULF antibacterial activity ratings are measured the same way and are equivalent to one another.

3. Methylglyoxal (MGO) content: In 2008, two laboratories showed that one of the major components that is responsible for the unique activity in Manuka honey is the presence of the naturally occurring compound methylglyoxal (MGO) (Adams et al. 2008; Mavric et al. 2008). The MGO is produced from a conversion of dihydroxyacetone that is present in nectar from the Manuka flower (Adams et al. 2009). The quantity of MGO in Manuka honey is now commonly used as an indicator of the strength of the antibacterial activity, as it correlates strongly with the NPA. MGO can be accurately measured in parts per million (ppm).

The following are estimates of the equivalence between NPA and MGO measurements:

NPA 5+ ~ MGO 100 ppm
NPA 10+ ~ MGO 260 ppm
NPA 15+ ~ MGO 450 ppm
NPA 20+ ~ MGO 800 ppm



Graph derived from the scientific research on Australian Leptospermum honey by Cokcetin, Pappalardo et al. 2016

Note – Care should be taken to check the actual system being used to report activity as the NPA/ULF and MGO numbers are derived from completely different types of tests.

MEDICINAL MANUKA HONEY - MGO GRADES EXPLAINED

Premium medicinal Manuka honey is defined as high-grade MGO Manuka honey that is scientifically tested for its curative compounds such as antioxidants, bioflavonoids and immune boosting enzymes.



This honey presents higher levels of MGO (Methylglyoxal) when compared to table-grade Manuka honey and is

naturally bio-active with anti-bacterial, antimicrobial, antiseptic, and anti-inflammatory properties for the holistic prevention and treatment of various health ailments.






Commonly known uses include: **burns, wounds, infection, oral care, ulcers, gut health, cold & flu, sore throat, cough, immunity building, general wellbeing,** and **beauty and skincare.**

The higher the MGO number of the Manuka Honey, the stronger its antimicrobial bioactivity levels and the more potent the natural medicinal benefits.

The MGO levels of Biosota's high strength medicinal Manuka honey ranges are regularly tested and verified authentic by independent accredited laboratories both in Australia and New Zealand.

ANTIBACTERIAL BIOACTIVITY & SUGGESTED USAGE

THIS CHART DISPLAYS THE SUGGESTED MEDICINAL USAGE FOR VARYING GRADES OF MANUKA HONEY. THE HIGHER THE MGO RATING, THE STRONGER AND MORE POTENT THE MANUKA HONEY AND ITS USES.

RANGE STRENGTH	MGO	NPA/ULF	SUGGESTED USE
 THE ULTIMATE STRENGTH MGO 2100+ (LIMITED EDITION)	2100+	36+	MEDICAL GRADE USE BURNS WOUNDS ULCERS INFECTIONS
	1717+	31+	
 SUPERIOR STRENGTH MGO 1000+ TO 1900+ (RARE)	1443+	28+	GUT HEALTH ORAL CARE
	1200+	25+	
 HIGH MEDICINAL STRENGTH MGO 500+ TO 990+	880+	20+	THERAPEUTIC GRADE USE COLDS & FLU SORE THROATS
	550+	15+	
 MEDICINAL STRENGTH MGO 300+ TO 490+ (MINIMAL BIOACTIVITY REQUIRED FOR MEDICINAL USE)	400+	12+	COUGHS
	300+	11+	
 EVERYDAY STRENGTH MGO 30+ TO 290+	260+	10+	TABLE GRADE USE WELLBEING IMMUNITY SKINCARE
	150+	5+	
	30+	2+	

Statements made have not been evaluated by the TGA (Australian Therapeutic Goods Administration) or FDA (US Food & Drug Administration). Products sold are not intended to diagnose, treat, cure, or prevent any disease.

HOW TO USE AND APPLY MEDICINAL MANUKA HONEY

Medicinal Manuka honey usage and dosage suggestions made are based on our own direct experience, from feedback received and collated from real customers, and from the latest published scientific journals and research papers.

Dosage and individual results may vary. Please consult your healthcare provider before beginning any treatment and if symptoms persist.

i (Allergy advice: honey should not be consumed by children under the age of 1 and people with allergies to honey and bee products).



FOR BURNS, WOUNDS AND INFECTIONS

[Try Superior Strength Medicinal Manuka Honey Range - MGO 1000+ to 1900+](#)

For topical application for a burn, wound, or infection liberally apply a superior grade medicinal Manuka honey directly to the affected areas. Cover with a clean and sterile dressing: bandage, gauze, compress, or cling film (optional). Leave overnight and change the dressing when saturated, or as needed.

At Biosota Organics, our passion lies in the natural healing power of Manuka honey.

Our premium Australian Manuka Honey is cultivated by nature with a whole lot of science and love as our secret ingredients.

MEDICINAL MANUKA HONEY

i Disclaimer: Statements made have not been evaluated by the TGA (Australian Therapeutic Goods Administration) or FDA (U.S. Food & Drug Administration). Products sold are not intended to diagnose, treat, cure, or prevent any disease. Manuka honey is not intended to be a substitute for other medicines or advice and is best used in conjunction with any existing treatment plans. Please consult your healthcare professional before beginning any treatment.

REFERENCES CITED:

- Molan, P.C., Russell, K.M. (1988). Non-peroxide antibacterial activity in some New Zealand honeys. *J. Apic. Res.* 27: 62–67.
- Cokcetin, N.N., Pappalardo, M., Campbell, L.T., Brooks, P., Carter, D.A., Blair, S.E. & Harry, E.J. (2016). The Antibacterial Activity of Australian *Leptospermum* Honey Correlates with Methylglyoxal Levels. *PLoS One*, vol. 11, no. 12, p.e0167780.
- Windsor S., Pappalardo M., Brooks P., Williams S., and Manley-Harris M. A. (2012). Convenient new analysis of dihydroxyacetone and methylglyoxal applied to Australian *Leptospermum* honeys. *Journal of Pharmacognosy and Phytotherapy*, 4(1).
- Blair, S.E., Cokcetin, N.N., Harry, E.J., and Carter, D.A. (2009). The unusual antibacterial activity of medical-grade *Leptospermum* honey: antibacterial spectrum, resistance and transcriptome analysis. *European Journal of Clinical Microbiology and Infectious Disease*, 28: 1199. <https://doi.org/10.1007/s10096-009-0763-z>.
- Thompson, J. (1989). A revision of the genus *Leptospermum* (Myrtaceae). *Telopea*. 3 (3): 301–449.
- Van Eaton, Cliff. (2014). *Manuka: The biography of an extraordinary honey*. Exisle Publishing.
- Fleming, C.A. (1975). The geological history of New Zealand and its biota. In Kuschel, G. (ed), *Biogeography and Ecology in New Zealand*. (W.Junk The Hague).
- Adams, C.J., Boulton, C.H., Deadman, B.J., Farr, J.M., Grainger, M.N.C., Manley-Harris, M., et al. (2008). Isolation by HPLC and characterisation of the Bioactive fraction of New Zealand Manuka (*Leptospermum scoparium*) honey. *Carbohydr. Res.* 343, 651–659. doi:10.1016/j.carres.2007.12.011
- Mavric, E., Wittmann, S., Barth, G. and Henle, T. (2008). Identification and quantification of methylglyoxal as the dominant antibacterial constituent of Manuka (*Leptospermum scoparium*) honeys from New Zealand. *Mol. Nutr. Food Res.*, 52: 483–489. doi:10.1002/mnfr.200700282
- Adams, C.J., Manley-Harris, M., and Molan, P.C. (2009). The origin of methylglyoxal in New Zealand Manuka (*Leptospermum scoparium*) honey. *Carbohydr. Res.* 344, 1050–1053. doi:10.1016/j.carres.2009.03.020 (superbugs), they cannot develop resistance to the activity of Manuka honey (Blair et al. 2009).

FOR MORE INFORMATION ON THE RESEARCH INTO AUSTRALIAN MANUKA:

<https://ozhoneyproject.wordpress.com/>

- Carter, D.A., Blair, S.E., Cokcetin, N.N., Bouzo, D., Brooks, P., Schothauer, R. & Harry, E.J. (2016). Therapeutic Manuka Honey: No Longer So Alternative. *Frontiers in Microbiology*, vol. 7, p. 569
- Irish, J., Blair, S. & Carter, D.A. (2011). The antibacterial activity of honey derived from Australian flora. *PloS one*, vol. 6, no. 3, p. e18229.
- Windsor, S., Pappalardo, M., Brooks, P., Williams, S., Manley-Harris, M. (2012). A convenient new analysis of dihydroxyacetone and methylglyoxal applied to Australian *Leptospermum* honeys. *Journal of Pharmacognosy and Phytotherapy* 4, no. 1: 6–11.
- Pappalardo M, Pappalardo L, Brooks P (2016) Rapid and Reliable HPLC Method for the Simultaneous Determination of Dihydroxyacetone, Methylglyoxal and 5-Hydroxymethylfurfural in *Leptospermum* Honeys. *PLoS ONE* 11(11): e0167006.

<https://doi.org/10.1371/journal.pone.0167006>