

Analytica Laboratories Limited Ruakura Research Centre 10 Bisley Road Hamilton 3214, New Zealand Ph +64 (07) 974 4740 sales@analytica.co.nz www.analytica.co.nz

Certificate of Analysis

Biosota Organics PTY Ltd Unit 10 / 475 Scottsdale Drive

Gold Coast QLD 4227 Attention: Andrey Zubko Phone: +61 0409839773

Email: andrey.zubko@biosota.com

Lab Reference: 23-16864

Submitted by:

Date Received: 3/06/2023
Testing Initiated: 7/06/2023
Date Completed: 8/06/2023
Order Number: 00016
Reference: 000016

Report Comments

Samples were collected by yourselves (or your agent) and analysed as received at Analytica Laboratories. Samples were in acceptable condition unless otherwise noted on this report.

Specific testing dates are available on request.

Results Summary

3in1 in Honey

Laboratory ID	Sample ID	Dihydroxyacetone (DHA)	Methylglyoxal (MG/MGO)	Non-Peroxide Activity* (NPA)	Hydroxymethylfurfural (HMF)
Units Reporting Limit		mg/kg 40	mg/kg 8	%w/v phenol eq. 1.3	mg/kg 1
23-16864-5	BN: 224436_MGO 1450+	1,620	1,410	27.6	56.8

3in1 in Honey Approver:

Alicia Laing, BSc. Technician

Leptosperin in Honey

Laboratory ID	Sample ID	Leptosperin
	mg/kg 20	
23-16864-5	BN: 224436_MGO 1450+	1,350

Leptosperin in Honey Approver:

Alicia Laing, BSc. Technician

All tests reported herein have been performed in accordance with the laboratory's scope of accreditation with the exception of tests marked *, which are not accredited.

This test report shall not be reproduced except in full, without the written permission of Analytica Laboratories.



Method Summary

3in1 Determination of Dihydroxyacetone (DHA), Methylglyoxal (MG/MGO) and Hydroxymethylfurfural (HMF) by aqueous

extraction, derivatisation, and UPLC (diode array) analysis in accordance with in-house procedures.

NPA Non-Peroxide Activity (NPA) values are not directly measured by the laboratory, but are calculated from the measured

methylglyoxal concentration in the honey according to the requirements of the client. The calculation is based on published data(†) comparing the NPA and methylglyoxal concentration measured in a range of honey samples. These calculated values are not accredited by IANZ and do not imply that the honey is or is not manuka honey.

NPA values less than 5 are an estimate based on extrapolation of the relationship between methylglyoxal and NPA

(†) Isolation by HPLC and characterisation of the bioactive fraction of New Zealand manuka (Leptospermum scoparium) honey. C. J. Adams, et al. Carbohydrate Research 343 (2008) 651-659. And, Corrigendum to "Isolation by HPLC and characterization of the bioactive fraction of New Zealand manuka (Leptospermum scoparium) honey" [Carbohydr. Res.

343 (2008) 651]. Carbohydrate Research 344 (2009) 2609. C. J. Adams, et al.

Leptosperin Aqueous extraction, dilution, analysis by UPLC-PDA/FLD in accordance with in-house procedures.