

MEDICAMENTOS DE HEEL



# LIBRO DE REFERENCIA

ALERGIAS

## 5 ALERGIAS

### Desplácese por los usos de los productos principales

Desplácese por los síntomas principales →

Productos principales de Heel para los siguientes síntomas y afecciones	Luffeel® Nebulizador nasal	Luffeel® Comprimididos	Curtis compositum	Oculoheel® (colirio)	Hepeel®	Mucosa comp.	Engystol®	Coenzyme compositum	Lympho- myosot®
<b>Producto principal</b> ● <b>Producto complementario</b>									
Estomudos									●
Rinorrea									●
Prurito en las mucosas nasofaríngeas									●
Obstrucción nasal									●
Congestión nasal y sinusal									●
Pólipos nasales									●
Síntomas cutáneos: prurito, pápulas, urticaria, tumefacción localizada*									●
Síntomas oculares: ardor, prurito, dolor, enrojecimiento, lagrimeo									
Actuación sobre la fisiopatología además de las recomendaciones terapéuticas anteriores									
Facilita la eliminación por el hígado de productos de desecho que desencadenan alergias, como la histamina									
Regulación y apoyo de las funciones de las mucosas									
Restablecimiento del equilibrio del sistema inmunitario									
Apoyo de la respiración celular en la enfermedad crónica									
Profilaxis en alergias estacionales (comenzar 12 semanas antes de la estación alérgica)									

\* En el eccema intenso, la inmunomodulación con Engystol y Curtis compositum debe realizarse durante al menos 4 semanas antes de iniciar la destoxicación con Lymphomyosot

Producto	Indicaciones	
Luffeel®/Luffa comp.	<ul style="list-style-type: none"> <li>Rinitis alérgica</li> </ul>	1.4
Cutis compositum	<ul style="list-style-type: none"> <li>Dermatitis</li> <li>Dermatitis alérgica</li> <li>Dermatitis seborreica</li> <li>Neurodermatitis</li> </ul>	1.7
Oculoheel®	<ul style="list-style-type: none"> <li>Conjunctivitis</li> <li>Irritación ocular</li> <li>Fatiga ocular</li> </ul>	1.8
Hepeel®	<ul style="list-style-type: none"> <li>Mejoría de la función metabólica del hígado</li> <li>Facilitación de la destoxificación</li> </ul>	1.9
Mucosa compositum	<ul style="list-style-type: none"> <li>Regulación y apoyo de las funciones de las mucosas</li> </ul>	1.10
Engystol®	<ul style="list-style-type: none"> <li>Apoyo del sistema inmunitario</li> </ul>	1.11
Coenzyme compositum	<ul style="list-style-type: none"> <li>Estimulación de sistemas enzimáticos intracelulares bloqueados</li> </ul>	1.12
Lymphomyosot®	<ul style="list-style-type: none"> <li>Trastornos del flujo linfático</li> <li>Infecciones recurrentes</li> <li>Drenaje y destoxificación</li> </ul>	1.13

**Nota importante:**

Este folleto contiene información médica útil basada en datos científicos. Esta información no constituye asesoramiento médico ni tratamiento y **no sustituye a la exploración médica**; los pacientes deben consultar a un médico antes de tomar decisiones sobre el tratamiento.



## Combata los síntomas de las alergias y de la rinitis alérgica estacional

- Rinitis alérgica (tratamiento y profilaxis)



### Recomendaciones posológicas



**Nebulizador nasal**

En general:	1 o 2 nebulizaciones en cada fosa nasal de 3 a 5 veces al día
Niños menores de 6 años:	1 nebulización de 3 a 4 veces al día

## Protéjase de las alergias y de la rinitis alérgica estacional

- Rinitis alérgica (tratamiento y profilaxis)



### Recomendaciones posológicas




**Comprimidos**

En general:	1 comprimido 4 veces al día
En afecciones agudas:	1 comprimido cada 15 minutos, durante 2 horas como máximo

## Datos clínicos

- En un ensayo de equivalencia aleatorizado y doble ciego con 146 pacientes, se compararon la eficacia y la tolerancia de Luffa comp.-Heel nebulizador nasal con las del cromoglicato disódico en nebulizador en el tratamiento de la rinitis alérgica estacional (fiebre del heno).



Referencia:

Weiser M., Gegenheimer L.H., Klein P. Seasonal allergic rhinitis: A homeopathic nasal spray as effective as cromoglicic acid. *Frosch Komplementärmed/ Research in Complementary Medicine*. 1999;6:142-148

- Luffa comp.-Heel mostró un inicio de acción comparable.
- Con ambos tratamientos el efecto terapéutico principal tuvo lugar en la primera semana.
- Sin efectos adversos sistémicos; efectos adversos locales muy poco frecuentes (p. ej., escozor en las fosas nasales)

## Perfiles de pacientes

- **Carla | 19 años de edad**

Soltera

Antecedentes médicos: dermatitis atópica en edad infantil confirmada por niveles elevados de anticuerpos IgE específicos contra el polen, la caspa animal y los ácaros del polvo.

**Enfermedad actual:** rinoconjuntivitis alérgica con prurito, enrojecimiento, dolor y lagrimeo de los ojos en caso de exposición a los alérgenos anteriormente indicados.



- **Ana | 26 años de edad**

Casada, 1 hijo

Trabajo: profesora escolar

Antecedentes médicos: prueba de punción cutánea positiva al polen y a ciertos tipos de gramíneas.

**Enfermedad actual:** crisis de estornudos, rinorrea y prurito en paladar y oídos seguido de congestión nasal que interfiere en su calidad de vida durante la estación alérgica.

La paciente desearía un tratamiento que pudiera contribuir a prevenir los síntomas.



- **Roberto | 32 años de edad**

Soltero

Occupation: Artist

Antecedentes familiares: la madre padece dermatitis atópica en situaciones de estrés y la hermana padece rinitis alérgica estacional.

Antecedentes médicos: rinitis alérgica estacional desde la adolescencia y sinusitis crónica.

**Enfermedad actual:** episodios de estornudos y rinorrea con congestión nasosinusal crónica, con malestar y somnolencia.



# Cutis compositum

## Tratamiento integral de los trastornos cutáneos

- Dermatitis
- Dermatitis alérgica
- Dermatitis seborreica
- Neurodermatitis



## Recomendaciones posológicas



### Solución inyectable

i.m., s.c., i.d., i.v.

POSOLOGÍA AGUDA:

1 ampolla al día

POSOLOGÍA ESTÁNDAR:

1 ampolla entre 1 y 3 veces por semana



### Comprimidos

i.m., s.c., i.d., i.v.

POSOLOGÍA AGUDA:

1 comprimido cada 1/2-1 hora el primer día hasta un máximo de 12 comprimidos; posteriormente, continuar con 1 comprimido 3 veces al día

POSOLOGÍA ESTÁNDAR:

1 comprimido 3 veces al día

# Oculoheel®

## Calma la inflamación ocular alérgica

- Conjunctivitis
- Irritación ocular
- Fatiga ocular



### Recomendaciones posológicas



Colirio

En general:

instilar 1 gota en el ojo afectado, 3 veces al día





## Facilita la eliminación de productos de desecho que desencadenan alergias

- Mejora la función metabólica del hígado
- Facilita la detoxificación



## Recomendaciones posológicas



### Comprimidos

En general: 1 comprimido disuelto en la boca, 3 veces al día



### Solución inyectable i.m., s.c., i.d., i.v.

Adultos  
(y niños  
> 12 años)

#### POSOLOGÍA AGUDA

1 ampolla diaria durante  
3 días; posteriormente,  
continuar con la posología  
estándar

#### POSOLOGÍA ESTÁNDAR

1 ampolla  
entre 1 y 3 veces  
por semana

Si desea más información,  
consulte el prospecto del envase.



# Mucosa compositum

Para un funcionamiento  
sano de las mucosas

- Regulación y apoyo de las funciones de las mucosas, incluido el tubo digestivo



## Recomendaciones posológicas

		POSOLOGÍA AGUDA	POSOLOGÍA ESTÁNDAR
 Tablets	Adultos (y niños > 12 años)	1 comprimido cada 1/2-1 hora, hasta un máximo de 12 comprimidos al día	1 comprimido 3 veces al día
	Niños de 0-2 años	1 comprimido cada 1-2 horas, hasta un máximo de 4 comprimidos al día	1 comprimido 1 vez al día
	Niños de 2-5 años	1 comprimido cada 1-2 horas, hasta un máximo de 6 comprimidos al día	1 comprimido 1-2 veces al día
	Niños de 6-11 años	1 comprimido cada 1-2 horas, hasta un máximo de 8 comprimidos al día	1 comprimido 2 veces al día
 Solución inyectable i.m., s.c., i.d., i.v.	Adultos (y niños > 12 años)	–	1 ampolla 1-3 veces por semana
	Niños de 2-5 años	–	1/2 ampolla 1-3 veces por semana
Si desea más información, consulte el prospecto del envase.	Niños de 6-11 años	–	2/3 ampolla 1-3 veces por semana





**Ayuda al sistema inmunitario  
a combatir las infecciones virales,  
como el resfriado y la gripe**

- Resfriado común
- Infecciones virales leves
- Enfermedades seudogripales



## Recomendaciones posológicas

		POSOLOGÍA AGUDA	POSOLOGÍA CRÓNICA
 <b>Tablets</b>	Adultos (y niños > 12 años)	1 comprimido cada 1/2-1 hora, hasta un máximo de 12 comprimidos al día	1 comprimido 3 veces al día
	Niños de 0-2 años	1 comprimido cada 1-2 horas, hasta un máximo de 4 comprimidos al día	1 comprimido 1 vez al día
	Niños de 2-5 años	1 comprimido cada 1-2 horas, hasta un máximo de 6 comprimidos al día	1 comprimido 1-2 veces al día
	Niños de 6-11 años	1 comprimido cada 1-2 horas, hasta un máximo de 8 comprimidos al día	1 comprimido 2 veces al día
 <b>Solución inyectable</b> i.m., s.c., i.d., i.v.	Adultos (y niños > 12 años)	1 ampolla al día	1 ampolla 1-3 veces por semana
	Niños de 2-5 años	1/2 ampolla al día	1/2 ampolla 1-3 veces por semana
	Niños de 6-11 años	2/3 ampolla al día	2/3 ampolla 1-3 veces por semana
Si desea más información, consulte el prospecto del envase.			






## Regulador celular multifocal

- Estimulación de sistemas enzimáticos intracelulares bloqueados



### Recomendaciones posológicas

		POSOLOGÍA AGUDA	POSOLOGÍA CRÓNICA
 <b>Tablets</b>	Adultos (y niños > 12 años)	1 comprimido cada 1/2-1 hora, hasta un máximo de 12 comprimidos al día	1 comprimido 3 veces al día
	Niños de 0-2 años	1 comprimido cada 1-2 horas, hasta un máximo de 4 comprimidos al día	1 comprimido 1 vez al día
	Niños de 2-5 años	1 comprimido cada 1-2 horas, hasta un máximo de 6 comprimidos al día	1 comprimido 1-2 veces al día
	Niños de 6-11 años	1 comprimido cada 1-2 horas, hasta un máximo de 8 comprimidos al día	1 comprimido 2 veces al día
 <b>Solución inyectable</b> i.m., s.c., i.d., i.v.	Adultos (y niños > 12 años)	1 ampolla al día	1 ampolla 1-3 veces por semana
	Niños de 2-5 años	1/2 ampolla al día	1/2 ampolla 1-3 veces por semana
	Niños de 6-11 años	2/3 ampolla al día	2/3 ampolla 1-3 veces por semana
Si desea más información, consulte el prospecto del envase.			





## Restablece el flujo linfático para acelerar la recuperación

- Trastornos del flujo linfático
- Infecciones recurrentes
- Drenaje y detoxificación



### Recomendaciones posológicas

		POSOLOGÍA AGUDA	POSOLOGÍA CRÓNICA
 Tablets	Adultos (y niños > 12 años)	1 comprimido cada 1/2-1 hora, hasta un máximo de 12 comprimidos al día	1 comprimido 3 veces al día
	Niños de 0-2 años	1 comprimido cada 1-2 horas, hasta un máximo de 4 comprimidos al día	1 comprimido 1 vez al día
	Niños de 2-5 años	1 comprimido cada 1-2 horas, hasta un máximo de 6 comprimidos al día	1 comprimido 1-2 veces al día
	Niños de 6-11 años	1 comprimido cada 1-2 horas, hasta un máximo de 8 comprimidos al día	1 comprimido 2 veces al día
 Solución inyectable i.m., s.c., i.d., i.v.	Adultos (y niños > 12 años)	1 ampolla al día	1 ampolla 1-3 veces por semana
	Niños de 2-5 años	1/2 ampolla al día	1/2 ampolla 1-3 veces por semana
	Niños de 6-11 años	2/3 ampolla al día	2/3 ampolla 1-3 veces por semana

Si desea más información, consulte el prospecto del envase.

## Product Information

### Coenzyme comp. (Coenzyme compositum)

Solution for injection

**Compositions:** 1 ampoule of 2,2 ml (=2,2 g) contains:

Acidum ascorbicum D6 22,0 mg, Acidum alpha-ketoglutaricum D8 22,0 mg, Acidum cis-aconiticum D8 22,0 mg, Acidum citricum D8 22,0 mg, Acidum fumaricum D8 22,0 mg, Acidum malicum D8 22,0 mg, Acidum succinicum D8 22,0 mg, Adenosinum triphosphoricum D10 22,0 mg, Barium oxalsuccinicum D10 22,0 mg, Beta vulgaris rubra D4 22,0 mg, Coenzym A D8 22,0 mg, Cysteinum D6 22,0 mg, Hepar sulfuris D10 22,0 mg, Nadidum D8 22,0 mg, Natrium pyruvicum D8 22,0 mg, Natrium riboflavinum phosphoricum D6 22,0 mg, Nicotinamidum D6 22,0 mg, Pulsatilla pratensis D6 22,0 mg, Pyridoxinum hydrochloricum D6 22,0 mg, Sulfur D10 22,0 mg, Thiaminum hydrochloricum D6 22,0 mg, Acidum thiocticum D6 22,0 mg, Cerium oxalicum D8 22,0 mg, Magnesium oroticum dihydricum D6 22,0 mg, Manganum phosphoricum D6 22,0 mg, Natrium diethylalacetatum D6 22,0 mg

**Indications:** Stimulation of blocked enzymatic systems in degenerative diseases as well as in defective enzymatic functions (cellular phases). Special note: If symptoms persist or worsen you should consult a doctor or healthcare professional. For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

**Contraindications:** None.

**Side effects:** None known.

**Interactions with other medication:** None known.

**Dosage:** In general, 1 ampoule i.m., s.c., i.d. 3-1 times weekly.

**Package sizes:** Packs containing 5, 10, 50 and 100 ampoules of 2,2 ml. (8511)

### Coenzyme comp. (Coenzyme compositum)

Tablets

**Compositions:**

Acidum cis-aconiticum D8 1,0 mg, Acidum ascorbicum D6 1,0 mg, Acidum citricum D8 1,0 mg, Acidum fumaricum D8 1,0 mg, Acidum alpha-ketoglutaricum D8 1,0 mg, Acidum malicum D8 1,0 mg, Acidum succinicum D8 1,0 mg, Adenosinum triphosphoricum D10 1,0 mg, Beta vulgaris rubra D6 1,0 mg, Coenzym A D8 1,0 mg, Cysteinum D6 1,0 mg, Nadidum D8 1,0 mg, Natrium pyruvicum D8 1,0 mg, Natrium riboflavinum phosphoricum D6 1,0 mg, Nicotinamidum D6 1,0 mg, Pulsatilla pratensis D6 1,0 mg, Pyridoxinum hydrochloricum D6 1,0 mg, Sulfur D10 1,0 mg, Thiaminum hydrochloricum D6 1,0 mg, Barium oxalsuccinicum D10 1,0 mg, Cerium oxalicum D8 1,0 mg, Hepar sulfuris D10 1,0 mg, Magnesium oroticum dihydricum D6 1,0 mg, Manganum phosphoricum D6 1,0 mg, Natrium diethylalacetatum D6 1,0 mg

**Indications:** Special note: For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

**Contraindications:** None.

**Side effects:** None known.

**Interactions with other medication:** None known.

**Dosage:** Standard dosage: Adults (and children 12 yrs. and older): 1 tablet to be dissolved in the mouth 3x daily.

**Package sizes:** (41946)

### Cutis comp. (Cutis compositum N)

Solution for injection

**Compositions:** 1 ampoule of 2,2 ml (=2,2 g) contains:

Acidum alpha-ketoglutaricum D10 22,0 mg, Acidum formicum D198 22,0 mg, Acidum fumaricum D10 22,0 mg, Acidum phosphoricum D6 22,0 mg, Aesculus hippocastanum D6 22,0 mg, Ammonium bituminosulfonicum D28 22,0 mg, Arctium D6 22,0 mg, Calcium fluoratum D13 22,0 mg, Cortisonum aceticum D28 22,0 mg, Cutis suis D8 22,0 mg, Funiculus umbilicalis suis D10 22,0 mg, Galium aparine D6 22,0 mg, Glandula suprarenalis suis D10 22,0 mg, Hepar suis D10 22,0 mg, Ledum palustre D4 22,0 mg, Mercurius solubilis Hahnemanni D13 22,0 mg, Natrium diethylalacetatum D10 22,0 mg, Placenta totalis suis D10 22,0 mg, Selenium D10 22,0 mg, Spleen suis D10 22,0 mg, Strychnos ignatii D6 22,0 mg, Sulfur D10 22,0 mg, Thallium sulfuricum D13 22,0 mg, Thuja occidentalis D8 22,0 mg, Urtica urens D4 22,0 mg

**Indications:** Stimulation of the defensive system in dermatitis, dermatosis, allergy, scleroderma, acne vulgaris, erythematodes, dermatomycosis, crusta lactea, kondylomatosa, impetigo contagiosa, neurodermatitis, microsporia, mycosis interdigitalis, decubitus, Dupuytren's contracture, alopecia, naevi, Quincke's oedema, urticariasis.

**Special note:** If symptoms persist or worsen you should consult a doctor or healthcare professional. For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

**Contraindications:** None.

**Side effects:** None known.

**Interactions with other medication:** None known.

**Dosage:** In acute disorders daily, otherwise 1 ampoule 3-1 times weekly i.m., s.c., i.d., or if necessary i.v.

**Package sizes:** Packs containing 5, 10, 50 and 100 ampoules of 2,2 ml. (9477)

### Cutis comp. (Cutis compositum)

Tablets

**Compositions:** 1 tablet contains:

Acidum formicum D198 1,0 mg, Acidum fumaricum D10 1,0 mg, Acidum phosphoricum D6 1,0 mg, Aesculus hippocastanum D6 1,0 mg, Ammonium bituminosulfonicum D28 1,0 mg, Arctium D6 1,0 mg, Cortisonum aceticum D28 1,0 mg, Cutis suis D8 1,0 mg, Funiculus umbilicalis suis D10 1,0 mg, Galium aparine D6 1,0 mg, Glandula suprarenalis suis D10 1,0 mg, Hepar suis D10 1,0 mg, Ledum palustre D4 1,0 mg, Placenta totalis suis D10 1,0 mg, Spleen suis D10 1,0 mg, Strychnos ignatii D6 1,0 mg, Sulfur D10 1,0 mg, Thuja occidentalis D8 1,0 mg, Urtica urens D4 1,0 mg, Calcium fluoratum D13 1,0 mg, Mercurius solubilis Hahnemanni D13 1,0 mg, Selenium D10 1,0 mg

Contains lactose

**Indications:** Supportive treatment of chronic skin conditions.

**Special note:** This remedy contains lactose. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption syndrome should not take this medicine. No studies on the effects on the ability to drive and use machines have been performed. None expected. For this product no clinical data on pregnancy and lactation are available.

Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation.

No adverse effects have so far been reported.

**Contraindications:** None known

**Side effects:** Not known

**Interactions with other medication:** None known

**Dosage:** Adults (and children over 12 years):

**Standard dose:** 1 tablet 3x/daily; Acute dose 1 tablet every 1/2 to 1 hr. for the first day up to 12x, then continue with 1 tablet 3x/daily.

**Package sizes:** (44179)

### Engystol®

Solution for injection

**Compositions:** 1 ampoule of 1,1 ml (=1,1 g) contains:

Sulfur D10 3,30 mg, Vincetoxicum hirundinaria D6 6,60 mg, Vincetoxicum hirundinaria D10 6,60 mg, Vincetoxicum hirundinaria D30 6,60 mg, Sulfur D4 3,30 mg

**Indications:** To activate the non-specific defensive mechanism, particularly in the case of influenza and questionable virus diseases.

**Special note:** If symptoms persist or worsen you should consult a doctor or healthcare professional. For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

**Contraindications:** None.

**Side effects:** In rare cases, skin rashes and itching might follow the use of Engystol. In this case the product should be discontinued and the patient's doctor consulted.

Le regarda sin traducir los textos específicos de su país

**Interactions with other medication:** None known.

**Dosage:** In acute disorders 1 ampoule daily, otherwise 3-1 times weekly i.m., s.c., i.d., i.v.

**Package sizes:** Packs containing 5, 10, 50 and 100 ampoules of 1,1 ml. (8546)

#### Engystol®

Tablets

**Compositions:** 1 tabl. cont.:

Sulfur Trit. D4 37,5 mg, Sulfur D10 37,5 mg, Vincetoxicum hirundinaria D6 75,0 mg, Vincetoxicum hirundinaria D10 75,0 mg, Vincetoxicum hirundinaria D30 75,0 mg

Contains lactose! Please see package insert!

**Indications:** To activate the non-specific defensive mechanism, particularly in the case of influenza and questionable virus diseases.

**Special note:** This remedy contains Lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product. If symptoms persist or worsen you should consult a doctor or healthcare professional. For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation.

No adverse effects have so far been reported.

**Contraindications:** None.

**Side effects:** In individual cases skin rashes may occur after use of Engystol, in which case the product should be discontinued and the patient's doctor consulted.

Interactions with other medication: None known.

**Dosage:** In general, 1-3 tablets to be dissolved in the mouth 3 times daily.

**Package sizes:** Packs containing 50 and 250 tablets. (9830)

#### Hepeel® (Hepeel® N)

Solution for injection

**Compositions:** 1 ampoule of 1,1 ml (= 1,1 g) contains:

Chelidonium majus D6 1,10 mg, Citrullus colocynthis D5 3,30 mg, Phosphorus D5 0,55 mg, Veratrum album D5 2,20 mg, Myristica fragrans D3 1,10 mg, Cinchona pubescens D2 1,10 mg, Lycopodium clavatum D2 1,10 mg, Silybum marianum D1 0,55 mg

**Indications:** Supportive treatment in functional disorders of the liver.

**Contraindications:** Because of the constituents Celandine (Chelidonium) and Veratrum do not use during pregnancy and breastfeeding. Do not use in case of hypersensitivity to Carduus marianus or other plants of the daisy family.

**Side effects:** In rare cases, reactions of hypersensitivity such as skin allergy or fever may occur after exposure to drugs that contain quinine. In such cases, you must see a doctor. Note: A quinine or quinidine sensitization may occur.

Interactions with other medication: None known.

**Dosage:** Unless otherwise prescribed,

**Standard dosage:**

Adult (and children > 12 years) 1 ampoule 1 to 3 times weekly

Paediatric: (Only if required 12 - 18 years: 1 ampoule 1 to 3 times weekly)

Acute or initial dosage:

Adult (and children > 12 years): 1 ampoule daily for 3 days, then continue with standard dosage

Paediatric: (Only if required 12 - 18 years: 1 ampoule daily for 3 days, then continue with standard dosage.)

Package sizes: Packs containing 5, 10, 50 and 100 ampoules of 1,1 ml. (46791)

#### Hepeel®

Tablets

**Compositions:** 1 tabl. cont.:

Chelidonium majus D4 30,0 mg, Citrullus colocynthis D6 90,0 mg, Myristica fragrans D4 1,10 mg, Phosphorus D6 15,0 mg, Veratrum album D6 60,0 mg, Cinchona pubescens D3 30,0 mg, Lycopodium clavatum D3 30,0 mg, Silybum marianum D2 5,5 mg

Contains lactose! Please see package insert!

**Indications:** Primary and secondary functional disorders of the liver; damaged liver. Special note: This remedy contains Lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product. If symptoms persist or worsen you should consult a doctor or healthcare professional.

**Contraindications:** Do not use in case of hypersensitivity to Carduus marianus or other plants of the daisy family (Asteraceae) or in case of quinine hypersensitivity. Due to the constituent Celandine (Chelidonium) do not use during pregnancy and lactation. In case of existing liver conditions or history thereof or simultaneous use of hepatotoxic substances use only after consultation with the physician.

**Side effects:** In rare cases, reactions of hypersensitivity such as skin allergy or fever may occur after exposure to drugs that contain quinine. In such cases, you must see a doctor. Note: A quinine or quinidine sensitization may occur. During the treatment with drugs containing Celandine alkaloids there have been individual reports of increased liver function enzymes (transaminases) and serum bilirubin elevations up to drug-induced jaundice (drug-related toxic hepatitis), which normalized or resolved after discontinuation of the product.

Interactions with other medication: None known.

**Dosage:** In general, 1 tablet to be dissolved in the mouth 3 times daily.

**Package sizes:** Packs containing 50 and 250 tablets. (9737)

#### Luffeel®

Nasal spray

**Compositions:** 100 g cont.:

Histaminum D12 5,0000 g, Histaminum D30 5,0000 g, Histaminum D200 5,0000 g, Luffa operculata D4 10,0000 g, Luffa operculata D12 10,0000 g, Luffa operculata D30 10,0000 g, Sulfur D12 5,0000 g, Sulfur D30 5,0000 g, Sulfur D200 5,0000 g, Thryallis glauca D4 10,0000 g, Thryallis glauca D12 10,0000 g, Thryallis glauca D30 10,0000 g, in isotonic sodium chloride solution. Preserved with benzalkonium chloride 0,01 %.

**Indications:** Hay fever. Special note: If symptoms persist or worsen you should consult a doctor or healthcare professional. For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

**Contraindications:** Do not use in case of hypersensitivity to benzalkonium chloride.

**Side effects:** Irritation of the nasal mucosae with burning and increased nasal secretion may occur in very rare cases after use of Luffeel nasal spray, in which case the product should be discontinued. In very rare cases bronchospasm may occur in predisposed asthmatic patients due to benzalkonium chloride.

Interactions with other medication: None known.

**Dosage:** In general, spray 1-2 shots into each nostril 3-5 times daily; for children under 6 years, 1 shot 3-4 times daily.

**Package sizes:** Atomiser without propellant, 20 ml. (9864)

#### Luffeel®

Tablets

**Compositions:** 1 tabl. cont.:

Arsenum iodatum D8 25,0 mg, Lobelia inflata D6 25,0 mg, Luffa operculata D12 25,0 mg, Aralia racemosa D1 25,0 mg, Contains lactose! Please see package insert!

**Indications:** Hay fever.

**Special note:** This remedy contains Lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product. If symptoms persist or worsen you should consult a doctor or healthcare professional. For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

**Contraindications:** Do not use in thyroid conditions without seeking medical advice.

**Side effects:** None known

Interactions with other medication: None known.

**Dosage:** In general, 1 tablet to be dissolved in the mouth 3 times daily. In acute disorders, 1 tablet every 15 minutes (over a period lasting up to 2 hours).

**Package sizes:** Packs containing 50 and 250 tablets. (9831)

#### Lymphomyosot® (Lymphomyosot® N)

Solution for injection

**Compositions:** 1,1 ml (1,1 g) cont.:

Araneus diadematus D6 0,55 mg, Calcium phosphoricum D12 0,55 mg, Equisetum hiemale D4 0,55 mg, Ferrum iodatum D12 1,10 mg, Fumaria officinalis D4 0,55 mg, Gentiana lutea D5 0,55 mg, Geranium robertianum D4 1,10 mg, Levothyroxinum D12 0,55 mg, Myosotis arvensis D3 0,55 mg, Nasturtium officinale D4 1,10 mg, Natrium sulfuricum D4 0,55 mg, Pinus sylvestris D4 0,55 mg, Scrophularia nodosa D3 0,55 mg, Smlax D6 0,55 mg, Teucrium scorodonia D3 0,55 mg, Veronica officinalis D3 0,55 mg

Le rogamos de introducir los textos específicos de su país

**Indications:** Lymphatism (tendency to hypertrophy of the lymphatic organs, tendency to formation of oedemas and to susceptibility to infection); glandular swelling; tonsillar hypertrophy and chronic tonsillitis. Special note: If symptoms persist or worsen you should consult a doctor or healthcare professional. For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

**Contraindications:** Do not use in thyroid conditions without seeking medical advice.

**Side effects:** In very rare cases allergic reactions may occur.

**Interactions with other medication:** None known.

**Dosage:** In acute disorders daily, otherwise 3-1 times weekly 1 ampoule i.m., s.c., i.d., i.v.

**Package sizes:** Packs containing 5, 10, 50 and 100 ampoules of 1,1 ml. (9487)

#### Lymphomyosot®

Tablets

**Compositions:** 1 tabl. cont.:

Araneus diadematus D6 15,0 mg, Calcium phosphoricum D12 15,0 mg, Equisetum hiemale D4 15,0 mg, Ferrum iodatum D12 30,0 mg, Fumaria officinalis D4 15,0 mg, Gentiana lutea D5 15,0 mg, Geranium robertianum D4 30,0 mg, Juglans regia ssp. regia D3 15,0 mg, Levothyroxinum D12 15,0 mg, Myosotis arvensis D3 15,0 mg, Nasturtium officinale D4 30,0 mg, Natrium sulfuricum Trit. D4 15,0 mg, Pinus sylvestris D4 15,0 mg, Scrophularia nodosa D3 15,0 mg, Smilax D6 15,0 mg, Teucrium scorodonia D3 15,0 mg, Veronica officinalis D3 15,0 mg

Contains lactose! Please see package insert!

**Indications:** Lymphatism (tendency to hypertrophy of the lymphatic organs, tendency to formation of oedemas and to susceptibility to infection); glandular swelling; tonsillar hypertrophy and chronic tonsillitis. Special note: This remedy contains Lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product. If symptoms persist or worsen you should consult a doctor or healthcare professional. For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

**Contraindications:** Do not use in thyroid conditions without seeking medical advice.

**Side effects:** In very rare cases allergic reactions may occur.

**Interactions with other medication:** None known.

**Dosage:** In general, 3 tablets to be dissolved in the mouth 3 times daily.

**Package sizes:** Packs containing 50, 100 and 250 tablets. (9832)

#### Mucosa comp. (Mucosa compositum)

Solution for injection

**Compositions:** 1 ampoule of 2,2 ml (= 2,2g) contains:

Argentum nitricum D6 22,0 mg, Atropa bella-donna D10 22,0 mg, Ceanothus americanus D4 22,0 mg, Colibacillinum Nosode D28 22,0 mg, Hydrastis canadensis D4 22,0 mg, Kalium bichromicum D8 22,0 mg, Kreosotum D10 22,0 mg, Lachesis D10 22,0 mg, Mandragora e radice siccata D10 22,0 mg, Marsdenia cundurango D6 22,0 mg, Momordica balsamina D6 22,0 mg, Mucosa coli suis D8 22,0 mg, Mucosa ductus choledochi suis D8 22,0 mg, Mucosa duodeni suis D8 22,0 mg, Mucosa ilei suis D8 22,0 mg, Mucosa jejuni suis D8 22,0 mg, Mucosa nasalis suis D8 22,0 mg, Mucosa oculi suis D8 22,0 mg, Mucosa oesophagi suis D8 22,0 mg, Mucosa oris suis D8 22,0 mg, Mucosa pulmonis suis D8 22,0 mg, Mucosa pylori suis D8 22,0 mg, Mucosa recti suis D8 22,0 mg, Mucosa vesicae felleae suis D8 22,0 mg, Mucosa vesicae urinae suis D8 22,0 mg, Oxalis acetosella D6 22,0 mg, Pankreas suis D10 22,0 mg, Phosphorus D8 22,0 mg, Psychotria ipecacuanha D8 22,0 mg, Pulsatilla pratensis D6 22,0 mg, Semecarpus anacardium D6 22,0 mg, Strychnos nux-vomica D13 22,0 mg, Sulfur D8 22,0 mg, Sturium D8 22,0 mg, Ventriculus suis D8 22,0 mg, Veratrum album D4 22,0 mg, Natrium diethylxalacetatum D8 22,0 mg

**Indications:** Stimulation of the bodily defences in diseases of the mucous membranes and catarrhs of various types and localisations, e.g. in the alimentary canal with or without ulcers, in the region of the upper and lower parts of the respiratory tract, the urinary passages and the conjunctiva. Special note: If symptoms persist or worsen you should consult a doctor or healthcare professional. For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

**Contraindications:** None.

**Side effects:** Skin reactions may occur in individual cases, in which case use of the product should be discontinued.

**Interactions with other medication:** None known.

**Dosage:** In acute disorders daily, otherwise 3-1 times weekly 1 ampoule i.m., s.c., i.d., or if necessary i.v.

**Package sizes:** Packs containing 5, 10, 50 and 100 ampoules of 2,2 ml. (8526)

#### Mucosa comp.-Heel (Mucosa compositum Heel)

Tablets

**Compositions:** 1 tabl. cont.:

Argentum nitricum D6 1,0 mg, Atropa bella-donna D10 1,0 mg, Ceanothus americanus D4 1,0 mg, Hydrastis canadensis D6 1,0 mg, Kalium bichromicum D8 1,0 mg, Kreosotum D12 1,0 mg, Lachesis D10 1,0 mg, Mandragora e radice siccata D10 1,0 mg, Marsdenia cundurango D6 1,0 mg, Momordica balsamina D6 1,0 mg, Mucosa coli suis D8 1,0 mg, Mucosa ductus choledochi suis D8 1,0 mg, Mucosa duodeni suis D8 1,0 mg, Mucosa ilei suis D8 1,0 mg, Mucosa jejuni suis D8 1,0 mg, Mucosa nasalis suis D8 1,0 mg, Mucosa oculi suis D8 1,0 mg, Mucosa oesophagi suis D8 1,0 mg, Mucosa oris suis D8 1,0 mg, Mucosa pulmonis suis D8 1,0 mg, Mucosa pylori suis D8 1,0 mg, Mucosa recti suis D8 1,0 mg, Mucosa vesicae felleae suis D8 1,0 mg, Mucosa vesicae urinae suis D8 1,0 mg, Oxalis acetosella D6 1,0 mg, Pankreas suis D10 1,0 mg, Phosphorus D8 1,0 mg, Psychotria ipecacuanha D8 1,0 mg, Pulsatilla pratensis D6 1,0 mg, Semecarpus anacardium D6 1,0 mg, Strychnos nux-vomica D13 1,0 mg, Sulfur D8 1,0 mg, Ventriculus suis D8 1,0 mg, Veratrum album D6 1,0 mg

Contains lactose! Please see package insert!

**Indications:** Stimulation of the bodily defences in diseases of the mucous membranes and catarrhs of various types and localisations, e.g. in the alimentary canal with or without ulcers, in the region of the upper and lower parts of the respiratory tract, the urinary passages and the conjunctiva. Special note: This remedy contains Lactose. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product. For this product no clinical data on pregnancy and lactation are available. Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation. No adverse effects have so far been reported.

**Contraindications:** None.

**Side effects:** Skin reactions may occur in individual cases, in which case use of the product should be discontinued. Phenol allergy

**Interactions with other medication:** None known.

**Dosage:** Tablets: Standard dosage: Adults (and children 12 yrs. and older): 1 tablet to be dissolved in the mouth 3x daily. Acute or initial dosage: Adults (and children 12 yrs. and older): 1 tablet every ½ to 1 hr., up to 12x daily, and then continue with standard dosage.

**Package sizes:** Packs containing 50 and 250 tablets. (52268)

#### Oculoheel®

Eye drops

**Compositions:** 1 phial with 0,45 ml (=0,45 g) contains:

Cochlearia officinalis D5 110,7 mg, Echinacea D5 110,7 mg, Euphrasia D5 110,7 mg, Pilocarpus D5 110,7 mg

**Indications:** Irritation and inflammation of the conjunctiva and marginal eyelid due to external stimuli, overstraining, or allergic reactions; dry eyes. Special note: The phial may be used only once. The contents of the phial are sufficient for one application: i.e., for dripping once into the left and/or right eye as needed. Do not open the phial until just before application. Discard any phial which may still contain solution, and not use later. If symptoms persist or worsen you should consult a doctor or healthcare professional. For this product no clinical data on pregnancy and lactation are available.

Homeopathic dilutions of the substances present in this medicament are not known to be toxic during pregnancy and lactation.

No adverse effects have so far been reported.

**Contraindications:** Hypersensitivity to one of the active ingredients or excipients or to plants of the daisy family (Asteraceae).

**Side effects:** Hypersensitivity reactions may occur in individual cases.

**Interactions with other medication:** None known.

**Dosage:** In general, allow 1 drop to drip into the affected eye, 3 times daily.

**Package sizes:** Packages with 15 phials, each containing 0,45 ml of solution. (12142)

Le rogamos Introducir los textos específicos de su país





