

ASCEplus IRLV ;

[Instructions]

[Vial 1 | Lyophilized
Powder]




[Vial 2 | Diluent]



1. Remove the rubber stopper from the second vial.
2. Mix the two liquid solutions in the first vial with lyophilized powder.
3. Place the vial in a horizontal place and completely dissolve the first lyophilized powder.
4. Leave it alone for up to 3 to 5 minutes to remove air bubbles and use it.
5. Please use the mixture with lyophilized powder for up to 3 hours to ensure the safety of the product.

ASCEplus IRLV <ProcedurePeriod>

ASCEplus IRLV ; [Procedure Period]



- 3~4 weeks interval / 1~3 sessions.
- Sessions may be extended further than 3 session depending on Patient's condition.

ASCEplus IRLV < Protocol >

Protocol ; [With Device Treatment]**Step 1.** [Cleansing]

Cleanse the application area with warm water

**Step 2.** [Anesthesia – Optional]

Apply anesthesia cream and wait for 20~30 minutes and remove the cream.

**Step 3.** [RF Needling/Laser Treatment – Optional]

It is recommended to select and apply an energy device suitable for the patient's affected area condition. Irradiate each part of the patient's affected area (CO2 Laser, RF, RF needling etc.)

**Step 4.** [ASCEplus IRLV Application]

Apply the product on the vaginal area.
Massage into the skin

**Step 5.** [After Care]

After the treatment, it is recommended not to wash the area with water for 12 hours and to avoid using chemical products for 24 hours. It is also advised to refrain from vigorous activity on the same day.