

Glycine Powder



Clinical Applications

- Helps Maintain Healthy Skin, Joint, and Bones*
- Supports Brain Health*
- Supports Liver Health and Function*
- Supports Detoxification*
- Supports Mood and Sleep*

Glycine Powder offers the amino acid glycine in a pure powder form for flexible dosing and convenience. Although glycine is synthesized in the human body, supplemental amounts may be beneficial to help meet the body's many biological and metabolic demands for this amino acid. Glycine powder may be beneficial in supporting normal muscle tissue repair after a workout.

All Absolute Health Formulas Meet or Exceed cGMP Quality Standards

Discussion

Glycine is a conditionally essential amino acid used for structural protein synthesis, detoxification pathways, neurotransmitter function, brain health, sleep support, and healthy blood sugar levels. It is the smallest amino acid, with a side chain consisting of just a single hydrogen atom. Although glycine is synthesized from serine in the human body, endogenous synthesis has been shown to fall far short of meeting the many biological and metabolic demands for this amino acid in healthy adults.¹

Structural and Transport Proteins Glycine is a primary constituent of collagen, making up over one third of the total amino acids in this key structural component of blood vessels, skin, bones, cartilage, tendons, ligaments, and other connective tissue. A reaction between glycine and succinyl-CoA resulting in δ -aminolevulinic acid is the rate-limiting first step in the synthesis of the heme constituent of hemoglobin, myoglobin, and the cytochrome proteins.² Glycine is also required for nucleic acid synthesis. For these reasons, glycine supplementation may be beneficial for supporting protein synthesis, tissue repair, and wound healing.³ In rat models of tendonitis, a high-glycine diet resulted in reduced inflammation and improved tissue repair, including higher concentrations of hydroxyproline and glycosaminoglycans.⁴ Rats undergoing laparotomy who were supplemented with a 1% glycine diet (plus arginine) had greater overall nitrogen retention, higher ratios of type III/type I collagen, and more hydroxyproline than unsupplemented controls, mostly due to increased collagen synthesis in the wounds.⁵ Mouse models of cancer cachexia indicate that glycine (injected subcutaneously) reduced loss of muscle and fat mass, and blunted increases in inflammatory markers. Similar improvements were not seen with isonitrogenous doses of alanine or citrulline, suggesting that the beneficial effects on tissue sparing are glycine specific.⁶

Detoxification, Glutathione Synthesis, and Liver Function Glycine is one of the amino acids that make up the glutathione tripeptide (a.k.a. gamma-glutamyl-cysteinyl-glycine). Reductions in glycine concentration from protein malnutrition, sepsis, or increased glycine oxidation due to diabetes or high glucagon levels may impair synthesis of adequate amounts of glutathione (GSH). Glycine availability is a determining factor in erythrocyte GSH synthesis in burn patients and children recovering from malnutrition; supplemental glycine improves hepatic GSH concentrations in protein-deficient rats.⁷ Glutathione is the body's master antioxidant and is also a critical compound for enabling healthy detoxification processes in the liver. Beyond glycine's role as a constituent of glutathione, however, glycine itself plays a role in phase II biotransformation and conjugation of xenobiotics and endogenously produced toxins.^{10,11} Impaired glycine conjugation may result in a buildup of compounds that adversely affect mitochondrial function, and impaired mitochondrial dysfunction is associated with a host of metabolic and neurological impairments.¹²

Glycine may be especially supportive of liver health after alcohol-induced damage and other forms of liver injury.¹³ Compared to control rats on a standard diet, those on a glycine-enriched diet experienced 30% faster recovery in aspartate aminotransferase (AST) and alanine aminotransferase (ALT) levels. After one-week post-ethanol-induced damage, hepatic steatosis was reduced significantly more in the glycine group than in the controls, and the glycine group also had significantly diminished numbers of infiltrating leukocytes and necrotic cells compared to controls.¹⁴ Liver injury due to ischemia/reperfusion (I/R), partial resection, or transplant may also be reduced by oral glycine supplementation. Studies in rats that underwent partial liver resection with I/R injury show that oral glycine supplementation led to significant improvements in survival rate and recovery rate of the remnant liver weight compared to untreated rats.¹⁵ Study authors concluded that glycine may be clinically beneficial to the prognosis of patients with liver resection, and it has also been shown to be cytoprotective for other tissues, including the kidneys and small intestine.¹⁵⁻¹⁸

Function as an Inhibitory Neurotransmitter Glycine is a well-known inhibitory neurotransmitter and is often included in combination products to combat stress and anxiety. The binding of glycine to its neuronal receptor results in an influx of chloride ions, leading to membrane hyperpolarization and inhibition of the response to excitatory neurotransmitters.¹⁷ Alterations in glycine receptor genes have been implicated in the pathology of autism, generalized epilepsy, and amyotrophic lateral sclerosis (ALS), while autoantibodies against subunits of the glycine receptor have been found in patients with progressive encephalomyelitis, including rigidity and myoclonus.²⁰ Owing to its inhibitory function, glycine may be helpful for improving sleep quality, which, in itself, may be beneficial for supporting healthy moods and responses to psychological stress. In a randomized crossover study, a single bolus of glycine (3 g) given one hour before bed to subjects who regularly experienced unsatisfactory sleep but were otherwise healthy resulted in subjective and objective improvements of sleep parameters. Compared to placebo, glycine supplementation led to improved subjective assessments of difficulty in getting to sleep, the time it took to fall asleep, and overall sleep efficiency (time in bed versus time asleep). Direct measurements showed that compared to placebo, glycine reduced sleep onset latency and the onset of slow wave sleep. Latency to rapid eye movement (REM) sleep remained unchanged, as did the total sleep architecture—the ratios of time spent in each sleep stage. Nevertheless, assessments of daytime sleepiness, cognitive function and reaction time the day after glycine ingestion showed that glycine supplementation slightly improved these parameters compared to placebo.²¹ Other studies support the role of glycine in improving sleep quality as well as "liveliness and peppiness," and "clear-headedness" upon awakening.²² Additionally, daytime ingestion of glycine does not lead to acute sleepiness, so individuals taking glycine for other reasons should not be at risk for this potentiality. Glycine is required along with glutamate to open ion channels on neuronal N-methyl-d-aspartate (NMDA) receptors.²³ Glycine supplementation and increasing brain glycine levels may be instrumental in improving treatment-resistant schizophrenia, particularly in patients with low baseline serum glycine levels. Randomized, placebo-controlled, crossover studies show that glycine supplementation added to prior antipsychotic treatment results in significant reductions in negative symptoms, with improvements in depressive and cognitive symptoms.²⁴⁻²⁶ Other calming and inhibitory effects of glycine in the central nervous system include reversing hyperactivity and schizophrenia-like psychosis induced by phencyclidine (PCP).²⁷

Absolute Health
7350 SW 60th Ave., Suite 2
Ocala, FL 34476
www.AbsoluteHealthOcala.com

Glycine Powder



Supplement Facts

Serving Size 3 grams (approx. one scoop)

Servings Per Container 60

Amount Per Serving	% Daily Value
Glycine	3g *

*Daily Value not established.

GLY180-2

Directions

As a dietary supplement take 3 grams (approx. 1/2 teaspoon) with water or any other beverage per day, or as directed by your health care practitioner. Glycine powder can be used post-workout for recovery and glycogen resynthesis, especially for those following a low carb diet and who prefer to avoid post-workout carbohydrates.

Consult your healthcare practitioner prior to use. Individuals taking medication should discuss potential interactions with their healthcare practitioner.

Does Not Contain

Gluten, soy, animal, or dairy products from genetically modified organisms (GMOs)



References

- Meléndez-Hevia E, De Paz-Lugo P, Cornish-Bowden A, Cárdenas ML. A weak link in metabolism: the metabolic capacity for glycine biosynthesis does not satisfy the need for collagen synthesis. *J Biosci.* 2009 Dec;34(6):853-72.
- Ajioka RS, Phillips JD, Kushner JP. Biosynthesis of heme in mammals. *Biochim Biophys Acta.* 2006 Jul;1763(7):723-36.
- Wang W, Wu Z, Dai Z, Yang Y, Wang J, Wu G. Glycine metabolism in animals and humans: implications for nutrition and health. *Amino Acids.* 2013 Sep;45(3):463-77.
- Vieira CP, De Oliveira LP, Da Ré Guerra F, et al. Glycine improves biochemical and biomechanical properties following inflammation of the achilles tendon. *Anat Rec (Hoboken).* 2015 Mar;298(3):538-45.
- Chyun JH, Griminger P. Improvement of nitrogen retention by arginine and glycine supplementation and its relation to collagen synthesis in traumatized mature and aged rats. *J Nutr.* 1984 Sep;114(9):1697-704.
- Ham DJ, Murphy KT, Chee A, Lynch GS, Koopman R. Glycine administration attenuates skeletal muscle wasting in a mouse model of cancer cachexia. *Clin Nutr.* 2014 Jun;33(3):448-58.
- Wu G1, Fang YZ, Yang S, Lupton JR, Turner ND. Glutathione metabolism and its implications for health. *J Nutr.* 2004 Mar;134(3):489-92.
- Sekhar RV, Patel SG, Guthikonda AP, et al. Deficient synthesis of glutathione underlies oxidative stress in aging and can be corrected by dietary cysteine and glycine supplementation. *The American Journal of Clinical Nutrition.* 2011;94(3):847-853.
- Sekhar RV, McKay SV, Patel SG, et al. Glutathione Synthesis Is Diminished in Patients With Uncontrolled Diabetes and Restored by Dietary Supplementation With Cysteine and Glycine. *Diabetes Care.* 2011;34(1):162-167.
- van der Sluis R, Badenhorst CP, Erasmus E, et al. Conservation of the coding regions of the glycine N-acyltransferase gene further suggests that glycine conjugation is an essential detoxification pathway. *Gene.* 2015 Oct 15;571(1):126-34.
- Badenhorst CP, van der Sluis R, Erasmus E, van Dijk AA. Glycine conjugation: importance in metabolism, the role of glycine N-acyltransferase, and factors that influence interindividual variation. *Expert Opin Drug Metab Toxicol.* 2013 Sep;9(9):1139-53.
- Knights KM1, Sykes MJ, Miners JO. Amino acid conjugation: contribution to the metabolism and toxicity of xenobiotic carboxylic acids. *Expert Opin Drug Metab Toxicol.* 2007 Apr;3(2):159-68.
- Yamashina S, Ikejima K, Enomoto N, Takei Y, Sato N. Glycine as a therapeutic immunonutrient for alcoholic liver disease. *Alcohol Clin Exp Res.* 2005 Nov;29(11 Suppl):162S-5S.
- Yin M, Ikejima K, Arteel GE, et al. Glycine accelerates recovery from alcohol-induced liver injury. *J Pharmacol Exp Ther.* 1998 Aug;286(2):1014-9.
- Ito K, Ozasa H, Noda Y, et al. Effect of non-essential amino acid glycine administration on the liver regeneration of partially hepatectomized rats with hepatic ischemia/reperfusion injury. *Clin Nutr.* 2008 Oct;27(5):773-80.
- Habib MM, Hodgson HJ, Davidson BR. The role of glycine in hepatic ischemiareperfusion injury. *Curr Pharm Des.* 2006;12(23):2953-67.
- Petrat F, Boengler K, Schulz R, de Groot H. Glycine, a simple physiological compound protecting by yet puzzling mechanism(s) against ischaemia-reperfusion injury: current knowledge. *British Journal of Pharmacology.* 2012;165(7):2059-2072.
- Zhong Z, Wheeler MD, Li X, et al. L-Glycine: a novel antiinflammatory, immunomodulatory, and cytoprotective agent. *Curr Opin Clin Nutr Metab Care.* 2003 Mar;6(2):229-40.
- Hofmann AF. The function of bile salts in fat absorption. The solvent properties of dilute micellar solutions of conjugated bile salts. *Biochemical Journal.* 1963;89(1):57-68.
- Dutertre S, Becker C-M, Betz H. Inhibitory Glycine Receptors: An Update. *The Journal of Biological Chemistry.* 2012;287(48):40216-40223.
- Yamadera W, Inagawa K, Chiba S, et al. Glycine ingestion improves subjective sleep quality in human volunteers, correlating with polysomnographic changes. *Sleep and Biological Rhythms.* 2007; 5: 126-131.
- Inagawa K, Hiraoka T, Kohda T, et al. Subjective effects of glycine ingestion before bedtime on sleep quality. *Sleep and Biological Rhythms.* 2006; 4: 75-77.
- Cummings KA, Popescu GK. Glycine-dependent activation of NMDA receptors. *The Journal of General Physiology.* 2015;145(6):513-527.
- Heresco-Levy U, Javitt DC, Ermilov M, et al. Double-blind, placebo-controlled, crossover trial of glycine adjuvant therapy for treatment-resistant schizophrenia. *Br J Psychiatry.* 1996 Nov;169(5):610-7.
- Yang CR, Svensson KA. Allosteric modulation of NMDA receptor via elevation of brain glycine and D-serine: the therapeutic potentials for schizophrenia. *Pharmacol Ther.* 2008 Dec;120(3):317-32.
- Heresco-Levy U, Javitt DC, Ermilov M, et al. Efficacy of high-dose glycine in the treatment of enduring negative symptoms of schizophrenia. *Arch Gen Psychiatry.* 1999 Jan;56(1):29-36.
- Javitt DC, Sershen H, Hashim A, Lajtha A. Reversal of phencyclidine-induced hyperactivity by glycine and the glycine uptake inhibitor glycyldodecylamide. *Neuropsychopharmacology.* 1997 Sep;17(3):202-4.

*These statements have not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure, or prevent any disease.