

Long-term use:

Do not use anticoagulants as permanent baits (e.g. for prevention of rodent infestation or to detect rodent activity). Using this product should eliminate rodents within 35 days. In case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), advice should be sought from the product supplier or a pest control service.

Medical advice:

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine. Antidote: Vitamin K1 administered by medical/veterinary personnel only.

In case of: Dermal exposure, wash skin with water and then with water and soap.

Eye exposure, rinse eyes with eye-rinse liquid or water, keep eyes lids open at least 10 minutes.

Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label. Contact a veterinary surgeon in case of ingestion by a pet. UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice. In Ireland, for information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-8092166), retain the label for reference.

Precautions:

Get medical advice/attention if you feel unwell. **IF SWALLOWED:** Immediately call a POISON CENTRE or doctor/physician. Keep out of reach of children. Read label before use. Keep /Store away from food, drink and animal feeding stuffs. Do not get on skin. Do not eat, drink or smoke when using this product. Store locked up in a closed container. Dispose of contents/container in accordance with national regulations.

Warning:

May cause damage to organs (Blood) through prolonged or repeated exposure.

To avoid risks to human health and the environment, comply with the instructions for use.

Other information:

Hazardous to wildlife. Because of their delayed mode of action, anticoagulant rodenticides may take from 4 to 10 days to be effective after consumption of the bait. This product contains a bittering agent and a dye. Bait stations must be labelled with the following information: 'do not move or open', 'contains a rodenticide', 'product name or authorisation number', 'active substance', 'in case of incident call a poison centre - UK: 111 / Ireland (01) 809 2166'.



UFI: AJ10-003E-500J-AQ1M / Batch & Expiry: See base.

Contact details:

Pelsis Ltd, Sterling House, Grimbald Crag Close,
Knaresborough HG5 8PJ
Tel: +44 (0) 800 988 5359

Authorisation holder: UK - PelGar International
Ltd., Alton, Hampshire, GU34 2QR. UK.
Ireland - PelGar International Ltd., 18 rue des
Remparts d'Ainay 69002 Lyon, France.

About Pest-Stop

Pest-Stop supplies high quality products for the control of all home and garden pests. See our full range at www.pest-stop.co.uk

PSWB04

35449

87822028

PEST-STOP

Super Rat & Mouse Killer Blocks

wax block RODENTICIDE

- ✓ For use indoors and outdoors around buildings
- ✓ Fast acting and highly effective
- ✓ For use in all weather conditions

TARGET



READY-TO-USE BAIT TO BE USED IN TAMPER RESISTANT BAIT STATIONS FOR THE CONTROL OF RATS AND MICE IN AND AROUND BUILDINGS.

15x20ge

UK-2016-0986-0002 / IE/BPA 70571-001

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it. Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered. Consider preventative control measures (block holes, remove potential food and drink as far as possible) to improve product intake and reduce the likelihood of reinvasion.

Directions for use:

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve. Use only in tamper resistant bait stations. Bait stations should be placed in the immediate vicinity where rodent activity has been observed (e.g. rodent runs, nesting sites, holes, burrows etc.). When used for the control of mice, bait stations must only be placed indoors.

Target	Infestation	Bait points	Advice
Rats (<i>Rattus norvegicus</i>)	Low	3-5 blocks in bait stations every 10 metres	The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
	High	3-5 blocks in bait stations every 5 metres	
Mice (<i>Mus musculus</i>)	Low	1 block in bait stations every 5 metres	The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.
	High	1 block in bait stations every 2 metres	

Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt. Where possible, bait stations must be fixed to the ground or other structures. Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals. Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these. Place the bait stations in areas not liable to flooding. Do not place bait stations near water drainage systems where they can come into contact with water. When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product. Remove the remaining bait or the bait stations at the end of the treatment period. Search for and remove dead rodents during treatment, at least as often as when bait stations are inspected.

Storage and disposal:

Store in a dry, cool and well-ventilated place. Keep only in original container. Keep the container closed and away from direct sunlight. Store in places prevented from the access of children, birds, pets and farm animals. At the end of the treatment, dispose of uneaten bait and the packaging in accordance with local requirements. In the UK waste bait should be double bagged in plastic bags and disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets, or taken to a civic amenity site. For information on civic amenity sites contact the local authority. In Ireland dispose of uneaten bait and the packaging in accordance with EPA requirements for the disposal of hazardous waste. Use of gloves is recommended. Dispose of dead rodents in accordance with local requirements. Poisoned rodents should be double-bagged using plastic bags and either disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets, or collected by a specialist waste contractor or (UK only - the local authority). Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing of them.

For general public use. Ready-to-use bait. For the protection of public health, stored products and materials. Use biocides safely and sustainably. It is illegal to use this product for uses or in a manner other than that prescribed on this label. Contains bromadiolone [CAS 28772-56-7] 0.0025% w/w and denatonium benzoate 0.001% w/w.