

S A SOUTHERN
5 0 AG®

CHELATED LIQUID IRON

**FOR THE CORRECTION AND PREVENTION OF FOLIAGE
YELLOWING DUE TO IRON DEFICIENCY OF FLOWERS,
SHRUBS, TREES, VEGETABLES AND LAWNS.**

MINIMUM GUARANTEED ANALYSIS:

Soluble Iron (Fe) 5.00%
3.50% Chelated Iron as (Fe)
Combined Sulfur as (S)..... 3.40%

DERIVED FROM: Iron Sulfate, Iron Lignin Sulfonate.

Chelating agent is Lignin Sulfonate
Guaranteed by: Southern Agricultural Insecticides, Inc.
P.O. Box 218 Palmetto, FL 34220

CAUTION:

KEEP OUT OF REACH OF CHILDREN

Avoid contact with skin and eyes.
Avoid breathing spray mist.
Do not take internally.



Southern Agricultural Insecticides, Inc.

Palmetto, FL
F672

Hendersonville, NC
J4

Boone, NC

Net Wt.: 9lbs. 13ozs.

Net Contents: 1 Gallon (3.785 liters)

USE PRECAUTIONS

THIS PRODUCT WILL SEVERELY STAIN CONCRETE, MASONRY AND PAINTED SURFACES.

Be extremely careful not to allow spray or runoff to contact with these materials. Do not mix with other fertilizers, fungicides or insecticides unless compatibility is known.

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin and eyes.

Avoid breathing spray mist. Do not take internally.

FIRST AID

IF SWALLOWED: Drink 2 glasses of water or milk and seek medical attention.

IF IN EYES: Flush eyes thoroughly with a large volume of water. If irritation persists seek medical attention.

IF ON SKIN: Rinse affected area with clear water and then wash thoroughly with soap and water.



LOT NO

0417067

SA-50 Brand Chelated Liquid Iron is intended for use in the correction and prevention of foliage yellowing due to iron deficiency of flowers, shrubs, trees, vegetables and lawns.

DIRECTIONS FOR USE

Shake well before using.

VEGETABLES AND FIELD CROPS:

Full volume conventional sprayers: Use 1-2 quarts per 100 gallons of water. Low Volume Applications: Use 1-2 quarts per acre in at least 5 gallons of water. For lower application volumes and on young sensitive crops, use the 1 quart rate. Repeat applications at 7-10 day intervals as needed.

CITRUS, APPLES

and other TREE CROPS:

Use 2-4 quarts per 500 gallons of dilute spray.

For concentrate sprays, increase the amounts used in proportion to the dilution.

ORNAMENTALS and young or sensitive crops:

Use 1-2 pints per 100 gallons of water.

LAWNS and TURF:

Use 2.5 - 5 gallons per acre (1 pint per 1,000 sq. ft.) Apply in a sufficient volume of water to thoroughly wet the foliage and runoff in the root zone.

Note: Iron deficiency in many plants is an indication of high pH or alkaline soil. If yellowing persists or returns rapidly after treatment, soil pH should be checked and adjusted if necessary.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions and precautions given herewith.