

S A M H S A

Opioid Overdose

TOOLKIT:

Five Essential Steps
for First Responders



TABLE OF CONTENTS

FIVE ESSENTIAL STEPS FOR FIRST RESPONDERS

STEP 1: CALL FOR HELP (DIAL 911)	3
STEP 2: CHECK FOR SIGNS OF OPIOID OVERDOSE	3
STEP 3: SUPPORT THE PERSON'S BREATHING	4
STEP 4: ADMINISTER NALOXONE	4
STEP 5: MONITOR THE PERSON'S RESPONSE	5
SUMMARY: DO'S AND DON'TS IN RESPONDING TO OPIOID OVERDOSE	5
ACKNOWLEDGMENTS, ETC.	7
■ Acknowledgments	
■ Disclaimer	
■ Public Domain Notice	
■ Electronic Access and Copies of Publication	
■ Recommended Citation	
■ Originating Office	

Also see the other components of this Toolkit:

- ★ Facts for Community Members
- ★ Information for Prescribers
- ★ Safety Advice for Patients & Family Members
- ★ Recovering from Opioid Overdose:
Resources for Overdose Survivors & Family Members

FIVE ESSENTIAL STEPS FOR FIRST RESPONDERS

Overdose is common among persons who use illicit opioids such as heroin and among those who misuse medications prescribed for pain, such as oxycodone, hydrocodone, and morphine. The incidence of opioid overdose is rising nationwide. For example, between 2001 and 2010, the number of poisoning deaths in the United States nearly doubled, largely because of overdoses involving prescription opioid analgesics [1]. This increase coincided with a nearly fourfold increase in the use of prescribed opioids for the treatment of pain [2].

To address the problem, emergency medical personnel, health care professionals, and patients increasingly are being trained in the use of the opioid antagonist naloxone hydrochloride (naloxone or Narcan), which is the treatment of choice to reverse the potentially fatal respiratory depression caused by opioid overdose. (Note that naloxone has no effect on non-opioid overdoses, such as those involving cocaine, benzodiazepines, or alcohol [3].)

Based on current scientific evidence and extensive experience, the steps outlined below are recommended to reduce the number of deaths resulting from opioid overdoses [4-9].

STEP 1: CALL FOR HELP (DIAL 911)

AN OPIOID OVERDOSE NEEDS IMMEDIATE MEDICAL ATTENTION.

An essential step is to get someone with medical expertise to see the patient as soon as possible, so if no EMS or other trained personnel are on the scene, dial 911 immediately. All you have to say is: "Someone is not breathing." Be sure to give a clear address and/or description of your location.

STEP 2: CHECK FOR SIGNS OF OPIOID OVERDOSE

Signs of **OVERDOSE**, which often results in death if not treated, include [3]:

- Face is extremely pale and/or clammy to the touch
- Body is limp
- Fingernails or lips have a blue or purple cast
- The patient is vomiting or making gurgling noises
- He or she cannot be awakened from sleep or is unable to speak
- Breathing is very slow or stopped
- Heartbeat is very slow or stopped

Signs of **OVERMEDICATION**, which may progress to overdose, include [3]:

- Unusual sleepiness or drowsiness
- Mental confusion, slurred speech, intoxicated behavior
- Slow or shallow breathing
- Pinpoint pupils
- Slow heartbeat, low blood pressure
- Difficulty waking the person from sleep

Because opioids depress respiratory function and breathing, one telltale sign of a person in a critical medical state is the "death rattle." If a person emits a "death rattle" — an exhaled breath with a very distinct, labored sound coming from the throat — emergency resuscitation will be necessary immediately, as it almost always is a sign that the individual is near death [8].

FIVE ESSENTIAL STEPS FOR FIRST RESPONDERS

STEP 3: SUPPORT THE PERSON'S BREATHING

Ideally, individuals who are experiencing opioid overdose should be ventilated with 100% oxygen before naloxone is administered so as to reduce the risk of acute lung injury [5,7]. In situations where 100% oxygen is not available, rescue breathing can be very effective in supporting respiration [5]. Rescue breathing involves the following steps:

- Be sure the person's airway is clear (check that nothing inside the person's mouth or throat is blocking the airway).
- Place one hand on the person's chin, tilt the head back and pinch the nose closed.
- Place your mouth over the person's mouth to make a seal and give 2 slow breaths.
- The person's chest should rise (but not the stomach).
- Follow up with one breath every 5 seconds.

STEP 4: ADMINISTER NALOXONE

Naloxone (Narcan) should be administered to any person who shows signs of opioid overdose, or when overdose is suspected [7]. Naloxone injection is approved by the FDA and has been used for decades by emergency medical services (EMS) personnel to reverse opioid overdose and resuscitate individuals who have overdosed on opioids.

Naloxone can be given by intramuscular or intravenous injection every 2 to 3 minutes [7-9]. The most rapid onset of action is achieved by intravenous administration, which is recommended in emergency situations [8]. The dose should be titrated to the smallest effective dose that maintains spontaneous normal respiratory drive.

Opioid-naive patients may be given starting doses of up to 2 mg without concern for triggering withdrawal symptoms [5-7, 9].

The intramuscular route of administration may be more suitable for patients with a history of opioid dependence because it provides a slower onset of action and a prolonged duration of effect, which may minimize rapid onset of withdrawal symptoms [5-7].

DURATION OF EFFECT. The duration of effect of naloxone is 30 to 90 minutes, and patients should be observed after this time frame for the return of overdose symptoms [7-9]. The goal of naloxone therapy should be to restore adequate spontaneous breathing, but not necessarily complete arousal [7].

More than one dose of naloxone may be needed to revive someone who is overdosing. Patients who have taken longer-acting opioids may require further intravenous bolus doses or an infusion of naloxone [7].

Comfort the person being treated, as withdrawal triggered by naloxone can feel unpleasant. As a result, some persons become agitated or combative when this happens and need help to remain calm.

SAFETY OF NALOXONE. The safety profile of naloxone is remarkably high, especially when used in low doses and titrated to effect [5-8]. When given to individuals who are not opioid-intoxicated or opioid-dependent, naloxone produces no clinical effects, even at high doses. Moreover, while rapid opioid withdrawal in tolerant patients may be unpleasant, it is not life-threatening.

Naloxone can safely be used to manage opioid overdose in pregnant women. The lowest dose to maintain spontaneous respiratory drive should be used to avoid triggering acute opioid withdrawal, which may cause fetal distress [7].

FIVE ESSENTIAL STEPS FOR FIRST RESPONDERS

STEP 5: MONITOR THE PERSON'S RESPONSE

All patients should be monitored for recurrence of signs and symptoms of opioid toxicity for at least 4 hours from the last dose of naloxone or discontinuation of the naloxone infusion. Patients who have overdosed on long-acting opioids should have more prolonged monitoring [5-7].

Most patients respond by returning to spontaneous breathing, with minimal withdrawal symptoms [7]. The response generally occurs within 3 to 5 minutes of naloxone administration. (Rescue breathing should continue while waiting for the naloxone to take effect. [5-7])

Naloxone will continue to work for 30 to 90 minutes, but after that time, overdose symptoms may return [8,9]. Therefore, it is essential to get the person to an emergency department or other source of medical care as quickly as possible, even if he or she revives after the initial dose of naloxone and seems to feel better.

SIGNS OF OPIOID WITHDRAWAL. The signs and symptoms of opioid withdrawal in an individual who is physically dependent on opioids may include, but are not limited to, the following: body aches, diarrhea, tachycardia, fever, runny nose, sneezing, piloerection, sweating, yawning, nausea or vomiting, nervousness, restlessness or irritability, shivering or trembling, abdominal cramps, weakness, and increased blood pressure. In the neonate, opioid withdrawal may also include: convulsions, excessive crying, and hyperactive reflexes [8].

NALOXONE-RESISTANT PATIENTS. If a patient does not respond to naloxone, an alternative explanation for the clinical symptoms should be considered. The most likely explanation is that the person is not overdosing on an opioid but rather some other substance or may even be experiencing a non-overdose medical emergency. A possible explanation to consider is whether the individual has overdosed on buprenorphine, a long-acting opioid partial agonist. Because buprenorphine has a higher affinity for the opioid receptors than do other opioids, naloxone may not be effective at reversing the effects of buprenorphine-induced opioid overdose [9.] In all cases, support of ventilation, oxygenation, and blood pressure should be sufficient to prevent the complications of opioid overdose and should be given priority if the response to naloxone is not prompt.

SUMMARY:

Do's and Don'ts in Responding to Opioid Overdose

- **DO** support the person's breathing by administering oxygen or performing rescue breathing.
- **DO** administer naloxone.
- **DO** put the person in the "recovery position" on their side, if he or she is breathing independently.
- **DO** stay with the person and keep him/her warm.
- **DON'T** slap or try to forcefully stimulate the person — it will only cause further injury. If you are unable to wake the person by shouting, rubbing your knuckles on the sternum (center of the chest or rib cage), or light pinching, he or she may be unconscious.
- **DON'T** put the person into a cold bath or shower. This increases the risk of falling, drowning or going into shock.
- **DON'T** inject the person with any substance (salt water, milk, "speed," heroin, etc.). The only safe and appropriate treatment is naloxone.
- **DON'T** try to make the person vomit drugs that he or she may have swallowed. Choking or inhaling vomit into the lungs can cause a fatal injury.

NOTE: All naloxone products have an expiration date, so it is important to check the expiration date and obtain replacement naloxone as needed.

FIVE ESSENTIAL STEPS FOR FIRST RESPONDERS

REFERENCES

1. Centers for Disease Control and Prevention (CDC), National Center for Health Statistics. CDC WONDER Online Database, 2012.
2. Harvard Medical School. Painkillers fuel growth in drug addiction; Opioid overdoses now kill more people than cocaine or heroin. *Harvard Ment Hlth Let.* 2011;27(7):4–5.
3. Hardman E, Limbird LE, eds. *Goodman and Gilman's the Pharmacologic Basis of Therapeutics*, 11th ed. New York, NY: McGraw-Hill, 2006, pp. 576–578.
4. Centers for Disease Control and Prevention (CDC). Community-based opioid overdose prevention programs providing naloxone — United States, 2010. *MMWR Morb Mortal Wkly Rep.* 2012;61(6):101–105.
5. Beletsky L, Rich JD, Walley AY. Prevention of fatal opioid overdose. *JAMA.* 2012 Nov 14;308(18):1863–1864.
6. Coffin PO, Sullivan SD. Cost-effectiveness of distributing naloxone to heroin users for lay overdose reversal. *Ann Int Med.* 2013;158:1–9.
7. BMJ Evidence Centre. Treatment of opioid overdose with naloxone. *British Medical Journal.* Updated October 23, 2012. [Accessed March 24, 2013, at www.bmjjournals.com]
8. Rx List [Accessed March 24, 2013, at www.rxlist.com]
9. Drugs.com [Accessed March 24, 2013, at www.drugs.com]

Acknowledgments

This publication was prepared for the Substance Abuse and Mental Health Services Administration (SAMHSA) by the Association of State and Territorial Health Officials, in cooperation with Public Health Research Solutions, under contract number 10-233-00100 with SAMHSA, U.S. Department of Health and Human Services (HHS). LCDR Brandon Johnson, M.B.A., served as the Government Project Officer.

Disclaimer

The views, opinions, and content of this publication are those of the authors and do not necessarily reflect the views, opinions, or policies of SAMHSA or HHS.

Public Domain Notice

All materials appearing in this volume except those taken directly from copyrighted sources are in the public domain and may be reproduced or copied without permission from SAMHSA or the authors. Citation of the source is appreciated. However, this publication may not be reproduced or distributed for a fee without the specific, written authorization of the Office of Communications, SAMHSA, HHS.

Electronic Access and Copies of Publication

This publication may be ordered from SAMHSA's Publications Ordering Web page at www.store.samhsa.gov. Or, please call SAMHSA at 1-877-SAMHSA-7 (1-877-726-4727) (English and Español).

Recommended Citation

Substance Abuse and Mental Health Services Administration. SAMHSA Opioid Overdose Prevention Toolkit: Five Essential Steps for First Responders. HHS Publication No. (SMA) 13-4742. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2013.

Originating Office

Division of Pharmacologic Therapies, Center for Substance Abuse Treatment, Substance Abuse and Mental Health Services Administration, 1 Choke Cherry Road, Rockville, MD 20857.



HHS Publication No. (SMA) 13-4742
Printed 2013