



Teaming confidence with nature since 1922



Wildflower Information Product Sheet

TEXAS REGIONAL POLLINATOR WILDFLOWER MIXTURE

THESE NATIVE SEEDS ARE THE FOUNDATION FOR YOUR BEE FRIENDLY GARDEN FIND OUT MORE AT:

WWW.BEEFRIENDLYGARDENING.ORG

TEXAS REGIONAL POLLINATOR WILDFLOWER MIXTURE					
States: Oklahoma, Texas					
Botanical Name	Common Name	Life Cycle	Flower Color	Bloom Period	Height inches
<i>Rudbeckia hirta</i>	Black Eyed Susan	A/P	Y	S	12-36
<i>Ratibida columnifera</i>	Prairie Coneflower	B/P	Y	S-F	12-36
<i>Coreopsis tinctoria</i>	Plains Coreopsis	A	Y-R	S-F	12-36
<i>Rudbeckia amplexicaulis</i>	Clasping Coneflower	A	Y	S-F	18-30
<i>Linum lewisii</i>	Blue Flax	P	B	S	18-30
<i>Dalea purpurea</i>	Purple Prairie Clover	P	Pu	Sp	12-36
<i>Gaillardia pulchella</i>	Indian Blanket	A	R/Y	S	12-24
<i>Symphyotrichum laeve</i>	Smooth Blue Aster	P	B	F	24-48
<i>Echinacea purpurea</i>	Purple Coneflower	P	Pu	S	24-36
<i>Salvia coccinea</i>	Scarlet Sage	P	R/Y	S-F	18-24
<i>Asclepias tuberosa</i>	Butterfly Weed	P	O	Sp-S	24-36
Approximate seeds per ounce: 32,700					
Life Cycle: P=Perennial, A=Annual, B=Biannual					
Flower Color: B=Blue, Pu=Purple, Pk=Pink, O=Orange, R=Red, Y=Yellow					
Bloom Period: Sp=Spring, S=Summer, F=Fall					
Coverage: 1 oz: 225 sq. ft. 4 oz: 900 sq. ft. 1 lb: 3,600 sq. ft.					

PLANTING INSTRUCTIONS:

Your plants will grow best in full sun, can tolerate a few hours of shade, and prefer moist, well- drained soil containing loam, gravel, or sand. The best time to sow seeds in most places is when the fall wet season starts, but seeds can be sown in winter or early spring as well. Sow seeds directly onto bare soil (remove vegetation/ mulch if necessary) and firmly press seeds into soil (but do not bury them more than 1/8 inch).

For more information scan the below QR code for a copy of our Establishment and Care of Wildflowers technical guide sheet



Serving California Since 1922

9180. SAN FERNANDO RD - P.O. BOX 1579 - SUN VALLEY, CALIFORNIA 91353- (213) 626-9668 - FAX (213) 626-4920
www.stoverseed.com