

NEURALGIA – JOINT & MUSCULAR PAIN

Patient Last Name _____ First name _____ DOB ____/____/____
Street Address _____ Phone ____-____-____
City, State, Zip _____
Drug Allergies _____

Extra Oral Neuropathic

- Guaifenesin, Ketoprofen, Menthol (10,10,0.1%) Anhydrous PLO Gel
- Guaifenesin, Ketoprofen, Menthol, Lidocaine HCL (10,10,0.1,1%) Anhydrous PLO Gel
- Guaifenesin, Ketamine, Ketoprofen (10,10,10%) Anhydrous PLO Gel
- Amitriptyline, Cyclobenzaprine (2,2%) Anhydrous PLO Gel
- Amitriptyline, Cyclobenzaprine, Lidocaine HCL (2,2,1%) Anhydrous PLO Gel
- Amitriptyline (4%) Anhydrous PLO Gel
- Amitriptyline, Lidocaine HCL (4,1%) Anhydrous PLO Gel
- Ibuprofen, Ketoprofen (2.5,10%) Anhydrous PLO Gel
- Ibuprofen, Ketoprofen, Lidocaine HCL (2.5,10,1%) Anhydrous PLO Gel
- Ketamine, Ketoprofen (4, 10%) Anhydrous PLO Gel
- Ketamine (4%) Anhydrous PLO Gel
- Gabapentin, Ketamine (4,4%) Anhydrous PLO Gel
- Gabapentin, Ketamine, Lidocaine HCL (4,4,1%) Anhydrous PLO Gel
- Amitriptyline, Ketamine, Gabapentin (4,4,4%) Anhydrous PLO Gel
- Amitriptyline, Carbamazepine, Gabapentin, Ketamine, Ketoprofen, Lidocaine (4,4,4,4,10,1%) Anhydrous PLO Gel

MD Pump 15g _____ 30g _____ 60g _____ 120g _____ 240g _____ Other _____ g

*Sig: Apply 1 pump (0.5ml) to affected area of skin 2-4 times daily, as needed.
Moisten skin with warm water & massage compound into skin for 2 minutes.*

Other Instructions: _____

Intra Oral Neuropathic Standard

- Carbamazepine, Gabapentin, Ketamine, Ketoprofen, Lidocaine HCL, Pregabalin (4,4,4,4,1,10%) in Anhydrous PLO Gel with oral adhesive agent
- Carbamazepine, Capsaicin, Gabapentin, Ketamine, Ketoprofen, Lidocaine HCL, Pregabalin (4,0.012,4,4,4,1,10%) in Anhydrous PLO Gel with oral adhesive agent

MicroClick Pen 0.05ml per click

Sig: Apply 1 click (0.05ml) to affected area 2-4 times daily, as needed.

Other Instructions: _____

Refills: _____ PRN _____

Prescriber Name: _____ Contact Phone: ____-____-____
NPI# _____ Email: _____
DEA# _____
Signature: _____ Date: _____

