

## ULCERS & VIRUSES

Patient Last Name \_\_\_\_\_ First name \_\_\_\_\_ DOB \_\_\_\_/\_\_\_\_/\_\_\_\_

Street Address \_\_\_\_\_ Phone \_\_\_\_-\_\_\_\_-\_\_\_\_

City, State, Zip \_\_\_\_\_

Drug Allergies \_\_\_\_\_

### Intra Oral Ulcers & Viruses

- Magic Mouth antacid, Diphenhydramine 0.25%, Lidocaine HCL 2% (equal amounts) oral rinse suspension
- Diphenhydramine 0.01%, Lidocaine HCL 1%, Misoprostol 0.0024% - Oral rinse suspension
- 2-Deoxy-D-Glucose 0.2%, Diphenhydramine 0.1%, Lidocaine HCL 1% in Nystatin 100,000 units/ml oral rinse suspension
- Acyclovir 2%, 2-Deoxy-D-Glucose 0.2% - oral rinse suspension

**Oral Rinse** \_\_\_\_\_ ml

*Sig: 2.5ml – 5ml orally q4-6h prn. Leave in mouth as long as possible and spit.*

### Anti-Viral Lip Balms

- 2-Deoxy-D-Glucose 0.2%, Lidocaine HCL 0.5%, 5g Lip Balm Tube
- Acyclovir 2%, 2-Deoxy-D-Glucose 0.2%, Lidocaine HCL 0.5%, 5g Lip Balm Tube

*Sig: Apply to affected area 2-4 times daily, as needed. Best to begin before virus erupts.*

### Lichen Plantis Oral

- Tretinoin, Clobetasol Propionate (0.1,0.05%) Oral MucoAdhesive rinse.

**Oral Rinse** \_\_\_\_\_ ml

*Sig: 2.5ml orally Q-8 hr. or PRN. Leave in mouth as long as possible and spit.*

### Oral MucoAdhesive Powder\* (Post Chemotherapy/Radiation)

- Collapsible Puffer Device (with/without APIs) 10g

\*May add anesthetics, steroids, antihistamines, antibiotics, antifungals, misoprostol, aloe, and other natural products.

*Sig: Puff light layer onto affected area and follow with small sip of water. May repeat prn to increase protective layering effect.*

Prescriber Name: \_\_\_\_\_ Contact Phone: \_\_\_\_-\_\_\_\_-\_\_\_\_

**NPI#** \_\_\_\_\_ **Email:** \_\_\_\_\_

**DEA#** \_\_\_\_\_

Signature: \_\_\_\_\_ Date: \_\_\_\_\_

