

ULCERS & VIRUSES

Patient Last Name _____ First name _____ DOB ____/____/____

Street Address _____ Phone ____-____-____

City, State, Zip _____

Drug Allergies _____

Intra Oral Ulcers & Viruses

- Magic Mouth antacid, Diphenhydramine 0.25%, Lidocaine HCL 2% (equal amounts) oral rinse suspension
- Diphenhydramine 0.01%, Lidocaine HCL 1%, Misoprostol 0.0024% - Oral rinse suspension
- 2-Deoxy-D-Glucose 0.2%, Diphenhydramine 0.1%, Lidocaine HCL 1% in Nystatin 100,000 units/ml oral rinse suspension
- Acyclovir 2%, 2-Deoxy-D-Glucose 0.2% - oral rinse suspension

Oral Rinse _____ ml

Sig: 2.5ml – 5ml orally q4-6h prn. Leave in mouth as long as possible and spit.

Anti-Viral Lip Balms

- 2-Deoxy-D-Glucose 0.2%, Lidocaine HCL 0.5%, 5g Lip Balm Tube
- Acyclovir 2%, 2-Deoxy-D-Glucose 0.2%, Lidocaine HCL 0.5%, 5g Lip Balm Tube

Sig: Apply to affected area 2-4 times daily, as needed. Best to begin before virus erupts.

Lichen Plantis Oral

- Tretinoin, Clobetasol Propionate (0.1,0.05%) Oral MucoAdhesive rinse.

Oral Rinse _____ ml

Sig: 2.5ml orally Q-8 hr. or PRN. Leave in mouth as long as possible and spit.

Oral MucoAdhesive Powder* (Post Chemotherapy/Radiation)

- Collapsible Puffer Device (with/without APIs) 10g

*May add anesthetics, steroids, antihistamines, antibiotics, antifungals, misoprostol, aloe, and other natural products.

Sig: Puff light layer onto affected area and follow with small sip of water. May repeat prn to increase protective layering effect.

Prescriber Name: _____ Contact Phone: ____-____-____

NPI# _____ **Email:** _____

DEA# _____

Signature: _____ Date: _____

