

NUTRITION
POULTRY

MITE TONIC
COMPLEMENTARY FEED FOR LAYERS

- Liquid product to administer to poultry through drinking water
- Promotes the natural resistance of the hens
- Harmless for the animals and won't influence the production results negatively
- Suitable for organic farming

Mite ToniC contributes to support the natural resistance of layers. It stimulates layer's natural defenses to create a natural toxin in the blood. This toxin is therefore an immune response.

Mite ToniC is a liquid feed additive to be administrated in drinking water. Safe to use and can be used at any time. Intake guaranteed thanks to the application in drinking water. Easy to apply.

PRODUCT INFORMATION

Composition per L: Magnesium chloride, Sodium chloride, Vitamin C, Plant extracts (Origanum vulgare, Thymus vulgaris, Inula helenium, Litsea cubéba).

RECOMMENDATIONS FOR USE

MiteTonic must be administered in drinking water at the dosage of 0,5 L for 1000 L of water, at the rate of 200 ml per hen.

Protocol 1. Continuous application from flock introduction:

- 1 day per week, during 6 weeks. Renew this treatment every month for 2 consecutive days.

Protocol 2. Intermediate application in case of deficiencies:

- For 4 consecutive days plus 2 days three weeks later and 2 days four weeks later. Renew this treatment every 15 to 20 weeks for 4 consecutive days.

STANDARD MITE TONIC TREATMENT PROTOCOL:

An effective protocol should always be farm-specific. Ask our advice in case of increased infection pressure.

STORAGE/DISPOSAL:

Store in a cool and dry place, between 5°C and 25°C, away from direct light and heat sources. Mite Tonic should not be stored near electrical devices.

LODI UK LIMITED

