

GREENHOUSE & SOIL FEED CHART (EXPANDED)



FEED PROGRAM															
All amounts per 1 gallon. Mix in order listed.	Clone		Vegetative				Flower								
	Pre Soak	Feed	W 1	W 2	W 3	W 4	W 1	W 2	w 3	W 4	W 5	W 6	W 7	W 8	W 9
K-Silicate****		mL EC	1.5mL .1 EC	4mL .2 EC	4mL .2 EC	4mL .2 EC	4mL .2 EC	4mL .2 EC	4mL .2 EC	4mL .2 EC	4mL .2 EC	4mL .2 EC	4mL .2 EC		
Base			24mL 1.3 EC	9.5mL .5 EC											
Grow			28mL 1.2 EC	28mL 1.2 EC	28mL 1.2 EC	28mL 1.2 EC									
Bloom		mL EC					28mL 1.2 EC	14mL .8 EC							
Amino			4mL	6.5mL	6.5mL	6.5mL	6.5mL	5.5mL	5.5mL	4mL	4mL	2.5mL	2.5mL		
PK 0-50-30									2.5mL .1 EC	5.5mL .2 EC	6.5mL .2C	6.5mL .2C			
Potassium										4mL .5 EC	5.5mL .7 EC	4mL .5 EC	2.5mL .3 EC	2.5mL .3 EC	2.5mL .3 EC
Cleaner	1.5	imL	2.5-6.5mL	2.5-6.5mL	2.5-6.5mL	4-10mL	4-10mL	2.5-6.5mL	2.5-6.5mL	2.5-6.5mL	2.5-6.5mL	2.5-6.5mL	2.5-6.5mL	6.5-13.5mL	13.5mL

FOLIAR PROGRAM														
Application Intervals*	Mothers	Vegetative				Flower								
	1 X W	W 1	W 2	W 3	W 4	W 1	W 2	W 3	W 4	W 5	W 6	W 7	W 8	W 9
Fulltek	0.15g	0.15g	0.15g	0.15g	0.15g	0.15g	0.15g							
Kelp	0.1g	0.1g	0.1g	0.1g	0.1g	0.1g	0.1g							
Yucca	0.7g	0.7g	0.7g	0.7g	0.7g	0.7g	0.7g							

Use KALIX Ph Up and Ph Down Super concentrate as needed for best results.

THE FOLLOWING INSTRUCTIONS ARE MEANT ONLY AS A GUIDELINE. ADJUST DOSE RATES DEPENDING ON CULTIVAR AND LEAF TISSUE LAB RESULTS.

* Foliar mother plants three days before taking cuttings.

www.kalixcpn.com Phone: 541.646.6042 sales@kalixcpn.com

KALIX Dosage Guidelines (25lb Bags)

MAKING A STOCK CONCENTRATE

A concentrate should never be fed directly to your plants.

The concentrate mix is meant for mixing into a batch resevoir.

Lbs / Gal	Grams / Gal
1.5	681

LDs / Gal	Glains / Gai
1.5	681

Base Nutrient EC Reference Table

Target Ref EC	Base	Grow / Bloom
0.5	4mL	6.5mL
1.0	8.4mL	13.9mL
1.5	12mL	20mL
2.0	16.5mL	27.8mL
2.5	20mL	33.5mL
2.7	22.5mL	37.2mL
3.0	24mL	40.1mL
3.5	28mL	46.8mL
4.0	32mL	53.5mL

Nutrient Preparation Steps:


1) Acquire an appropriately sized mixing container (20 gallon or larger recommended. One 25lb bag of product will make approximately 15 gallons of stock concentrate).

2) Fill stock concentrate mixing tank with water of total desired liquid concentrate being made. One 25lb bag makes 15 gallons of concentrate

3) Slowly pour weighed product into water while agitating then continue mixing thoroughly.

4) Continue to mix thoroughly until all visible product goes into solution.

5) Store in a dark area, preferably at room temperature.



If unsure whether or not the mix is in solution or suspension, always perform a glass jar test by taking a sample of the mix with a glass jar when it's believed to be in solution and let set 24-48 hrs. If the product falls to the bottom the mix is NOT

- * All values calculated using Reverse Osmosis water. Agitate center of container to avoid clumping.
- ** Not responsible for clogged filters, emitters, or irrigation lines due to improper mixing.
- *** For better results use a transfer pump with a filter to a new tank and always make sure to clean tanks and lines after every feeding.
- **** Depending on the water source. The water for K-Silicate may need to be pH'd to 11 to go into solution faster.