HAND PHANTOM METACARPAL FRACTURES

Age Category

Adult

Body Region

Extremities

Target Modality

Radiography, CT

Diagnostic Features

Bone fractures



This phantom simulates a right hand with two metacarpal fractures. It covers the entire hand including fingers, metacarpus, carpus and the distal parts of the ulna and radius.

The phantom can be used in radiography and CT (including CBCT) to evaluate and optimize imaging performance and post-processing applications, including Al-enabled applications. It is also suited for training purposes.

The phantom provides a detailed and realistic simulation of soft and bone tissue.





HAND PHANTOM METACARPAL FRACTURES



Specifications

Size Approx. 110 x 200 x 69 mm

Weight Approx. 450 g

Base material Cellulose-polymer composite


Optimal 120 kVp (cf page 3)

tube voltage - adaptable upon request -

Diagnostic features

Realistic simulation of bones and soft tissue.

• Fractures of the 3rd and 4th metacarpal



For more information visit www.phantomx.de


HAND PHANTOM METACARPAL FRACTURES


General indications

- The phantom is made of a cellulose-polymer composite material with properties similar to hardwood. If handled carefully, it will last a long time.
- The phantom is coated with a protective layer. If the protective layer is undamaged, the phantom can be cleaned using a damp cloth (water or mild detergent).
- Protect from direct sunlight.
- Maintain a storage temperature of 10 °C to 30 °C. If the phantom is exposed to temperatures below -10 °C or above 45 °C, it can be severely damaged.
- The phantom is not equipped for dose measurements with dosimeters and it is not suited for material characterization with dual energy CT.
- The phantom is not certified as medical device.

Attenuation properties Soft Tissue


Linear attenuation coefficients [cm-1] (calculated)






Bone Tissue

Linear attenuation coefficients [cm-1] (calculated)





Tissue Reference: Woodard HQ, White DR. The composition of body tissues. Br J Radiol. 1986