



## **BBL Hair Removal aftercare**

Patient response can vary after a hair reduction treatment. Erythema (redness) and edema (swelling) around the hair follicles can happen immediately after the treatment and typically resolves within 24-48 hours. A sunburn sensation in the area treated is also normal and expected.

The treated area must be treated with care. **BE GENTLE!** Do not scratch or pick at your skin.

- Until sensitivity has completely subsided, avoid all of the following:
  - Sun exposure
  - Applying make-up over the treated areas
  - Use of scented lotions or soaps, deodorants, exfoliant creams (Retin-A, AHA's or BHA's), acne creams or gels, loofa sponges and aggressive scrubbing
  - Hot or cold water - wash with tepid water
  - Shaving
  - Swimming pools and spas with multiple chemicals/chlorine
  - Activities that cause excessive perspiration
- Strictly avoid any sun exposure to the treated area for a minimum of 7 days after the procedure for laser and 14 days after BBL. The treated area is more prone to sunburn and pigmentation change. Keep the area covered and use a sun block with SPF 30+.
- Delayed blistering secondary to sun exposure has been noted up to 72 hours post treatment.
- A cold compress or an ice pack can be used to provide comfort if the treated area is especially warm. This is typically only needed within the first 12 hours after the treatment.
- Vitamin E or Aloe Vera applied to the treatment area may provide a soothing effect.
- Post treatment discomfort is typically minimal but if the area is very uncomfortable, oral pain reliever, i.e. Extra Strength Tylenol, may be used.
- In the unusual case of crusting of the skin in the treated area, apply an antibiotic ointment twice a day to the affected areas. Do not pick at these areas, as this may result in infection or scarring.
- If the skin is broken or a blister appears, apply an antibiotic ointment and contact the

clinic immediately. Keep the affected area moist and avoid direct sunlight.

- In rare cases, hives have been reported after laser hair reduction treatments. If you experience an irritated raised rash after treatment, please contact our office. Benadryl may be taken to help relieve the itchiness.
- Do not use any other hair removal methods or products on the treated area during the course of your laser treatments (tweezing, waxing and depilatories), as it will prevent you from achieving your best results.

Hair may take **up to 4 weeks**, or in some cases, as little as a few days, to fall out following treatment. Do not be alarmed if hairs appear to be “growing” during this time.

- Subsequent treatments are based upon your clinician’s recommendation and are typically 4-8 weeks apart. The number of treatments required will depend upon the body location, the amount of hair present, the character of the hair and your skin type.

**On-Call medical team can be reached anytime at 403-763-3464. For non-emergency calls, please call the clinic at 403-475-4025.**