

EN

WARNING: THIS PRODUCT IS NOT RECOMMENDED FOR USE BY CHILDREN AND NON-SMOKERS.

Please read this entire leaflet carefully before you start using this product. It contains important information. Keep this leaflet; you may need it again.

DISPOSABLE DEVICE, DO NOT REFILL OR ATTEMPT TO RECHARGE.

PLEASE DISPOSE OF THE DEVICE IF YOU GET A DRY BURN HIT.



1. Avoid taking eLiquid into your mouth or ingesting it.
2. Please do not force eLiquid out of the device.
3. Please do not drop, throw or mishandle this product as the outer shell or inner components may become damaged.
4. In order to prolong the operation of your product, please keep it away from excessive temperatures and humid environments.
5. Please keep this product away from children and pregnant women.

STORAGE CONDITIONS

Store the product at an ambient temperature (between 18-30°C), out of direct sunlight.

Do NOT refrigerate or freeze or store the product above 30°C.

CONTRA-INDICATIONS

Due to propylene glycol:

Adverse effects may occur in patients treated with disulfiram or metronidazole.
Individuals with a history of atopy, allergic rhinitis (hay fever) or childhood eczema may be susceptible to developing allergic reactions.
Lactic acidosis may occur when propylene glycol is consumed in large quantities by children, pregnant women, and patients with hepatic or renal failure.

Due to glycerol:

Chances of dehydration are increased in elderly or already dehydrated patients.
Caution must be observed in diabetic patients as when metabolised, glycerol may cause slight hyperglycaemia and glycosuria.

SPECIFIC RISK GROUPS

This product is not recommended for use by children, and pregnant or lactating women.

POSSIBLE ADVERSE EFFECTS

New users of electronic cigarettes may experience dry throat, dry cough, mild pain or irritation of the mouth or respiratory tract. If the symptoms become severe or last for more than 48 hours, please see a general practitioner and report the incident to the distributor/manufacturer listed below and on the product box.

Possible nicotine withdrawal adverse effects – irritability or aggression, feeling low, anxiety, restlessness, poor concentration, increased appetite or weight gain, urges to smoke (cravings), night time awakening or sleep disturbance, lowering of heart rate.

ADDICTIVENESS

This product contains nicotine which is a highly addictive substance and is thus not recommended for use by non-smokers. No other ingredient of this product is known to be addictive.

TOXICITY

This product contains nicotine which may produce toxic effects if consumed orally, especially in infants and children. Please DO NOT drink this product. In case it is consumed, please take this leaflet and immediately call a POISON CENTRE or general practitioner, reference **H301**. In case of contact with skin, rinse immediately with cold running water for at least 15 minutes.

REPORTING ADVERSE EFFECTS

If you experience any side effects as a result of using this product, please help us by informing us about it so that we can continue to improve the safety of our products.

INFORMATION OF THE PRODUCER

Diamond Mist

128 Cannon Workshops, Cannon Drive

LONDON

E14 4AS

info@diamondmist.com

Tel : +44(0)2081249988