

WildBrew Sour Pitch[™] is a high-performance, high-purity lactic acid bacteria specifically selected for its ability to produce a wide range of sour beer styles.

WildBrew Sour Pitch[™] produces a clean and balanced citrus flavor profile typical of both traditional and modern sour beer styles. When inoculated at optimal temperature and the right conditions, it is a powerful, safe and easy way to handle bacteria for various beer souring techniques, such as a typical kettle souring process. Besides providing an outstanding performance, WildBrew Sour Pitch[™] is capable of delivering consistent results for brewers.

Styles brewed with this bacteria include, but are not limited to, Berliner Weisse, Gose, lambic, American Wild, and Sour IPA.

MICROBIOLOGICAL PROPERTIES

Classified as Lactiplantibacillus plantarum, a facultative hetero-fermentative strain.

VIABLE BACTERIA	> 10 ¹¹ CFU/g	SALMONELLA	Absent in 25g
DRY MATTER	> 92%	ACETIC BACTERIA	< 104 CFU/g
COLIFORM	< 10 ² CFU/g	MOULDS	< 10 ³ CFU/g
E. COLI	Absent in 1g	YEAST	< 10 ³ CFU/g
S. AUREUS	Absent in 1g		


In line with Lallemand's strict quality control and assurance procedures, every single batch of WildBrew Sour Pitch[™] is released to the market only after passing a rigorous series of tests, guaranteeing the highest standards of purity, quality and performance.

BREWING PROPERTIES

In Lallemand's Standard Wort conditions, WildBrew Sour Pitch™ bacteria exhibits:

- Fast pH drop that can be completed within 1-2 days.
- High lactic acid and low acetic acid production.
- Aroma and flavor is citrus and tangy with a hint of fruit.
- The optimal temperature range for WildBrew Sour Pitch[™] when producing sour beer styles is between 30 40°C (86 104°F).
- Souring rate, fermentation time and pH drop are dependent on inoculation density, bacteria handling, fermentation temperature and nutritional quality of the wort.

🛞 FLAVOR & AROMA



QUICK FACTS

BEER STYLES Sours

акома Strong citrus, tangy, sour

ph range 3.2 - 3.5

temperature range 30 - 40°C (86 - 104°F)

HOP TOLERANCE Alpha acid: 4 ppm (IBU) Beta acid: 8 ppm

INOCULATION RATE 10g/hL

TECH Data Sheet

BACTERIA

WILDBREW SOUR PITCH™





TECHNICAL DATA SHEET

WILDBREW SOUR PITCH'

INSTRUCTIONS

Open the sachet and add WildBrew Sour Pitch[™] directly to unhopped wort at a rate of 10g/hL (1g/10L), ensuring temperature is 30 - 40°C (86 - 104°F).

For better distribution rehydrate WildBrew Sour Pitch[™] in a mix of unhopped wort and water (50/50) 10 times the weight of inoculation at 30°C +/- 5°C (86°F +/- 8°F).

Monitor pH drop and test regularly.



inoculation 10g/hL **temperature** 30 - 40°C (86 - 104°F)



Pour WildBrew Sour Pitch™ into unhopped wort

PACKAGING & STORAGE

рH

> 3.4

WildBrew Sour Pitch[™] is currently available in 250g packs (for 25hL / 660 US gal) and 10g packs (for 1hL.)

This product is stable and tolerant to different storage conditions in its original packaging:

- Shelf life of 36 months when stored at -18°C (0°F)

- Shelf life of 18 months when stored at 4°C (40°F)

- Tolerant to 3 week exposure to ambient temperatures (<25°C / 77°F) with no significant loss of viability

While it is recommended to use the entire sachet of WildBrew Sour Pitch[™] after opening, extended storage is possible under specific conditions. Please contact us with any questions you may have.



BREWERS CORNER

For more information on our yeasts including:

- Technical Documents
- Best Practices Documents
- Recipes
 Ritch Det
- Pitch Rate Calculator and other brewing tools
- Scan this QR code to visit the Brewers Corner on our website.

CONTACT US

If you have questions, do not hesitate to contact us at **brewing@lallemand.com**. We have a team of technical representatives happy to help and guide you in your fermentation journey.

www.lallemandbrewing.com brewing@lallemand.com

LALLEMAND BREWING

